



UNIVERSITI PUTRA MALAYSIA

FLOWER ABSCISSION OF CHILLI (*CAPSICUM ANNUUM* L. VAR. KULAI) UNDER HUMID TROPICAL CONDITIONS AND THE USE OF PLANT GROWTH REGULATORS TO IMPROVE FRUIT SET AND YIELD

INUWA SHEHU USMAN

FP 1997 17

FLOWER ABSCISSION OF CHILLI (*CAPSICUM ANNUUM* L. VAR. KULAI) UNDER HUMID TROPICAL CONDITIONS AND THE USE OF PLANT GROWTH REGULATORS TO IMPROVE FRUIT SET AND YIELD.

By

INUWA SHEHU USMAN

**Dissertation Submitted in Fullfilment of the Requirements for the Degree
of Doctor of Philosophy in the Faculty of Agriculture,
Universiti Putra Malaysia.**

June 1997



This work is dedicated to 'El-Nizamiah' - a school of thought established in remembrance of Shehu Usman Danfodio and his Jama'a.



ACKNOWLEDGEMENTS

I wish to express my sincere appreciation to Dr. Awang Soh Mamat, Chairman of my supervisory committee for his keen interest, invaluable assistance and guidance throughout the study. My profound gratitude goes to Dr. Siti Aishah Hassan, Dr. Anuar Abd. Rahim, Dr. Mohd Fauzi Ramlan and Hj. Ahmad Abd. Wahab members of my supervisory committee, for their efforts and contributions in making the program successful.

Gratitude is also extended to Professor Sharifuddin Abd. Hamid, who had keen interest on the program and offered valuable suggestions whenever approached. The study accorded me opportunity to contact personalities within and outside UPM, whose co-operation and assistance contributed to the success of program. In this regard I wish to send a word of thanks to the following: En. Jaafar Husin and Hj. Abdul Malek Yaacob, En. Abd. Karim Latif and Hj. Moh Esa Saiman of the Basic Sciences Laboratory, Mardi; En. Zainal Kassim of the Chemistry Department, UPM; Pn Latifah Md. Nor and her assistant En. Abd. Rahman of the Food Technology Centre, Mardi; Pn. Amina Jusof, Cik Azilah Ab. Jalil and Mr. L. Y. Ho of the Electron Microscopy unit, UPM; and En. Mohd. Yunus Abd. Wahab of the Hydroponic Unit, UPM. The same appreciation is extended to staff of the Agronomy Department who have helped in all ways possible.



I am indebted to the Kano State Polytechnic for granting me a study leave and to the Malaysian Technical Co-operation Program for offering me a scholarship. The companionship of friends like Omar el Sanussi (Libya), Mohammad Abdel Malek (Libya), Ahmad Ganfoud (Libya), Nasir Umar Tsafe (Nigeria), Khairul Naim Hj. Adham (Malaysia), Syed Mohammad Syed Ibrahim (Malaysia), and Saad Osman Abdalla (Sudan), which served as a motivation throughout the program is also acknowledged.



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
LIST OF TABLES	ix
LIST OF FIGURES.....	xi
LIST OF PLATES.....	xiii
ABSTRACT.....	xiv
ABSTRAK.....	xvii
CHAPTER	
I	
INTRODUCTION.....	1
Role of Plant Growth Regulators in Growth and Development.....	1
Role of Growth Regulators in Abscission.....	4
Chilli Crop.....	6
Economic and Nutritional Importance	6
Botany of the Crop	8
Physiological Effects of Temperature	10
Objectives of the Study	13
II	
LITERATURE REVIEW.....	16
Effect of Heat Stress on Reproductive Growth of Chilli.	16
Mechanism of Abscission in Higher Plants	19
Hormonal Basis of Abscission of Chilli Reproductive Structures.....	25
Endogenous Hormone Analysis in Abscissing Chilli Flowers	29
Determination of Ethylene Production Rate and ACC Accumulation in Chilli Flowers.....	26
Analysis of Indole-3-acetic Acid and Abscisic Acid in Chilli Flowers	32
Role of Plant Growth Regulators in Fruit Set and Yield of Chilli	34
Auxins in Flowering and Fruit Development.....	34
Role of Gibberellins in the Induction of Flowering.....	36
Paclobutrazol (PP333) Increases Plant's Resistance to Stress	38
Triacantanol a Natural Constituent of Plants Increases the Growth and Yield of Crops	39
III	
GENERAL MATERIALS AND METHODS	42
Location of the Experiments.	42
Variety.....	43



	Nursery Operations	44
	Cultural Operations	44
	Assessment of Parameters.....	45
	Statistical Analysis.....	46
IV	FLOWER ABSCISSION OF CHILLI UNDER HIGH TEMPERATURE AND HUMID CONDITIONS.....	48
	Introduction.....	48
	Materials and Method	50
	Scanning Electron Microscopy of the Abscission Layer in Chilli Flowers	51
	Energy Dispersive X-ray Analysis (EDX).....	52
	Acetocarmin Staining of Pollen Grains.....	53
	Fluorescence Microscopy of Pollen Tube Behavior in Chilli Flowers.....	54
	Seed Production and Viability Test.....	55
	Results.....	55
	Trends in Abscission of Chilli Flowers under Heat Stress	55
	Electron Microscopy of Abscission Layer in Chilli Flowers.....	59
	X-ray Microanalysis of Calcium Changes in the Abscission Zone.....	62
	Pollen Stainability.....	62
	Pollen Tube Behaviour of Chilli under Heat Stress.....	67
	Seed Production and Germination Test.....	68
	Discussion	68
	Abscission Trends of Chilli Flowers under Stress Conditions.....	68
	Abscission Layer Development in Chilli Flowers	70
	Pollen - Pistil Interaction in Chilli Flowers under Stress Conditions.....	72
	Seed Production and Viability	74
V	HORMONE CHANGES IN THE ABSCISSION OF CHILLI FLOWERS.....	75
	Introduction.....	75
	Materials and Method	76
	Ethylene and ACC Analysis.....	77
	Analysis of Indole-3-acetic Acid and Abscissic Acid in Chilli flowers.....	81
	Results and Discussion.....	87
	Changes in Ethylene and 1-Aminocyclopropane-1-Carboxylic Acid (ACC) in Chilli Flowers in Relation to Abscission.....	87



	Analysis of IAA and ABA in Chilli flowers.....	90
	HPLC Analysis of IAA and ABA.....	90
	GC-MS Analysis of IAA and ABA.....	91
	Changes in IAA Levels in Relation to Abscission of Flowers.....	98
	Changes in ABA Levels in Relation to Abscission of Flowers.....	100
VI	USE OF PLANT GROWTH REGULATORS TO IMPROVE FRUIT SET AND YIELD OF CHILLI I - EFFECT ON FRUIT SET AND YIELD	103
	Introduction.....	103
	Materials and Method	104
	Combined Analysis of Results from the Three Growing Conditions.....	106
	Results.....	107
	Total Number of Flowers.....	107
	Total Abscission of Flowers.....	108
	Total Fruit Count.....	112
	Fruit Set of Chilli	113
	Fruit Yield.....	114
	Fruit Length of Chilli.....	118
	Fruit Diameter.....	120
	Manova Test Criteria and Exact-F Statistics on Effect of Growing Condition on Measured Parameters	122
	Discussion	124
	Effect of PGRs on Fruit Set and Yield of Chilli	124
VII	USE OF PLANT GROWTH REGULATORS TO IMPROVE FRUIT SET AND YIELD OF CHILLI II - THE EFFECT OF TIME AND FREQUENCY OF APPLICATION ON THE GROWTH AND YIELD OF CHILLI.....	130
	Introduction.....	130
	Materials and Method	131
	Capsaicin Determination.....	133
	Determination of Ascorbic Acid in Chilli Fruit.....	136
	Results.....	139
	Plant Height.....	139
	Number of Days to Maturity	142
	Number of Fruits per Plant.....	144
	Fruit Yield.....	145
	Fruit Size.....	151
	Ascorbic Acid Content.....	154
	Capsaicinoid Content.....	154



Discussion	156
The Effect of GA ₄₊₇ Application at Different Stages of Plant Growth.....	156
Effect of PP333 Application at Different Stages of Plant Growth.....	162
Effect of PGR Application on Capsaicinoid Content of Chilli.....	163
VIII GENERAL DISCUSSION AND CONCLUSIONS.....	166
Abscission Tendency of Chilli Flowers under Heat Stress.....	166
Use of Plant Growth Regulators to Improve Fruit Set and Yield of Chilli.....	170
Recommendations for Future Research	173
REFERENCES.....	176
APPENDIX A.....	194
Climatological Data of the Study Area	194
APPENDIX B	196
Sample of Output Generated Using the SAS Repeated Measurement. Analysis Variance Procedure.....	196
APPENDIX C	201
GC-MS Total Ion Mass Chromatogram of Capsaicinoids.....	201
Vita	202



LIST OF TABLES

Table		Page
1	Abscission parameters of chilli as influenced by different microclimates.....	56
2	Pollen stainability of chilli as influenced by different microclimates.....	65
3	Scores of pollen tube growth of chilli grown under glasshouse and open conditions as viewed under the fluorescence microscope.....	67
4	Viability of chilli seeds as influenced by different microclimates.....	68
5	Ethylene production rate and ACC accumulation in chilli flowers relative to changes in abscission.....	88
6	Concentrations of IAA, c-ABA, t-ABA and total ABA in chilli flowers relative to changes in abscission.....	100
7	Total count of chilli flowers as influenced by PGRs under protective structure and field conditions.....	109
8	Total abscission of chilli flowers as influenced by PGRs under protective structure and field conditions.....	111
9	Total fruit count of chilli as influenced by PGRs under protective structure and field conditions.....	123
10	Percentage fruit set of chilli as influenced by PGRs under protective structure and field conditions.....	115
11	Fresh fruit yield of chilli (g/plant) as influenced by PGRs under protective structure and field conditions.....	116
12	Fruit length (cm) of chilli as influenced by PGRs under protective structure and field conditions.....	118
13	Fruit diameter (cm) of chilli as influenced by PGRs under protective structure and field conditions.....	121



15	Plant height(cm) of chilli at various stages of plant growth as influenced by PGR application	141
16	Number of days to maturity of chilli as influenced by PGR application.....	143
17	Number of fruit per plant of chilli as influenced by PGR application.....	146
18	Fresh fruit yield of chilli (g/plant) as influenced by PGR application.....	148
19	Fruit size of chilli as influenced by PGR application.	153
20	Ascorbic acid content of chilli as influenced by PGR application.....	155
21	Capsaicinoid content of chilli as influenced by PGR application.....	158



LIST OF FIGURES

Figure	Page
1 Mean weekly rate of flower opening and abscission as influenced by different microclimates	58
2 Time course of abscission of chilli flowers in the glasshouse as recorded from the date of tagging (DAT)	60
3 Energy dispersive X - ray spectra profile of the abscission zone.....	63
4 Schematic presentation of sample preparation method used in this study for HPLC and GCMS analysis of IAA and ABA.....	85
5 Ethylene production rate and ACC accumulation in chilli flowers at different stages of development relative to abscission.....	89
6 Scatter plot of mean ethylene production and ACC accumulation in chilli flowers.....	89
7 HPLC chromatogram of (a) standard and (b) sample showing the separation of IAA	93
8 HPLC chromatograms of (a) standard and (b)sample showing the separation of IAA, c-ABA and t- ABA.....	94
9(a) Total ion chromatogram and mass spectral plot of IAA components in standard IAA solution.....	95
9(b) Total ion chromatogram and mass spectral components of ABA in Standard ABA solution	96
9(c) Total ion chromatogram and mass spectral components of IAA and ABA in a typical sample run.....	97
10 Concentrations of IAA and ABA in chilli flowers relative to changes in abscission	101
11 Mean total flower count of chilli as influenced by PGR application under glasshouse and open conditions	110
12 Mean fresh fruit yield of chilli as influenced by PGR application under the shade house and open conditions	117



13	Mean fruit length (CM) of chilli as influenced by PGR application under shade house condition.	119
14	The effect of PGR application on the flower count of chilli averaged over the growing conditions	125
15	The effect of PGR application on fresh fruit yield of chilli averaged over the growing conditions	125
16	Box - Whiskers plot of mean fruit length of chilli as influenced by PGR application under glasshouse and field conditions.....	152
17	HPLC chromatograms showing the separation of capsaicinoids.....	157
18	Diurnal fluctuation of temperature of the study area	194
19	Diurnal fluctuation of relative humidity of the study area.....	195



LIST OF PLATES

Plate		Page
1	Scanning electron micrographs of the abscission zone at various stages of flower development relative to abscission.....	61
2	Stainability of grains from chilli flowers formed under varying microclimates.....	64
3	Pollen tube behaviour in chilli flowers under different micro climates as viewed under fluorescence microscope.....	66
4	Modification of Wien et al. (1989) method of ethylene determination.	80
5	Modification of Wien et al. (1989) method of ACC determination.....	81
6	Photographs showing the depressing effects of PP333 on plant growth .	128
7	Fruit ripening of chilli at early harvest as influenced by PGR application in the glasshouse.	160
8	Fruit ripening of chilli at early harvest as influenced by PGR application in the field.	161



Abstract of Dissertation submitted to the senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Doctor of Philosophy.

FLOWER ABSCISSION OF CHILLI (*CAPSICUM ANNUUM* L. VAR. KULAI) FLOWERS UNDER HUMID TROPICAL CONDITIONS AND THE USE OF PLANT GROWTH REGULATORS TO IMPROVE FRUIT SET AND YIELD.

BY

USMAN INUWA SHEHU

JUNE 1997

Chairman: Awang Soh Mamat, Ph. D.

Faculty: Agriculture

A series of experiments were carried out under glasshouse and field conditions, at the Universiti Pertanian Malaysia (UPM), Serdang (latitude 30°2'N and longitude 101°42'E), during the period January 1994 to December 1996. The main objective of the research is to study the nature of abscission of chilli flowers and to use plant growth regulators (PGRs) to improve fruit set and yield of chilli under heat stressed tropical conditions. Most of the flowers fall 3 days after anthesis. This suggest that flower abscission under heat stress is related to the ontogenic changes taking place in the flowers at this stage (pollination and fertilization). Pollen viability and pollen tube growth were not impaired by the stress. Also seed production and viability were similar under both glasshouse and open conditions.



Electron microscopic observations revealed an abscission zone between the peduncle and the twig. The first morphological evidence of abscission by way of cell separation became visible at anthesis. Energy dispersive X - ray scan of this zone showed higher accumulation of Ca^{2+} relative to the surrounding areas. The visibility of cell separation at anthesis indicated that the process of abscission begun prior to pollination and fertilisation.

Endogenous hormone analysis in abscising chilli flowers showed that peak production of ethylene and its precursor 1- aminocyclopropane-1- carboxylic acid (ACC) was at anthesis. Similarly, highest levels of indole acetic acid (IAA) were detected at this stage. The level of abscissic acid (ABA) increased with advancement of abscission.

On the use of PGRs to improve fruit set and yield of chilli, naphthalene acetic acid (NAA), paclobutrazol (PP333), gibberellic acid (GA_{4+7}), and triacontanol (TRIA) were applied to pot grown chilli plants. The GA_{4+7} treatment proved superior over other PGRs in flower count, fruit count, fruit size and total fruit yield. The yield increase following NAA application has not been consistent. The application of TRIA proved effective only in the open. PP333 application increased fruit set and gave the least recorded abscission. In addition this treatment gave yields that are comparable to other PGRs. GA_{4+7} and PP333 were selected for the next stage of the study.

In the second stage, GA₄₊₇ maintained superiority in plant height, fruit yield, fruit size and ascorbic acid content of fruits. Its application hastened fruit maturity and favoured heavier early harvests. The yield of fruits increased with change in application time of GA₄₊₇ from early flowering (EF) to peak flowering (PF) and then double application (EF+PF). The application of PP333 resulted in delayed maturity and heavier late season harvest. The depressing effect of PP333 on plant height was more pronounced in the glasshouse than in the field. PP333 applied at PF tended to be superior in yield parameters, followed by double application (EF+PF) and then EF treatment. Application of GA₄₊₇ increased the total capsaicinoid content of chilli fruit.



Abstrak Disertasi yang diserahkan kepada Senat Universiti Putra Malaysia sebagai memenuhi sebahagian daripada keperluan untuk ijazah Doktor Falsafah

KECENDERONGAN PELURUHAN BUNGA CILI (*CAPSICUM ANNUUM* L. VAR. KULAI) DALAM KEADAAN TROPIKA YANG LEMBAB DAN PENGGUNAAN PENGAWALATUR PERTUMBUHAN POKOK UNTUK MENINGKATKAN PEMBENTUKAN BUAH DAN HASIL.

OLEH

USMAN INUWA SHEHU

JUNE 1977

Pengerusi: Awang Soh Mamat, Ph.D.

Fakulti: Pertanian

Satu siri ujikaji telah dijalankan di dalam rumah kaca dan di ladang di Universiti Putra Malaysia (UPM), Serdang (garis lintang $30^{\circ}2'U$ and garis bujur $101^{\circ}42'T$), Selangor, Malaysia mulai Januari 1994 hingga Disember 1996. Objektif utama penyelidikan ini ialah untuk mengkaji tabiat peluruhan bunga cili dan penggunaan pengawalatur pertumbuhan pokok untuk meningkatkan pembentukan buah dan hasil cili dalam keadaan tegasan haba tropika. Sebahagian besar daripada bunga-bunga cili luruh dalam masa 3 hari selepas antesis. Ini menunjukkan bahawa peluruhan bunga dalam keadaan tegasan haba berkait dengan perubahan ontogenik yang berlaku dalam bunga pada peringkat ini (contoh: pendebungaan dan persenyawaan). Kebolehhidupan debunga dan



pertumbuhan tiub debunga tidak berubah disebabkan tegasan haba. Pengeluaran dan kebolehhidupan biji juga didapati sama diantara di dalam rumah kaca dan di ladang.

Pemerhatian dengan menggunakan imbasan mikroskop electron (scanning electron microscopy) menunjukkan kewujudan zon peluruhan di pertemuan antara tangkai bunga dan ranting. Bukti morfologi yang pertama yang menunjukkan bahawa peluruhan berlaku melalui pemisahan sel-sel jelas kelihatan semasa antesis. Imbasan penyebaran tenaga sinaran-X (Energy dispersive x-ray) pada zon ini menunjukkan kehadiran Ca^{2+} yang banyak berbanding dengan kawasan keliling. Pemisahan sel-sel yang dilihat semasa antesis menunjukkan bahawa proses peluruhan bermula sebelum pendebungaan dan persenyawaan.

Analisa hormon endogen keatas bunga-bunga cili yang luruh menunjukkan pengeluaran etilena dan prekursornya [1-aminocyclopropane-1-carboxylic acid (ACC)] yang memuncak semasa antesis. Paras asid indola-asetik (IAA) yang tinggi juga dikesan pada peringkat ini. Paras asid absisik (ABA) bertambah dengan penambahan peluruhan.

Mengenai penggunaan pengawalatur pertumbuhan untuk meningkatkan pembentukan buah dan penghasilan cili, pokok cili di dalam polibag dirawat dengan asid asitik naftalena (NAA), paklobutrazol (PP333), asid gibralik



(GA₄₊₇) dan triakontanol (TRIA). GA₄₊₇ didapati lebih baik daripada pengawalatur pertumbuhan yang lain, disegi peningkatan jumlah bunga, jumlah buah, saiz buah dan jumlah hasil. Pertambahan hasil selepas menggunakan NAA didapati tidak tetap. Penggunaan TRIA berkesan hanya di ladang. Penggunaan PP333 meningkatkan pembentukan buah dan memberi jumlah bunga luruh yang paling rendah. Rawatan ini juga memberi hasil yang setara dengan pengawalatur pertumbuhan yang lain. GA₄₊₇ dan PP333 kemudiannya dipilih untuk ujikaji seterusnya.

Dalam peringkat kedua, GA₄₊₇ didapati masih meningkatkan ketinggian pokok, penghasilan buah, saiz buah dan kandungan asid askorbik. Penggunaan GA₄₊₇ mempercepatkan kematangan buah dan lebih banyak buah dipungut awal. Pengeluaran hasil buah cili berubah mengikut masa penggunaan GA₄₊₇. Semburan semasa pembungaan awal (PA) memberi hasil paling rendah. Hasil bertambah apabila semburan dibuat semasa pembungaan puncak (PP). Seterusnya, hasil paling tinggi dipungut daripada pokok yang menerima semburan berganda (PA+PP). Penggunaan PP333 menghasilkan buah yang mempunyai kandungan melewati kematangan dan lebih banyak buah dipungut lewat. Kesan pengurangan ketinggian pokok oleh PP333 lebih jelas di dalam rumah kaca berbanding dengan di ladang. PP333 yang disemur semasa pembungaan puncak (PP) lebih berkemungkinan memberi hasil yang lebih baik. Ini diikuti oleh penyemburan berganda (PA+PP) dan penyemburan

semasa pembungaan awal (PA). Penggunaan GA₄₊₇ meningkatkan jumlah kandungan kapsaisinoid di dalam buah cili.

CHAPTER I

INTRODUCTION

Role of Plant Growth Regulators in Growth and Development

Communications within the plant as in animals is mediated by the action of chemical substances called hormones. A hormone is defined as any organic compound synthesised in one organ in response to a stimulus, and transported to another organ where a response is produced. In recent years, the hormone concept has been questioned by different sources. The chemicals involved do not necessarily move from their site of synthesis. In view of this, and because many aspects of their action are on plant growth they have been called 'growth substances' synonymous to 'growth regulators' (Van Overbeek, 1954).

Plant growth substances regulate the physiological and morphological processes taking place in the plant. This control is achieved through a complex interaction between plant internal factors and the environment. Auxins, gibberellins and cytokinins have their most pronounced effects on growth, flowering, fruit and seed set. Whereas ethylene and abscisic acid are



associated with ripening, senescence and abscission of plant organs. For a normal growing plant that is at equilibrium with its environment, sufficient endogenous levels of the individual hormones are usually available. A deviation from this equilibrium results in changes in the levels of the hormones which is usually accompanied with plant response to this change. For instance endogenous levels of ABA and ethylene increase under water stress and the plants respond to this stimulus by slowed growth, stomatal closure and increase in protein synthesis. This is probably a survival mechanism for the plant to overcome the stress. Similarly, modifications in growth would occur due to high levels of these growth retardants in the case of air pollution, pest damage, flooding, etc.

The growing plant passes through several stages of development that are associated with definite morphological and physiological changes. Similarly, a single organ of the plant may pass through several developmental sub-phases. In the stem for instance, older internodes are completing their development when younger ones are just beginning. This decreasing rate of elongation has been correlated with the increasing ability of the tissues to destroy IAA in Japanese morning glory (Yoneda and Stonier, 1966).

The transition from vegetative to flowering stage is a complex morphological event. The ability to flower is attained when the plant has reached a certain size or age. In some plants flowering occur independent of



the environment (autonomous induction). Other plants require exposure to appropriate environmental conditions. Day length and temperature are the most important environmental factors controlling flowering.

As in the case of growth, there is a hormonal control of flowering. This response occurs in the growing points, not yet ascribed to any particular day length. A specific flowering hormone in the name of "florigen" was postulated in the 1930's by Michael Chailakhyan working in Russia. Many attempts have been made to extract and isolate this hormone but without much success (Zeevart and Boyer, 1987; Zeevart, 1984; and Lang et al., 1977). Thus, despite unequivocal data showing that a transmissible factor regulates flowering, the involved substances are yet to be characterised. Gibberellins play an important role in flowering. Exogenous gibberellin has been found to substitute for photoperiodic induction in some plants (Lang, 1957). In particular, GAs cause flowering when applied to long day plants that grow as rosettes in short days. In these plants the flowering response is accompanied by elongation of the flowering stem.

Senescence also called ageing of the plant, is the stage preceding death or in the case of organs, preceding abscission. Many physiological changes accompany the ageing process. Translocation of food materials away from the senescing organ has been put forward as a basis for senescence. However, studies indicated that hormones are responsible. According to this view a



deficiency of one or more of the growth regulators leads to senescence (Osborne, 1963). Thus, auxin content declines with age of leaf and towards death is too low to measure accurately. This decline in auxin is accompanied by a decline of other substances, perhaps because of a direct effect of auxin on phloem transport. Therefore, older leaves senesce more rapidly in the presence of young auxin-rich leaves above them. Senescence can actually be delayed by treating leaves with growth regulators. Kinetin delays sequential ageing and auxin delays synchronous ageing. Growth retardants may also be used successfully (Kesler et al., 1967) even when cytokinins fail.

ABA greatly accelerates senescence of leaves. Ethylene has been found to play a role in the senescence of oat leaf segments, but ABA appears to be the initiating agent. Whereas ethylene exerts its effect at a later stage (Gepstein and Thimann, 1981). Leaf senescence has been studied extensively and the physiological, anatomical and biochemical changes that take place during this process have been described (Nooden and Leopold, 1988).

Role of Growth Regulators in Abscission

Ethylene appears to be the primary regulator of the abscission process, with auxin acting as a suppressor of the ethylene effect. A model of hormonal control of leaf abscission that explains the process in three distinct phases has

