



UNIVERSITI PUTRA MALAYSIA

**EFFECT OF PHYTATE PHOSPHORUS ON THE UTILISATION OF
RICE BRAN IN BROILER CHICKENS)**

MD ASADUZZAMAN UKIL

FP 1999 19



**EFFECT OF PHYTATE PHOSPHORUS ON THE UTILISATION OF RICE
BRAN IN BROILER CHICKENS**

BY

MD ASADUZZAMAN UKIL

**Thesis Submitted in Fulfilment of the Requirements for the Degree of Doctor
of Philosophy in the Faculty of Agriculture
Universiti Putra Malaysia**

November 1999



Dedication

To the memory of my late parents



ACKNOWLEDGEMENTS

At the outset, I would like to express my most sincere gratitude and deep appreciation to Associate Prof. Dr. Abdul Razak Alimon, chairman of the supervisory committee, for his advice, tutelage, invaluable guidance, encouragement and providing graduate assistantship from the IRPA Project (No. 51297, UPM) throughout the course of study.

I am also indebted to the members of my supervisory committee, Associate Prof. Puan Ramlah Hamid and Dr. Zulkifli Idrus for their constructive suggestions, guidance and critical reviewing of this dissertation.

I gratefully acknowledge the assistance of the Bangladesh Open University authority for allowing me to pursue the study programme smoothly by providing me study leave throughout the period. A very special thanks are due to Professor Dr. M. Aminul Islam, vice-chancellor, Bangladesh Open University who always encouraged me during the course of the study.

The author gratefully acknowledge the BASF Malaysia for providing the enzyme microbial phytase (Natuphos® 5000).

I would like to express my appreciation to Mr. Ibrahim Mohsin, Laboratory in charge, Mr. Saporin, Mr. Bakari, Mr. Chan Tin Wan for helping me during the laboratory work. I want to extend my appreciation to Mr. Islah Uddin, Mr. Shahril



Hassan, Mr. Mazlan Hamzah, Mr. Ponnosamy, during experimentation. I gratefully acknowledge the staff members of the department of animal science for helping me in any means. I also record my appreciation to the staff of graduate school

I want to extend my sincere thanks to all of my post-graduate friends and well wishers including Dr. Rafiqul Islam, Mr. Sheikh Ashraf Ali, Ms Yusnita, who have assisted me by providing different inputs directly or indirectly which contributed to the successful accomplishment of this work.

Last but not least, very special thanks go to my wife Parvin Akther Rumi for her sacrifice, patience, understanding, help and encouragement throughout the study. My Sons, Azizur Rahman Anik and Asif Rahman Antik also deserve appreciation for their co-operation.



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS.....	iii
LIST OF TABLES.....	xii
LIST OF FIGURES.....	xvii
LIST OF ABBREVIATIONS.....	xix
ABSTRACT.....	xxii
ABSTRAK.....	xxiv
 CHAPTER	
I GENERAL INTRODUCTION.....	1
Objectives.....	5
 II LITERATURE REVIEW.....	6
Introduction.....	6
Concept of Rice Bran.....	7
Types/Classification of Rice Bran.....	10
Composition of Rice Bran.....	12
Proximate Components.....	14
Chemical Constituents.....	15
Amino Acid Contents of Rice Bran.....	15
Vitamins.....	16
Minerals.....	17
Fats and Fatty Acid Characteristics of Rice Bran.....	18
Antinutritional Factors in Rice Bran.....	19
Trypsin Inhibitor.....	20
Hemagglutinin.....	20
Phytate Phosphorus.....	21
Rice Bran in Poultry Ration.....	22
Processing of Rice Bran.....	29
Stabilisation.....	30
Dry-Heat Stabilisation.....	30
Added Moisture Heat Stabilisation.....	30
Retained Moisture / Wet heat stabilisation.....	31
Parboiling.....	31
Deoling.....	31
Autoclaving.....	31
Extrusion Cooking.....	32



Effect of Rice Bran on Efficiency of Energy and Protein Utilisation.....	32
Occurrence, Distribution and Structure of Phytate Phosphorus in Foodstuffs	33
Hydrolysis of Phytate Phosphorus and its Bioavailability...	36
Factors Influencing Phytate Phosphorus Utilisation.....	39
Dietary Calcium and Phosphorus Levels.....	39
Dietary Vitamin D ₃ (1,25-Dihydroxycholecalciferol) Levels.....	40
Age of Birds.....	41
Type of Dietary Feed Ingredients	41
Effect of Fibre.....	42
Genotype of Birds.....	42
Feed Processing.....	42
Effect of Phytate on Bioavailability of Other Nutrients.....	44
Effects on the Mineral Bioavailability	44
Effect on Protein (Amino Acid) Availability.....	45
Characteristics, Sources and Nutritional Effects of Phytase.....	47
Use of Phytase in Poultry Ration.....	48
Effect of Feed Processing on Phytase Activity.....	48
Potential of Microbial Phytase to Increase the Availability of Nutrients.....	49
Effects of Microbial Phytase on Phytate Phosphorus Availability.....	49
Effect of Microbial Phytase on the Availability of Cations.....	50
Effects of Phytase on the Availability of Protein and Amino Acids.....	52
Role of Dietary Calcium and Calcium:Phosphorus Ratio on the Efficacy of Microbial Phytase.....	53
Effect of Phytase and Vitamin D ₃ on the Level of Calcium and Phosphorus and Ca:P Ratios.....	54
Effect of Microbial Phytase Supplementation on the Performance of Broiler Chickens.....	55
Effect of Vitamin D ₃ Supplementation on the Utilisation of Phytate Phosphorus.....	57
Effects on Metabolisable Energy (ME).....	57
Effects on Varying Levels of Calcium, Available Phosphorus and Dietary Protein.....	58
Responses of Phytase on Laying Hens	58
Response of Phytase on Phosphorus Excretion.....	59

III	GENERAL MATERIALS AND METHODS.....	61
	Animals.....	61
	Housing.....	61
	Sanitation of the House.....	62
	Rearing and Management of the Chicks up to 21 Days.....	62
	Vaccination.....	63
	Growth Trial.....	63
	Ambient Temperature.....	63
	Digestibility Trial.....	64
	Data Collection.....	65
	Sampling and Chemical Composition of the Feed Ingredients.....	65
	Formulations of the Experimental Diets.....	66
	Mixing of Vitamin Mineral Premix, Minerals and Other Feed Additives.....	66
	Preparation of the Diets.....	66
	Sampling of the Mixed Diet.....	67
	Proximate Analysis.....	67
	Determination of Moisture and Dry Matter.....	67
	Determination of Ash.....	68
	Crude Protein.....	69
	Ether Extract.....	70
	Determination of Crude Fibre.....	71
	Determination of Acid Detergent Fibre.....	72
	Determination of Acid Detergent Lignin.....	73
	Neutral Detergent Fibre.....	74
	Determination of Minerals.....	75
	Wet Ashing.....	75
	Dry Ashing.....	76
	Determination of Phosphorus.....	77
	Determination of Calcium, Zinc, Copper and Manganese....	78
	Determination of Gross Energy.....	78
	Preparation of Samples for Determination of Amino Acid...	80
	Preparation of Glassware.....	80
	Determination of Amino Acid.....	81
	Required Eluents and Reagents.....	81
	Acid Hydrolysis.....	82
	Drying.....	82
	Redrying Procedure.....	82
	Derivatisation Procedure.....	83
	Start up and Operation.....	83
	Standard Preparation.....	84
	Calculation of Amino Acids.....	84



	Determination of Chromic Oxide.....	85
IV	CHEMICAL COMPOSITION OF RICE BRAN AVAILABLE IN MALAYSIA.....	86
	Introduction.....	86
	Materials and Methods.....	87
	Collection of Rice Bran Samples.....	87
	Fractionation of Rice Bran into Different Particles.....	87
	Preparation of Samples for Chemical Analysis	88
	Chemical Analysis.....	88
	Results	88
	Discussion.....	91
V	REPLACEMENT OF CORN WITH RICE BRAN IN BROILER FINISHER DIET.....	95
	Introduction.....	95
	Materials and Methods.....	96
	Experiment 1a.....	96
	Animals Treatments and Experimental Design.....	96
	Feed Formulation.....	97
	Feeding and Management.....	97
	Meat Yield.....	98
	<i>In Vivo</i> Digestibility.....	98
	Collection of Faecal Samples.....	98
	Chemical Analysis.....	100
	Experiment 1b.....	100
	Animals, Treatments and Experimental Design.....	100
	Feed Formulation.....	101
	Feeding and Management.....	101
	Data Collection.....	101
	Statistical Analysis.....	103
	Results.....	103
	Experiment 1a.....	103
	Body Weight.....	103
	Weight Gain	103
	Feed Intake.....	104
	Feed Conversion Ratio.....	104
	Carcass Quality.....	107
	Digestibility.....	109
	Experiment 1b.....	110
	Body Weight.....	110
	Weight Gain.....	110
	Feed Intake.....	110

	Feed Conversion Ratio.....	111
	Discussion.....	112
	Conclusion.....	118
VI	EFFECTS OF SUPPLEMENTAL PHYTASE AND 1,25 DI-HYDROXYCHOLECALCIFEROL ON THE GROWTH PERFORMANCE OF BROILER CHICKENS FED RICE BRAN BASED DIETS.....	119
	Introduction.....	119
	Materials and Methods.....	122
	Animals, Treatments and Experimental Design.....	122
	Lay Out of the Diets.....	123
	Feeding and Management.....	124
	Meat Yield.....	124
	<i>In Vivo</i> Digestibility.....	126
	Collection of Faecal Samples.....	126
	Chemical Analysis.....	126
	Determination of Toe Ash.....	127
	Determination of Tibia Ash.....	127
	Statistical Analysis.....	127
	Curve Fitting.....	128
	Results.....	128
	Body Weight.....	128
	Weight Gain.....	129
	Feed Intake and Feed Conversion Ratio.....	129
	Meat Yield.....	129
	<i>In Vivo</i> Digestibility.....	134
	Intake and Excretion of phosphorus.....	134
	Toe and Tibia Ash.....	140
	Discussion.....	143
	Conclusion.....	148
VII	EFFECTS OF SUPPLEMENTAL PHYTASE AND DIFFERENT LEVELS OF PROTEIN ON THE PERFORMANCE OF BROILER CHICKENS (FINISHER) FED RICE BRAN BASED DIETS.....	149
	Introduction.....	149
	Materials and Methods.....	151
	Animals, Treatments and Experimental Design.....	151
	Feed Formulation.....	151
	Feeding and Management.....	152
	<i>In Vivo</i> Digestibility.....	153
	Collection of Faecal Samples for Total and Indicator	



Method.....	153
Sampling for Ileal Digestibility Digestibility of Amino Acids	154
Determination of Digestibility.....	154
Chemical Analysis.....	155
Apparent Ileal Digestibility.....	155
Determination of Toe Ash.....	155
Determination of Tibia Ash.....	156
Statistical Analysis.....	156
Results.....	156
Body Weight	156
Weight gain.....	157
Feed Intake.....	159
Feed Conversion Ratio.....	160
Toe and Tibia Ash.....	162
<i>In Vivo</i> Digestibility.....	164
Indicator Methods for Determining the Digestibility of DM, CP, Ca and P and Ileal Digestibility of Amino Acids.....	167
Discussion.....	170
Conclusion.....	173

VIII	EFFECT OF CALCIUM TO PHOSPHORUS RATIO ON THE EFFICACY OF PHYTASE IN BROILER CHICKENS (FINISHER) FED RICE BRAN BASED DIETS HAVING TWO LEVELS OF PROTEIN.....	174
	Introduction.....	174
	Materials and Methods.....	176
	Animals Treatments and Experimental Design.....	176
	Feed Formulation.....	176
	Feeding and Management.....	176
	<i>In Vivo</i> Digestibility.....	178
	Collection of Faecal Samples	179
	Sampling for Ileal Digestibility of Amino Acids.....	179
	Chemical Analysis.....	179
	Determination of Toe Ash.....	180
	Determination of Tibia Ash.....	180
	Statistical Analysis.....	180
	Results	181
	Body Weight.....	181
	Weight Gain	181
	Feed Intake.....	182
	Feed Conversion Ratio.....	182
	<i>In Vivo</i> Digestibility	187



Ileal Digestibility of Amino Acids.....	192
Toe and Tibia Ash Contents.....	192
Excretion of Calcium and Phosphorus.....	192
Discussion.....	196
Conclusion.....	199
IX GENERAL DISCUSSION AND CONCLUSION.....	200
Conclusions.....	207
BIBLIOGRAPHY.....	209
APPENDIX	
A Eluents and Reagents for Determination of Amino Acids.....	231
PUBLICATIONS FROM THIS WORK.....	233
VITA.....	234



LIST OF TABLES

Table		Page
1	Types of Discrete Particles Identified in Spanish Commercial Rice Bran.....	10
2	Proximate Analyses and Some Chemical Constituents of Rice Bran (%)......	14
3	Amino Acid Composition of Rice Bran (g/16.g of N).....	16
4	Vitamin Content of Rice Bran.....	17
5	Mineral Content (macro and micro) of Rice Bran.....	18
6	Fatty Acid Composition of Rice Bran (weight % of total acid)...	19
7	Phytate Phosphorus Contents of Some Feed Ingredients Commonly Used in Chicken Feed.....	34
8	Phytase Activity of Some Feed Ingredients.....	38
9	Effect of Microbial Phytase on Body Weight (BW) and Retention of Phosphorus in Broiler Chickens fed Corn-Soybean Meal Diets.....	56
10	Effect of Phytase Supplementation on the Performance of Laying Hens Over the Period from 24 to 54 Weeks of Age.....	60
11	Chemical Composition of Whole Rice Bran and its Fractions.....	89
12	Mineral Contents of Rice Bran and its Fractions.....	90
13	Fractions of Different Particle Size in Whole Rice Bran.....	91
14	Composition of the Diets Using Different Levels of Rice Bran...	99
15	Composition of the Diets Using Different Levels of Rice Bran...	102



16	The Average Body Weight and Weight Gain of Broiler (finisher) Fed Different Levels of Rice Bran from Day 21-42.....	105
17	The Average Feed Intake and FCR of Broiler (finisher) Fed Different Levels of Rice Bran from Day 21-42.....	106
18	Carcass Composition and Meat yield (% of body Weight) of Broiler Chickens fed Different Levels of Rice Bran.....	108
19	Digestibility (%) of Nutrients and Apparent Metabolisable Energy (AME) Contents of Diets.....	109
20	Mean \pm SE of Live Weight, Weight Gain, Feed Intake and FCR of Broiler (finisher) Fed Different Levels of Rice Bran.....	112
21	Composition of the Formulated Rice Bran Based Diets for Broiler (finisher).....	125
22	Mean \pm SE of Body Weight and Weight Gains of Broiler (finisher) fed Rice Bran Based Diet Supplemented with Phytase Natuphos® Vitamin D ₃ and Dicalcium Phosphate.....	131
23	Mean \pm SE of Feed Intake and FCR (weekly) of Broiler (finisher) fed Rice Bran Based Diet Supplemented with Phytase Natuphos® Vitamin D ₃ and Dicalcium Phosphate.....	132
24	Mean \pm SE of Dressing (%) Gizzard, Liver, Heart, Abdominal Fat, Skin and Shank Weight (% of body weight) of Birds fed Low Phosphorus Diet Supplemented with Phytase or 1,25(OH) ₂ D ₃	133
25	Apparent Metabolisable Energy (AME kcal/kg) of diets and Digestibility (%) of Dry Matter (DM), Crude Protein (CP), Calcium and Phosphorus.....	135
26	Intake and Excretion of Phosphorus in Broiler Chicken fed Rice Bran Based Diets Supplemented with Phytase or Vitamin D ₃	138



27	Toe and Tibia Ash Contents of Birds fed Supplemental Phytase and Vitamin D ₃ (1,25(OH) ₂ D ₃).....	140
28	Composition of the Formulated Rice Bran Based Diets for Broiler Chicken (finisher).....	152
29	Body Weights (g) of Broiler Chickens (finisher) fed Rice Bran Based Diets Consisting of Two Levels of Protein Supplemented with Four Levels of Phytase Natuphos®.....	158
30	Weight Gain (g) of Broiler Chicken (finisher) fed Rice Bran Based Diet Consisting of Two Levels of Protein Supplemented with Four Levels of Phytase Natuphos®.....	159
31	Feed Intake (g) of Broiler Chicken (finisher) Fed Rice Bran Based Diet Consisting of Two Levels of Protein Supplemented with Four Levels of Phytase. Natuphos®.....	161
32	Feed Conversion Ratio of Broiler Chicken (finisher) fed Rice Bran Based diet Consisting of Two Levels of Protein Supplemented with Four Levels of Phytase Natuphos®.....	162
33	Effect of Phytase on Toe and Tibia Ash Content of Broiler Chicken (finisher) fed Rice Bran Based Diets Consisting of Two Levels of Protein and Four Levels of Phytase Natuphos®.....	163
34	Digestibility (g) of Dry Matter (DM), Crude Protein (CP) of Broiler Chicken fed Rice Bran Based Diet Consisting of Two Levels of Protein Supplemented with Four Levels of Phytase Natuphos® in Total Collection Method.....	166
35	Digestibility (%) of Calcium and Phosphorus in Broiler Chicken fed Rice Bran Based Diet Consisting of Two Levels of Protein Supplemented with Four Levels of Phytase Natuphos® in Total Collection Method.....	167
36	Digestibility of Dry Matter (DM) and Crude Protein (CP) of Broiler Chicken (finisher) fed Rice Bran Based Diet Consisting of Two Levels of Protein Supplemented with Four Levels of Phytase in Indicator Method.....	168



37	Digestibility of Calcium and Phosphorus of Broiler Chicken (finisher) fed Rice Bran Based Diets Consisting of Two Levels of Protein and Four Levels of Phytase in Indicator Method.....	168
38	Main effects of Protein and Phytase on the Ileal Digestibility (%) of Amino acids.....	169
39	Composition of the Formulated Rice Bran Based Diets for Broiler (finisher).....	177
40	Body Weights (g/bird) of Broiler Chicken (finisher) Fed Rice Bran Based Diet Consisting of Two Levels of Protein and Three Ca:tP Ratios as Influenced by Supplemental Phytase Natuphos®.....	183
41	Weight Gain (g/bird) of Broiler Chicken (finisher) Fed Rice Bran Based Diet Consisting of Two Levels of Protein and Three Ca:tP Ratios as Influenced by Supplemental Phytase Natuphos®.....	184
42	Feed Intake (g/bird) of Broiler Chicken (finisher) Fed Rice Bran Based Diet Consisting of Two Levels of Protein and Three Ca:tP Ratios as Influenced by Supplemental Phytase Natuphos®.....	185
43	Feed Conversion Ratio of Broiler Chicken (finisher) Fed Rice Bran Based Diet Consisting of Two Levels of Protein and Three Ca:tP Ratios as Influenced by Supplemental Phytase Natuphos®.....	186
44	Digestibility (%) of Dry Matter (DM) and Protein (Total Collection Method) of Broiler Chicken (finisher) Fed Rice Bran Based Diet Consisting of Two Levels of Protein and Three Ca:tP Ratios as Influenced by Supplemental Phytase Natuphos®.....	188
45	Digestibility (%) of Calcium and Phosphorus (total collection method) of Broiler Chicken (finisher) Fed Rice Bran Based Diet Consisting of Two Levels of Protein and Three Ca:tP Ratios as Influenced by Supplemental Phytase Natuphos®.....	189



46	Digestibility (%) of Dry Matter (DM) and Crude Protein (CP) of Broiler Chicken (finisher) Fed Rice Bran Based Diet Consisting of Two Levels of Protein and Three Ca:tP Ratios as Influenced by Supplemental Phytase Natuphos® (Indicator method).....	190
47	Digestibility of Calcium and Phosphorus of Broiler Chicken (finisher) Fed Rice Bran Based Diet Consisting of Two Levels of Protein and Three Ca:tP Ratios as Influenced by Supplemental Phytase Natuphos® (Indicator method).....	191
48	Main Effects of Crude Protein (CP) and Calcium to Total Phosphorus Ratio on the Ileal Digestibility (%) of Amino Acids.	193
49	Toe and Tibia Ash Contents of Broiler Chickens (finisher) Fed Rice Bran Based Diet Consisting of Two Levels Protein and Three Ca:tP Ratios as Influenced by Supplemental Phytase Natuphos®.....	194
50	Excretion of Calcium and Phosphorus in Broiler Chickens (finisher) Fed Rice Bran Based Diet Consisting of Two Levels of Protein and Three Ca:tP Ratios as Influenced by Supplemental Phytase Natuphos®.....	195



LIST OF FIGURES

Figure		Page
1	The Different Fractions of Paddy Rice as a Result of Milling....	12
2	Structure of Phytic Acid as Proposed by Anderson (1914).....	35
3	Phytic Acid Chelated at Neutral pH.....	35
4	Feed Intake of Broiler Chickens fed Different Levels of RB.....	107
5	Feed Intake of Broiler Chickens fed Different Levels of RB in the Diet.....	111
6	Digestibility of Calcium of Broiler Chicken Fed Rice Bran Based Diets Supplemented with Phytase.....	136
7	Digestibility of Phosphorus of Broiler Chicken Fed Rice Bran Based Diets Supplemented with Phytase.....	136
8	Digestibility of Calcium of Broiler Chicken Fed Rice Bran Based Diets Supplemented with Vitamin D ₃	137
9	Digestibility of Phosphorus of Broiler Chicken Fed Rice Bran Based Diets Supplemented with Vitamin D ₃	137
10	Phosphorus Excretion of Broiler Chickens fed Rice Bran Based Diets Supplemented with Phytase.....	139
11	Phosphorus Excretion of Broiler Chickens fed Rice Bran Based Diets Supplemented with Vitamin D ₃	139
12	Toe Ash Contents of Broiler Chickens fed Different Levels of Phytase	141
13	Toe Ash Contents of Broiler Chickens fed Different Levels of Vitamin D ₃	141
14	Tibial Ash Contents (%) of Broiler Chicken fed Rice Bran Based Diets at the Finishing Stage Supplemented with	

	Phytase.....	142
15	Ash Contents of Tibia of Broiler Chicken fed Rice Bran Based Diets Supplemented with Vitamin D ₃	142
16	Toe Ash Contents of Broiler Chickens fed Different Levels of Protein in Rice bran Based Diets Supplemented with Microbial Phytase.....	163
17	Tibia Ash Contents of Broiler Chickens fed Different levels of Protein in Rice bran based Diets Supplemented with Microbial Phytase.....	164
18	Digestibility of Calcium of Broiler Chickens given Different Levels of Protein in Rice Bran Based Diets Supplemented with Phytase.....	165
19	Digestibility of Phosphorus of Broiler Chickens given Different Levels of Protein in Rice Bran Based Diets Supplemented with Phytase.....	165



LIST OF ABBREVIATIONS

AA	Amino Acid
AAS	Atomic Absorption Spectrophotometer
ADF	Acid Detergent Fibre
ADL	Acid Detergent Lignin
AME	Apparent Metabolisable Energy
aP	Available Phosphorus
Ca	Calcium
CF	Crude Fibre
CP	Crude Protein
cm	Centimetre
Cr	Chromium
Cu	Copper
DFRB	De-fatted Rice Bran
DM	Dry Matter
d	Day
EE	Ether Extract
FCR	Feed Conversion Ratio
Fe	Iron
FFRB	Full Fatted Rice Bran



FM	Fishmeal
FTU	Unit of phytase
g	Gram
HPLC	High Pressure Liquid Chromatography
HU	Hemagglutinin Unit
IU	International Unit
k cal	Kilo Calorie
kg	Kilogram
MCP	Monocalcium phosphate
ME	Metabolisable Energy
Mg	Magnesium
mg	Milligram
ml	Mililitre
Mn	Manganese
NDF	Neutral Detergent Fibre
NFE	Nitrogen Free Extract
NRC	National Research Council
P	Phosphorus
PITC	Phenylisothyanate
pP	Phytate Phosphorus
PRP	Parboiled Rice Polish

RB	Rice Bran
rrf	Relative Response Factor
SBM	Soybean meal
Sd	Standard deviation
S.D.W	Sample Dry Weight
SE	Standard error
SEM	Standard Error of Means
sq	Square
tP	Total Phosphorus
TIU	Trypsin Inhibitor Unit
UPM	Universiti Putra Malaysia.
wt	Weight
Zn	Zinc
μl	Micro litre



Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Doctor of Philosophy.

EFFECT OF PHYTATE PHOSPHORUS ON THE UTILISATION OF RICE BRAN IN BROILER CHICKENS

By

MD ASADUZZAMAN UKIL

November 1999

Chairman: Assoc. Prof. Abdul Razak Alimon, Ph. D.

Faculty: Agriculture

Rice bran (RB) is a by-product of rice milling industry and is produced in many parts of the world including Malaysia. It is a high energy source for both ruminant and nonruminant animals with a crude protein content of 12-13%. Its fibre content is about 7-16%. Rice bran is also rich in phytate phosphorus (pP) which is not easily available to poultry. This study examines the utilisation of pP by broiler chickens fed RB based diets at the finishing stage. The role of phytase enzyme and vitamin D₃ in improving the utilisation of pP was also examined.

Two growth trials were carried out with 21 day-old broiler chickens to determine the effects of replacing corn with RB on their performance. It was found that the performance of birds were adversely affected when RB was included at 45% level. The results obtained from these two studies suggest that RB can be included up to 35% in the diet of broiler chicken without any adverse effects on their performance.



The third experiment was carried out to investigate the effects of supplementing phytase and vitamin D₃ in a phosphorus (P) deficient corn-RB-soybean meal based diet. Birds fed P deficient diets supplemented with phytase or vitamin D₃ showed similar performance with those fed diets adequate in phosphorus. The bioavailability of calcium (Ca) and P was increased and P excretion decreased with supplemental phytase or vitamin D₃.

In the fourth experiment, two P deficient diets consisting of 18 and 20% protein were formulated and were supplemented with 0, 350, 700 and 1050 FTU of phytase. The birds showed similar performance regardless of protein and phytase levels. Higher mineralisation was noted in the toe and tibia ash contents with higher levels of supplemental phytase. Toe and tibia ash contents were not affected by varying levels of proteins. At the same time P excretion was reduced with higher levels of phytase.

In the last study, two diets, deficient in available P with 18 and 20% crude protein were formulated. Varying levels of calcium was used to obtain three Ca:total P (Ca:tP) ratios, 1.1:1, 1:1 and 0.9:1 and six diets were formulated. The diets were supplemented with 700 FTU of phytase. The narrower Ca:tP ratio (0.9:1) had adverse effect on the performance of birds and P utilisation. Phosphorus excretion was increased with Ca:tP ratio of 0.9:1. It was observed that Ca:tP ratio could be reduced from 1.1:1 to 1:1 without any deleterious effect on the performance of broiler chickens.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi syarat keperluan untuk Ijazah Doktor Falsafah.

**KESAN FOSFORUS PHYTATE KEATAS PENGGUNAAN DEDAK PADI
PADA AYAM PEDAGING**

Oleh

MD. ASADUZZAMAN UKIL

November 1999

Pengerusi: Prof. Madya Abdul Razak Alimon, Ph. D.

Fakulti: Pertanian

Dedak padi adalah hasil sampingan daripada industri perkilangan beras yang dihasilkan di kebanyakan tempat di dunia termasuk Malaysia. Ia adalah sumber tenaga yang tinggi untuk haiwan ruminan dan bukan ruminan dengan kandungan protein kasar 12-13%. Kandungan serat ialah 7-16%. Dedak padi juga kaya dengan phytate fosforus, bentuk yang tidak mudah digunakan oleh ayam. Kajian ini menilai penggunaan phytate fosforus oleh ayam pedaging yang diberi makanan berasaskan dedak padi diperingkat penamat. Tindakan enzim phytase dan vitamin D₃ dalam meningkatkan penggunaan phytate fosforus juga dikaji.

Dua ujian pembesaran telah dilakukan ke atas ayam pedaging berumur 21 hari untuk melihat kesan penggantian jagung dengan dedak padi. Kesan sampingan dedak padi telah dilihat pada 45% aras rangsum. Keputusan kedua ujian ini menunjukkan bahawa dedak padi boleh di gunakan sehingga kearas 35% dalam rangsum ayam tanpa sebarang kesan sampingan keatas prestasi ayam.