



UNIVERSITI PUTRA MALAYSIA

**SELECTED HERBAL PLANTS AS GROWTH AND HEALTH
PROMOTERS IN BROILER CHICKENS**

**SEYED REZA HASHEMI
FP 2009 33**

**SELECTED HERBAL PLANTS AS GROWTH AND HEALTH PROMOTERS
IN BROILER CHICKENS**

By

SEYED REZA HASHEMI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

December 2009



THIS THESIS IS DEDICATED TO

MY WIFE, MY PARENTS

WITH

LOVE AND GRATITUDE

AND ALSO TO

THE KIDS WHO HAVE THE ABILITY

BUT

DON'T HAVE FACILITIES AND EDUCATION

**SELECTED HERBAL PLANTS AS GROWTH AND HEALTH PROMOTERS
IN BROILER CHICKENS**

By
SEYED REZA HASHEMI
December 2009

Chairman: Zulkifli B Idrus, PhD

Faculty: Agriculture

A series of experiments were conducted to examine the effect of herbal supplementation as a growth and health promoters. An *in vitro* procedure was used to determine the antibacterial activity of the fruit of *Solanum torvum*, whole plant of *Euphorbia hirta*, and rhizomes of *Zingiber officinale*, *Curcuma longa* and *Zingiber zerumbet*. Among the five herbal plants studied only the aqueous extract of *Euphorbia hirta* showed considerable growth-inhibiting activity against *Staphylococcus aureus*.

Phytochemical screening and a toxicity evaluation of the selected plants was carried out in experiment II. Phytochemical screening study showed that selected plants contained volatile oils, tannins, alkaloids, saponins, flavonoids. Alkaloids and steroids were only detected in the aqueous extract of *Euphorbia hirta*. Toxicity evaluation also showed that based on liver enzyme and histopathology the herbal aqueous extracts were not toxic when administered 2,000 mg/kg body weight by the oral route.

Experiment III was conducted to examine the effects of dietary *Euphorbia hirta* supplementation on growth performance, nutrient digestibilities, digesta pH and ileal microbial population. 600 one-day-old broilers (Cobb-500) were housed in 30 pens of 20 birds each until day 42. The dietary treatments were: (1) basal diet (control), (2) basal diet + 0.02 g/kg virginiamycin (VM), (3) basal diet + 1.5 g/kg acidifier (OrgacidsTM) (OA), (4) basal diet + 2.5 g/kg *E. hirta* (EH2.5), (5) basal diet + 5 g/kg *E. hirta* (EH5.0) and (6) basal diet + 7.5 g/kg *E. hirta* (EH7.5). The higher levels of *E. hirta* inclusion (7.5 g/kg) improved FCR of broiler chickens at 42 days of age. The highest AMEn and protein digestibility were observed in EH7.5 group. On day 21, the total aerobic bacteria counts in the ileum content of the EH2.5, VM and control groups were significantly higher ($P<0.05$) than the other groups. OA and VM supplementation significantly increased ($P<0.05$) *Lactobacillus* count at day 21 and the lowest *Lactobacillus* count was noted in the EH7.5 birds. At 42 days, feeding of EH2.5 promoted greater numbers of *lactobacillus*, to compare with other groups and the lowest *Lactobacillus* count was enumerated in the control birds. On day 21, except for ileum and cecum, OA significantly reduced ($P<0.05$) pH value. The trend in pH value of the gastrointestinal tract on day 42 and 21 was similar.

In experiment IV the effect of *E. hirta* and acidifier supplementation on gut colonization and organ invasion by *Salmonella enteritidis* was carried out. 120 day-old chicks were assigned to one of the three treatments: (1) basal diet (control), (2) basal diet + 7.5 g/kg *E. hirta* (EH), and (3) basal diet + 2 g/kg acidifier (OrgacidsTM) (OA). On day 3, each chick was inoculated orally with 1 mL of *S. enteritidis*

(1.5×10^8 cfu/bird). The control birds had significantly higher ($P < 0.05$) caecal *S. enteritidis* number than other groups following 7 and 14 days of challenge. Incidence of *S. enteritidis* colonization on the liver and spleen was significantly lower ($P < 0.05$) in OA birds following 7 days of challenge. The average of shedding time was longer in the control group than in the OA and EH7.5 treatments throughout the duration of the experiment.

Response of heat-stressed broiler chickens to *Zingiber zerumbet* and *Zingiber officinale* supplementation were investigated in the fifth experiment. A total of 300 male broiler chicks were randomly assigned to one of the following treatment. 1) Basal diet (control), 2) Basal diet +1% *Zingiber zerumbet* (ZZ1%), 3) Basal diet +2% *Zingiber zerumbet* (ZZ2%), 4) Basal diet +1% *Zingiber officinale* (ZO1%) and 5) Basal diet +2% *Zingiber officinale* (ZO2%). Diets were fed to the birds from day 28-42. At 35 days of age, half of the chickens were exposed to $38 \pm 1^\circ\text{C}$ for 2 h/d to another environmentally controlled chamber. Following 2 days of heat treatment heterophile/lymphocyte ratio and corticosterone were significantly elevated ($P < 0.05$). Significant feeding regimes \times heat treatment interaction was noted for serum glucose and potassium level ($P < 0.05$). Except for serum level of albumin, chloride, heterophile/lymphocyte ratio and plasma corticosterone seven days of heat treatment had no significant effect on blood parameters. Heat treatment markedly increases Hsp70 expression on day 42. Birds fed ZO2%, ZZ1% and ZZ2% had significantly greater ($P < 0.05$) Hsp70 response.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

TUMBUHAN HERBA YANG TERPILIH SEBAGAI PROMOTER PERTUMBUHAN DAN KESIHATAN DALAM AYAM PEDAGING

Oleh

SEYED REZA HASHEMI

Disember 2009

Pengerusi : Zulkifli B Idrus, PhD

Fakulti : Pertanian

Satu siri eksperimen telah dijalankan untuk memeriksa kesan penambahan herba sebagai penggalak pertumbuhan dan kesihatan. Suatu prosedur *in vitro* telah digunakan untuk menentukan aktiviti anti-bakteria daripada buah *Solanum torvum*, pokok *Euphorbia hirta*, rizom-rizom *Zingiber officinale*, *Curcuma longa* dan *Zingiber zerumbet*. Di antara lima tumbuhan herba yang dikaji, hanya cecair estrak daripada *Euphorbia hirta* dapat merencatkan aktiviti pembiakan *Staphylococcus aureus*.

Semakan fitokimia dan suatu penilaian ketoksidan terhadap tumbuh-tumbuhan yang terpilih telah dijalankan dalam eksperimen II. Kajian semakan fitokimia telah menunjukkan bahawa tumbuh-tumbuhan yang terpilih mengandungi minyak-minyak teruap, tannin, alkaloid, saponin dan flavonoid. Hanya alkaloid dan steroid didapati dalam cecair ekstrak daripada *Euphorbia hirta*. Penilaian ketoksidan telah menunjukkan

bahawa estrak herba adalah tidak toksik semasa diberi melalui mulut pada 2,000 mg/kg berat badan berdasarkan kajian pada enzim hati dan histopatologi.

Eksperimen III telah dijalankan untuk memeriksa kesan-kesan penambahan *Euphorbia hirta* dalam diet terhadap prestasi pertumbuhan, pencernaan nutrien, pH hasil pencernaan dan populasi mikrobia dalam usus ileum. 600 ekor ayam pedaging jenis Cobb 500 telah dikaji selama 42 hari di dalam 30 sangkar di mana setiap sangkar sebanyak 20 ekor ayam. Enam rawatan diet adalah: satu kawalan dan lima diet yang mengandungi: (1) diet basal (kawalan), (2) diet basal + 0.02 g/kg virginiamycin (VM), (3) diet basal + 1.5 g/kg asid organik (OrgacidsTM) (OA), (4) diet basal + 2.5 g/kg *E. hirta* (EH2.5), (5) diet basal + 5 g/kg *E. hirta* (EH5.0) dan (6) diet basal + 7.5 g/kg *E. hirta* (EH7.5). Asid organik tersebut mengandungi asid-asid formik, fosforik, laktik, tartarik, sitrik, dan malik yang dihasilkan oleh Sunzen Corporation Sdn. Bhd. Malaysia. Kandungan *E. hirta* yang tertinggi (7.5 g/kg) dapat memperbaikkan nisbah penukaran makanan dalam ayam-ayam pedaging pada umur 42 hari. AMEn dan pencernaan protein yang tertinggi dapat diperhatikan dalam kumpulan EH7.5. Pada umur 21 hari, jumlah bilangan bakteria aerobik dalam usus ileum yang terdapat dalam kumpulan EH2.5, VM dan kumpulan kawalan adalah signifikasinya lebih tinggi daripada kumpulan lain. Penambahan OA dan VM telah meningkatkan bilangan *Laktobasillus* dengan signifikan pada hari ke-21 dan bilangan *Lactobacillus* yang terendah adalah dicatatkan pada ayam EH7.5. Pada hari ke-42, pemberian kandungan makanan dengan EH2.5 telah menggalakkan penambahan bilangan *Laktobasillus* berbanding dengan kumpulan lain dan didapati bilangan *Laktobasillus* terlalu sedikit dalam ayam kawalan.

Pada hari ke-21, selain usus ileum dan sekum, OA dengan signifikan menurunkan nilai pH. Perubahan nilai pH dalam salur percernaan adalah sama pada hari 42 dan 21. Dalam eksperimen IV kesan penambahan *E. hirta* dan asid organik terhadap pembentukan koloni dalam usus dan serangan organ oleh *Salmonella enteritidis* telah dijalankan. Anak-anak ayam telah dibahagikan kepada salah satu daripada tiga kumpulan: (1) diet basal (kawalan), (2) diet basal + 7.5 g/kg *E. hirta* (EH), dan (3) diet basal + 2 g/kg asid organik (OrgacidsTM) (OA). Asid organik mengandungi asid-asid formik, fosforus, laktik, tartarik, sitrik, dan malik yang dihasilkan oleh Sunzen Corporation Sdn. Bhd. Malaysia. Pada hari ke-3, setiap anak ayam telah diberi 1 mL *Salmonella enteritidis* (1.5×10^8 cfu/ekor) secara oral. Ayam-ayam kawalan mempunyai nombor SE dalam sekum yang signifikasinya lebih tinggi daripada kumpulan lain selepas 7 dan 14 hari dikaji. Keadaan pembentukan koloni SE dalam hati dan limpa adalah signifikasinya lebih rendah dalam ayam OA selepas 7 hari dikaji. Masa purata pembebasan adalah lebih lama dalam kumpulan kawalan berbanding kumpulan rawatan OA dan EH7.5 di sepanjang tempoh eksperimen.

Tindakbalas tekanan haba oleh ayam-ayam pedaging dengan penambahan *Zingiber zerumbet* dan *Zingiber officinale* telah dikaji dalam eksperimen kelima. Sebanyak 300 ekor ayam pedaging telah dikaji dengan diet-diet eksperimen telah diberi kepada ayam dari umur 28-42 hari. Lima rawatan diet adalah: 1) diet basal (kawalan), 2) diet basal + 1% *Zingiber zerumbet* (ZZ1%), 3) diet basal + 2% *Zingiber zerumbet* (ZZ2%), 4) diet basal + 1% *Zingiber officinale* (ZO1%), 5) diet basal + 2% *Zingiber officinale* (ZO2%).

Pada umur 35 hari, separuh daripada ayam-ayam tersebut telah dikekalkan pada suhu

23°C manakala yang selebihnya telah didedahkan dengan suhu $38\pm1^{\circ}\text{C}$ selama 2 jam/hari di tempat lain yang persekitarannya terkawal. Cara-cara pemakanan dan rawatan haba tidak mempunyai kesan yang signifikan pada kadar kematian di sepanjang tempoh kajian. Rawatan haba selama dua hari dengan signifikan telah meningkatkan nilai albumin, jumlah protein, kolesterol, glukose, sodium dan klorida dalam serum. Selepas 2 hari rawatan haba, nisbah H/L dan CS didapati meningkat dengan signifikan. Interaksi di antara cara-cara pemakanan \times rawatan haba yang dicatatkan adalah signifikan bagi tahap glukosa dan potasium dalam serum. Selain nilai albumin dalam serum, klorida, nisbah H/L, dan kortikosteron dalam plasma, tujuh hari rawatan haba tidak memberikan kesan yang signifikan kepada parameter-parameter dalam darah. Rawatan haba telah meningkatkan pendedahan Hsp70 pada 42 hari. Ayam-ayam yang diberi ZO2%, ZZ1% dan ZZ2% mempunyai tindakbalas Hsp70 yang lebih banyak secara signifikan.

ACKNOWLEDGEMENTS

An effort such as this dissertation is only made possible through the immeasurable support, mentorship, time and patience of many individuals. I am indebted to all who have helped along the way and made the current work a reality. This work was made possible by a grant (05-01-04-SF0182) from the Ministry of Science, Technology and Innovation, Malaysia.

First and foremost, the author would like to thank Our God for teaching him the way of life. I would like to thank my family for their love and support given to me and most of all their trust and open door policy that kept me inspired through my program in oversea.

I wish to convey my heartfelt gratitude to the Malaysian government, University Putra Malaysia and my supervisor Professor Dr. Zulkifli for financial support.

My thanks and appreciations are expressed to my committee members, especially with patience, encouragement and instruction from Professor Dr. Zulkifli, I have enhanced my knowledge in both my class work and research.

Additionally, my thanks also are expressed to Ponnusamy Muniandy, our farm technician and Mr. Yap lab technician in hormonology lab, have given me lots of help in my research and farm work.

The work would not have been made possible without the efforts of Iranian students in Malaysia. My graduate fellows have given lots of help in my research, Abdul Reza Soleimani, Saeed Navid, Meisam Tabatabaei, Mehdi Ebrahimi and specially Tang Siew Ching gave great help for data collection and thankful to Tang Siew Ching and Nor Azwati for thesis abstract translation to Bahasa Melayu.

I would like to extend my gratitude to all the teachers from Iran who provided us the indispensable theoretical knowledge especially my former supervisor Dr. Behroz Dastar and Mr. Foladian from high school.

Finally, I am grateful to have found someone to stand beside me and continually remind me not to take life so seriously, especially when an experiment fails. So my final thanks go to my wife Homa Davoodi. The only way to thank you properly would be to promise to always give you the level of love and support that you've given me.

I certify that a Thesis Examination Committee has met on 21 December 2009 to conduct the final examination of Seyed Reza Hashemi on his thesis entitled "Selected Herbal Plants as Growth and Health Promoters in Broiler Chickens " in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the degree of Doctor of Philosophy (Phd).

Members of the Thesis Examination Committee were as follows:

Ismail Idris, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Chairman)

Norhani binti Abdullah, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

Abdul Razak Alimon, PhD

Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Internal Examiner)

D. Michael Denbow, PhD

Professor

Department of Animal and Poultry Sciences, Virginia Polytechnic Institute and State University

USA

(Independent Examiner)

BUJANG BIN KIM HUAT, PhD

Professor and Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Zulkifli B Idrus, PhD

Professor

Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Mohd Hair B Bejo, PhD

Professor

Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Muhammad Nazrul Hakim Bin Abdullah, PhD

Associate Professor

Faculty of Medicine And health Sciences
Universiti Putra Malaysia
(Member)

Loh, Teck Chwen, , PhD

Associate Professor

Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Zunita Binti Zakaria, PhD

Associate Professor

Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean

School of Graduate studies
Universiti Putra Malaysia

Date: 11 February 2010

DECLARATION

I declare that the thesis is my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously and it is not concurrently submitted for any other degree at Universiti Putra Malaysia or other institutions.

SEYED REZA HASHEMI

Date: 11 February 2010

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	x
APPROVAL	xii
DECLARATION	xiv
LIST OF TABLE	xix
LIST OF FIGURES	xxii
LIST OF ABBRIVATION	xxiii
1 INTRODUCTION	
1.1 General introduction	1
1.2 General objectives	7
1.3 Specific objectives	7
2 LITERATURE AND REVIEW	
2.1 The Important of Herbal Plants	9
2.2 <i>Curcuma longa</i>	
2.2.1. Botanical characteristics of <i>Curcuma longa</i> plant	11
2.2.2. The use of <i>Curcuma longa</i> as a traditional medicinal plant	11
2.2.3. Chemical compound of <i>Curcuma longa</i>	12
2.2.4. Previous Studies	13
2.3 <i>Zingiber officinale</i>	
2.3.1 Botanical characteristics of <i>Zingiber officinale</i> plant	17
2.3.2 The use of <i>Zingiber officinale</i> as a traditional medicinal plant	17
2.3.3 Chemical compound of <i>Zingiber officinale</i>	18
2.3.4 Previous Studies	19
2.4 <i>Zingiber zerumbet</i>	
2.4.1 Botanical characteristics of <i>Zingiber zerumbet</i> plant	23
2.4.2 The use of <i>Zingiber zerumbet</i> as a traditional medicinal plant	23
2.4.3 Chemical compound of <i>Zingiber zerumbet</i>	24
2.4.4 Previous Studies	25
2.5 <i>Euphorbia hirta</i>	
2.5.1 Botanical characteristics of <i>Euphorbia hirta</i> plant	28
2.5.2 The use of <i>Euphorbia hirta</i> as a traditional medicinal plant	29
2.5.3 Chemical compound of <i>Euphorbia hirta</i>	30
2.5.4 Previous Studies	31
2.6 <i>Solanum torvum sw.</i>	
2.6.1 Botanical characteristics of <i>Solanum torvum</i> plant	42
2.6.2 The use of <i>Solanum torvum</i> as a traditional medicinal plant	42

2.6.3	Chemical compound of <i>Solanum torvum</i>	43
2.6.4	Previous Studies	44
2.7	Use of herbal plants and herbal derivatives in poultry research	45

3 MATERIALS AND METHODS

CHAPTER 3: ANTIMICROBIAL ACTIVITY OF AQUEOUS EXTRACT OF HERBAL PLANTS

3.1	Introduction	48
3.2	Materials and Methods	
3.2.1	Plant materials	51
3.2.2	Preparation of extract	51
3.2.3	Test Microorganism	52
3.2.4	Antimicrobial sensitivity test	52
3.3	Results	53
3.4	Discussion	57
3.5	Conclusion	59

CHAPTER 4: PHYTOCHEMICAL SCREENING AND ACUTE TOXICITY STUDY OF SELECTED HERBAL AQUEOUS EXTRACT IN BROILER CHICKENS

4.1	Introduction	61
4.2	Materials and Methods	
4.2.1	Plant materials	63
4.2.2	Preparation of extract	64
4.2.3	Preliminary phytochemical analysis	65
4.2.4	Acute toxicity study	68
4.2.5	Experimental design	69
4.2.6	Behavioral study	70
4.2.7	Histopathological studies	70
4.2.8	Blood sample collection	71
4.2.9	Statistical analysis	71
4.3	Results	
4.3.1	Phytochemical screening	72
4.3.2	Behavioral study	74
4.3.3	Effect of extract on tissue morphology of chickens	76
4.3.4	Serum metabolites and liver function test (LFT)	78
4.3.5	Growth performance	80
4.4	Discussion	81
4.5	Conclusion	85

CHAPTER 5: EFFECTS OF *EUPHORBIA HIRTA* SUPPLEMENTATION ON GUT MICROFLORA, ILEAL DIGESTIBILITY AND GROWTH PERFORMANCE OF BROILER

CHICKENS

5.1	Introduction	87
5.2	Materials and Methods	
5.2.1	Herbal supplement preparation	90
5.2.2	Birds and housing environment	91
5.2.3	Experimental design	92
5.2.4	Performance parameters	94
5.2.5	Enumeration of Bacteria	94
5.2.6	Ileal Digestibility and AMEn	95
5.2.7	Measurement of digestive tract pH	97
5.2.8	Liver function enzymes	97
5.2.9	Statistical analysis	98
5.3	Results	
5.3.1	Growth performance	99
5.3.2	Bacteria enumeration	101
5.3.3	AMEn and protein digestibility values	103
5.3.4	Digestive tract pH	105
5.3.5	liver function enzymes	105
5.4	Discussion	107
5.5	Conclusion	113

CHAPTER 6: EFFECTS OF *EUPHORBIA HIRTA* SUPPLEMENT ON RESISTANCE OF BROILER CHICKENS AGAINST *SALMONELLA ENTERITIDIS* INFECTION

6.1	Introduction	115
6.2	Materials and Methods	117
6.2.1	Chickens	118
6.2.2	Experimental Design	118
6.2.3	<i>Salmonella</i> Source for Challenge and Inoculation Procedure	120
6.2.4	SE shedding	120
6.2.5	Sampling Protocol for SE organ invasion and SE Concentration in Cecal Content	121
6.2.6	Statistical analysis	123
6.3	Results	
6.3.1	Cecal colonization by SE	124
6.3.2	Organ Colonization by SE	126
6.3.3	Cloacal swabs SE enumeration	128
6.4	Discussion	129
6.5	Conclusion	133

CHAPTER 7: CHANGE IN GROWTH, BLOOD PARAMETERS AND HEAT SHOCK PROTEIN 70 EXPRESSION IN HEAT STRESSED BROILER CHICKENS IN RESPONSE TO DIETARY

SUPPLEMENTATION OF *ZINGIBER ZERUMBET* AND *ZINGIBER OFFICINALE*

7.1	Introduction	135
7.2	Materials and Methods	138
7.2.1	Plant materials	139
7.2.2	Birds and diets	139
7.2.3	Experimental procedure	141
7.2.4	Performance parameters and mortality of broiler chickens	142
7.2.5	Measurement of body temperatures	142
7.2.6	Measurement of Corticosterone, heterophil/lymphocyte ratio, blood biochemical parameters	143
7.2.7	Determination of Hsp70 concentration	145
7.2.8	Statistical analysis	147
7.3	Results	147
7.4	Discussion	156
7.5	Conclusion	161
8	GENERAL DISCUSSION AND RECOMMENDATIONS FOR FUTURE WORK	
7.6	General Discussion	162
7.7	Recommendations for future work	166
9	REFERENCES	166
10	APPENDICES	206
11	BIODATA OF STUDENT	219
12	LIST OF PUBLICATIONS	220

LIST OF TABLES

Table	Page
3.1. Antibacterial activity of various concentrations of aqueous extracts of <i>Zinziber officinale</i> , <i>Curcuma long</i> , <i>Zinziber zerumbet</i> , <i>Euphorbia hirta</i> and <i>Solanum torvum</i> .	55
3.2. Mean inhibition zones of <i>Euphorbia hirta</i> aqueous extracts in comparison with standard antibiotics.	56
4.1. Phytochemical screening of selected herbal plants.	73
4.2. Toxicity signs observed in broiler (first 24h) that received oral single dose (2000 mg/kg BW) of the aqueous extract of herbal plants.	75
4.3. Clinical blood chemistry values of female broilers administered with aqueous extract of herbal plants (mean \pm SEM).	79
4.4. The effect of high dose aqueous extract of selected herbal plants on body weight gain, feed intake and feed conversion in broilers.	81
5.1. Ingredients and nutrient composition of diets.	93
5.2. The effect of <i>Euphorbia hirta</i> , acidifier and virginiamycin supplementation on body weight, body weight gain, feed intake, feed conversion ratio (FCR) and mortality rate in broilers (mean \pm SEM).	100

- 5.3. The effect of *Euphorbia hirta*, acidifier and virginiamycin supplementation on viable cell counts of microflora in ileum of broiler chickens (mean±SEM). 102
- 5.4. The effect of *Euphorbia hirta*, acidifier and virginiamycin supplementation on apparent metabolizable energy (AME), apparent metabolizable energy nitrogen-corrected (AMEn), protein and energy digestibility in broiler chickens (mean ± SEM). 104
- 5.5. The effect of *Euphorbia hirta*, acidifier and virginiamycin on pH in gastrointestinal contents at 21and 42 d (mean ± SEM). 106
- 5.6. The effect of *Euphorbia hirta*, acidifier and virginiamycin supplementation on liver function enzymes values of broiler chickens (mean ± SEM). 107
- 6.1. The effect of *Euphorbia hirta* (EH) and acidifier (OA) supplementation on *Salmonella enterica* serovar *Enteritidis* cecal colonization of broiler (mean±SEM). 125
- 6.2. The effect of *Euphorbia hirta* (EH) and acidifier (OA) supplementation on *Salmonella enterica* serovar *Enteritidis* (SE) colonization of the crop, liver, spleen, bile bladder and apex of broiler. 127
- 6.3. The effect of *Euphorbia hirta* (EH) and acidifier (OA) supplementation on *Salmonella enterica* serovar Enteritidis (SE) cloacal swab in broiler. 129
- 7.1. Ingredients and nutrient composition of basal diets. 140

- 7.2. The effects of feeding regimes and heat treatment on body weight (BW), body weight gain (BWG), feed intake (FI) and feed conversion ratio (FCR) of broiler chickens from d 35-42 (mean \pm SEM). 151
- 7.3. The effects of feeding regimes and heat treatment on blood parameters and body temperature value of broiler chickens on d 37 (mean \pm SEM). 152
- 7.4. Blood parameter values of broiler chickens where feeding regimes \times heat treatment interaction were significant on d 37 (means \pm SEM). 153
- 7.5. The effects of feeding regimes and heat treatment on blood parameters and body temperature value of broiler chickens on d 42 (mean \pm SEM). 154
- 7.6. Blood parameter values and body temperature of broiler chickens where feeding regimes \times heat treatment interaction were significant on d 42 (means \pm SEM). 155
- 7.7. The effect of feeding regimes and heat treatment on HSP70 density value of broiler chickens on d 42 (mean \pm SEM). 156

LIST OF FIGURES

Figure		Page
3.1.	Antibacterial inhibition zone by <i>E.hita</i> .	57
3.2.	Significant increase in the inhibitory zone with increasing the concentration of the <i>E. hita</i> against <i>S. aureus</i> at 50, 100 and 200 mg/mL concentration.	57
4.1.	Liver histology of untreated and treated chicks with aqueous extract of selected herbal plants (2000mg/kg bw) at 35 days of age.	76
4.2.	Kidney histology of untreated and treated chicks with aqueous extract of selected herbal plants (2000mg/kg bw) at 35 days of age.	77

LIST OF ABBREVIATIONS

Abs	Antibodies
Ag	Antigens
ALT	Alanine aminotransferase
ALP	Alkaline phosphatase
AME	Apparent metabolizable energy
AMEn	Apparent metabolizable energy nitrogen corrected
AP	<i>Andrographis paniculata</i>
AST	Aspartate aminotransferase
ATCC	Ameriacan Type Culture Collection
BCA	Bicinchoninic acid
<i>B. cereus</i>	<i>Bacillus cereus</i>
BSA	Bovine Serum Albumin
<i>B. subtilis</i>	<i>Bacillus subtilis</i>
BW	Body weight
Caco-2	Human colon carcinoma epithelial cell line
CDC	Center for Disease Control and Prevention
<i>C. longa</i>	<i>Curcuma longa</i>
CORT	Corticosterone
COX-2	Cyclooxygenase -2 (an enzyme responsible for inflammation)
<i>E. coli</i>	<i>Escherichia coli</i>
EDTA	Ethylenediaminetetraacetic
<i>E. hirta</i>	<i>Euphorbia hirta</i>
ELISA	Enzyme-Linked ImmunoSorbent Assay
EPEC	Enteropathogenic <i>E. coli</i>
EU	European Union
FAO	Food and Agriculture Organization
FDA	United States Food and Drug Administration
FEDESA	Fédération Européenne de la Santé Animale (European Federation of Animal Health)

FCR	Feed conversion ratio
FR	Feed Restriction
GE	Gross energy
GC	Gas Chromatography
GLC	Gas Liquid Chromatography
GOT	Glutamine oxaloacetic transaminase
GPT	Glutamine pyruvic transaminase
HDL	High density lipoprotein
<i>H. influenzae</i>	<i>Haemophilus influenzae</i>
H/L ratio	Heterophile/Lymphocyte Ratio
HPLC	High Performance Liquid Chromatography
IC ₅₀	Median Inhibition Concentration (the concentration of substance that provides 50% inhibition to certain reaction)
IBS	Institute of Biosciences
IgA	Immunoglobulin A
IgG	Immunoglobulin G
IgM	Immunoglobulin M
IL	Interleukins
iNOS	Inducible Nitric oxide synthases
LC ₅₀	Lethal Concentration 50 (concentration in water having 50% chance of causing death to aquatic life)
LDL	Low density lipoprotein
LD50	lethal dose of 50% of treated target animals
<i>L. monocytogenes</i>	<i>Listeria monocytogenes</i>
MAP kinase	Mitogen-activated protein kinase
MBC	Minimum Bactericidal Concentration
ME	Metabolisable energy
MIC	Minimal inhibitory concentration
MR	Mortality rate
MRLs	Maximum residue limits
n	Number