



UNIVERSITI PUTRA MALAYSIA

**EFFECTS OF POLYPHENOL AND pH ON COCOA FLAVOUR
PRECURSORS AND FLAVOUR COMPOUNDS USING LIPIDIC
MODEL SYSTEM**

**NOOR SOFFALINA SOFIAN SENG
FSTM 2009 32**



**EFFECTS OF POLYPHENOL AND pH ON COCOA FLAVOUR PRECURSORS
AND FLAVOUR COMPOUNDS USING LIPIDIC MODEL SYSTEM**

NOOR SOFFALINA SOFIAN SENG

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2009



**EFFECTS OF POLYPHENOL AND pH ON COCOA FLAVOUR PRECURSORS
AND FLAVOUR COMPOUNDS USING LIPIDIC MODEL SYSTEM**

By

NOOR SOFFALINA SOFIAN SENG

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

October 2009



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science.

EFFECTS OF POLYPHENOL AND pH ON COCOA FLAVOUR PRECURSORS AND FLAVOUR COMPOUNDS USING LIPIDIC MODEL SYSTEM

By

NOOR SOFFALINA SOFIAN SENG

October 2009

Chairman : Jinap Selamat, PhD

Faculty : Food Science and Technology

Polyphenol is a naturally occurring compound in cocoa beans reported to reduce the concentration of flavour compounds, mainly pyrazines after cocoa roasting. This study was carried out to determine the effect of polyphenol concentrations and pH on cocoa flavour precursors (i.e. reducing sugars and amino acids) and flavour compounds (i.e. alkylpyrazines) during roasting. The study was conducted using a lipidic model system at roasting temperature 120⁰C for 45 min. The concentration of amino acids and reducing sugars incorporated in the model systems was based on the results of the concentration found in fermented Malaysian cocoa beans.

To determine the effect of polyphenol concentration, crude polyphenol extracted from cocoa bean was added into the lipidic model system at 0, 58, 116 and 174 g kg⁻¹. Results showed that the higher concentrations of polyphenols (174 g kg⁻¹) significantly (p<0.05) decreased the concentration of reducing sugars and amino acids during roasting. As the concentration of polyphenol increased from 58, 116 to 174 g kg⁻¹,



fructose was degraded at 33, 35, and 41% and glucose at 29, 30 and 33% respectively. Seventeen individual amino acids were detected, among all; alanine, tyrosine, valine and isoleucine were significantly ($p < 0.05$) degraded higher than the others. These amino acids which are in the hydrophobic group were degraded at 41, 44, 45 and 43 % respectively at polyphenol concentration 174 g kg^{-1} . Changes in procyanidins monomeric and polymeric constituents were correlated with the reduction of the precursors. High polyphenol concentration (174 g kg^{-1}) significantly decreased the formation of alkylpyrazines in the model system.

The Response Surface Methodology (RSM) plots revealed that lower concentration of amino acids and reducing sugars was obtained at higher polyphenol concentration (120 g kg^{-1}) and lower pH value (4.5). Based on the constraints set on the obtained surface plots, the optimum region for polyphenol concentration of $43 - 58 \text{ g kg}^{-1}$ and pH of 7.0 -7.5 was obtained for the optimum production of flavour precursors; while the optimum region polyphenol concentration of $< 75 \text{ g kg}^{-1}$ and pH of > 6.0 was obtained for the production of alkylpyrazines. The optimum point was determined using response optimizer and it was found that the optimum polyphenol concentration of 49.5 g kg^{-1} and pH of 7.2 for flavor precursors and polyphenol concentration of 52.3 g kg^{-1} and pH of 7.0 for generated alkylpyrazines. Thus, it is recommended that the acidity of the beans should be controlled during processing and extracting the excessive amount of polyphenol higher than 52.3 g kg^{-1} from the cocoa beans should be considered to lower its negative effect on reducing sugars and amino acids.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains.

KESAN POLIFENOL DAN pH KE ATAS PELOPOR RASA KOKO DAN SEBATIAN RASA KOKO MENGGUNAKAN MODEL SISTEM BERLIPID

Oleh

NOOR SOFFALINA SOFIAN SENG

Oktober 2009

Pengerusi : Jinap Selamat, PhD

Fakulti : Sains Makanan dan Teknologi

Polifenol adalah salah satu sebatian yang terkandung secara semulajadi di dalam biji koko, dan dikatakan dapat merendahkan komponen perasa terutamanya pirazin selepas proses pemanggangan koko. Kajian ini dijalankan untuk mengenalpasti kesan kepekatan polifenol dan pH ke atas pelopor rasa koko (gula penurunan dan asid amino) dan juga komponen rasa koko (alkilpirazin) selepas pemanggangan koko. Kajian ini dijalankan menggunakan model sistem berlipid pada suhu pemanggangan 120⁰C selama 45 minit. Kepekatan asid amino dan gula penurunan yang digunakan di dalam model sistem adalah berdasarkan data yang diperolehi daripada biji koko tertapai Malaysia.

Polifenol kasar yang diekstrak daripada biji koko kemudiannya ditambah ke dalam model sistem berlipid pada kepekatan 0, 58, 116 and 174 g kg⁻¹. Keputusan mendapati kepekatan polifenol yang tinggi (174 g kg⁻¹) telah merendahkan kepekatan gula penurunan dan asid amino secara signifikan (p<0.05). Apabila kepekatan polifenol meningkat dari 58, 116 ke 174 g kg⁻¹, fruktosa menurun sebanyak 33, 35, dan 41%; dan glukosa

sebanyak 29, 30 dan 33% masing-masing. Sebanyak tujuh belas asid amino dikesan dalam kajian ini, di mana alanina, tirosina, valina dan isoleusina didapati lebih menurun berbanding yang lain. Asid amino - asid amino ini, yang tergolong di dalam kumpulan hidrofobik telah menurun sebanyak 41, 44, 45 dan 43 % masing-masing pada kepekatan polifenol 174 g kg⁻¹. Perubahan pada komponen monomer dan polimer prosianidin juga dikaji dan dihubungkan dengan penurunan pelopor rasa. Kepekatan polifenol yang tinggi (174 g kg⁻¹) juga didapati mempunyai kesan yang signifikan dalam merendahkan penghasilan alkilpirazina di dalam model sistem.

Plot yang dihasilkan menggunakan Kaedah Permukaan Respons menunjukkan bahawa kepekatan asid amino dan gula penurun yang rendah diperolehi pada kepekatan polifenol yang tinggi (120 g kg⁻¹) dan pH yang rendah (4.5). Berdasarkan kepada pencirian yang ditetapkan, kawasan kepekatan polifenol yang optimum ialah pada 43 – 58 g kg⁻¹ dan pH pada 7.0 -7.5, bagi pelopor rasa koko. Manakala, kawasan kepekatan polifenol yang optimum pada <75 g kg⁻¹ dan pH pada > 6.0 dapat mengoptimumkan penghasilan alkilpirazina. Titik optimum diperolehi menggunakan pengoptimum respons mendapati titik optimum polifenol pada 49.5 g kg⁻¹ dan pH 7.2 bagi pelopor perisa koko, manakala titik optimum bagi penghasilan alkilpirazina adalah pada kepekatan polifenol pada 52.3 g kg⁻¹ dan pH 7.0 . Oleh itu, adalah dicadangkan tahap keasidan biji koko dikawal semasa pemprosesan dan pengestrakan polifenol yang tinggi dari 52.3 g kg⁻¹ pula dapat menurunkan kepekatannya ke tahap yang diperlukan bagi menurunkan kesan negatifnya ke atas gula penurun dan asid amino.

ACKNOWLEDGEMENTS

In the name of Allah, Most Gracious Most Merciful and *salawat* and *salam* to His righteous messenger, Prophet Muhammad SAW.

This research project would not have been possible without the support of many people. The author wishes to express her gratitude and thanks to her supervisor, Prof. Dr. Jinap Selamat. Deepest gratitude and appreciations are also due to the members of the supervisory committee, Prof. Dr. Nazamid Saari and special thanks go to Dr. Nazimah Sheikh Abdul Hamid who constantly have faith in the author and helped make some sense of the confusion.

Sincere appreciation goes to the lecturers and staffs of the Faculty of Food Science and Technology for their help and cooperation. The author would also like to convey her utmost gratitude to Universiti Putra Malaysia (UPM), in which she had the opportunity to be part of this magnificent institution for so many years. The author would also like to convey thanks to the Ministry of Science, Technology and Innovation (MOSTI) Malaysia and the Faculty of Food Science and Technology, UPM for providing the financial means and laboratory facilities, respectively without which the research work would not had been possible.

The author would like to take this opportunity to express gratitude to her family, whose constant encouragement and love she hold high through the duration of her studies especially her beloved mother, Hajjah Fatimah Abdul Rahman whose compassion and



love are what every daughter sought for. Deepest gratitude to her beloved father, Mr. Lim Khai Seng @ Sofian Seng Bin Abdullah and her one and only brother, Muhammad Noorsoffazli for being the family in which she found her pillar of strength. As a gift for enduring this long process with her, it is only to thee that she dedicates this work.

The author is indebt to all her friends whose unflinching conviction towards her keeps her always inspired. Thank you for the friends she holds dear, who share both work and play with her, who offer her support and love that she relied throughout the process. Many thanks to her bestfriend, Yang Hayati Abdul Razak that hear her unspoken needs, recognize her secret dreams, and understand the silent things that only true friends know. Thank you so very much.

The author also expresses sincere remembrance and gratitude to those who contribute to her being today but are no longer here to share her joy. May Allah bless you. And to the people pursuing in the name of science, the author hopes to continue, in her own way though her contribution might be very small, the noble mission to which others had sacrificed their lives for.

I certify that a Thesis Examination Committee has met on 23rd October 2009 to conduct the final examination of Noor Soffalina Binti Sofian Seng on her thesis entitled " Effect of polyphenol and pH on cocoa flavour precursors and alkylpyrazines in lipidic model system " in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science degree).

Members of the Thesis Examination Committee were as follows:

Azis Ariffin, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Lasekan Olusegun, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Internal Examiner)

Seyed Hamed Mirhosseini, PhD

Faculty of Food Science and Technology
Universiti Putra Malaysia
(Internal Examiner)

Azhar Mat Easa, PhD

Associate Professor
School of Industrial Technology
Universiti Sains Malaysia
(External Examiner)

BUJANG KIM HUAT, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 15 January 2010

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Jinap Selamat, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Nazamid Saari, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Nazimah Sheikh Abdul Hamid, PhD

Lecturer
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia
Date: 11 February 2010

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

NOOR SOFFALINA BT SOFIAN SENG

Date: 25 February 2010

TABLE OF CONTENTS

	Page
DEDICATION	2
ABSTRACT	3
ABSTRAK	5
ACKNOWLEDGEMENT	7
APPROVAL	9
DECLARATION	11
LIST OF TABLES	16
LIST OF FIGURES	18
LIST OF ABBREVIATIONS	21
 CHAPTERS	
1 GENERAL INTRODUCTION	
1.1 Introduction	22
1.2 Statement of Problem	28
1.3 Objectives of the Study	30
2 LITERATURE REVIEW	
2.1 Cocoa Beans	31
2.2 Cocoa Fermentation	32
2.2.1 Acidity and pH	35
2.2.2 Changes in Protein	38
2.2.3 Changes in Sugars	42
2.2.4 Changes in Polyphenol	43
2.2.5 Importance of fermentation on cocoa flavour precursors development	45
2.3 Cocoa Roasting	48
2.3.1 Maillard Reaction	50
2.3.2 Model System in Maillard Reactions	55
2.3.3 Importance of roasting on cocoa flavour Development	58
2.4 Cocoa Alkalization	59
2.5 Cocoa Flavour	62
2.5.1 Volatiles Compounds	65
2.5.2 Pyrazines	66
2.5.3 Extraction of pyrazines using Solid-Phase Microextraction (SPME)	66



2.6	Cocoa Polyphenol	71
2.6.1	Chemistry and Biosynthesis of Polyphenol	76
2.6.2	Polyphenol on Health	78
2.6.3	Polyphenol Interactions	78
2.6.4	Astringency and Bitterness	80
3	GENERAL MATERIALS AND METHODS	
3.1	Materials	
3.1.1	Source of Cocoa Beans For Extraction of Cocoa Butter and Crude Polyphenol	83
3.1.2	Source of Cocoa Beans for Development of Lipidic Model	83
3.1.3	Reagents	84
3.2	Methodology	
3.2.1	Lipidic Model System for Cocoa Roasting Process	84
3.2.2	Cocoa Butter Extraction	85
3.2.3	Crude Polyphenol Extraction	86
3.2.4	Polyphenol Analyses	86
3.2.5	Free Amino Acids Analysis	88
3.2.6	Sugars Analysis	90
3.2.7	Statistical Analysis	91
4	EFFECT OF POLYPHENOL ON COCOA FLAVOUR PRECURSORS AND ALKYLPIRAZINES IN A LIPIDIC MODEL SYSTEM	
4.1	Introduction	92
4.2	Materials and Methods	94
4.2.1	Chemicals and Standards	94
4.2.2	Source of Cocoa Beans for Extraction of Cocoa Butter and Crude Polyphenol	94
4.2.3	Cocoa Butter Extraction	94
4.2.4	Crude Polyphenol Extract	95
4.2.5	Determination of Free Amino Acids	95
4.2.6	Determination of Sugars	95
4.2.7	Determination of Alkylpyrazines	95
4.2.8	Lipidic Model System	96
4.2.8	Experimental Design	97
4.2.9	Statistical Analysis	97

4.3	Results and Discussion	99
4.3.1	Development of Lipidic Model System	99
4.3.2	Effect of Temperature on Sugars, Amino Acids and Procyanidins	102
4.3.3	Effect of Polyphenol Concentration on Amino Acids During Roasting and Changes in Procyanidins Polymers	106
4.3.4	Effect of Polyphenol Concentration on Reducing Sugar During Roasting and Changes in Procyanidins Polymers	115
4.3.5	Method Development for SPME Conditions for Detection of Alkylpyrazines in Lipidic Model System	118
4.3.6	Effect of Polyphenol Concentration on Alkylpyrazines During Roasting and Changes in Procyanidins Polymers	124
4.4	Conclusion	131

5 COMBINED EFFECT OF POLYPHENOL AND pH ON COCOA FLAVOUR PRECURSORS AND ALKYLPIRAZINES IN A LIPIDIC MODEL SYSTEM USING RESPONSE SURFACE METHODOLOGY (RSM)

5.1	Introduction	133
5.2	Materials and Methods	135
5.2.1	Chemicals and Standards	135
5.2.2	Source of Cocoa Beans for Extraction of Cocoa Butter and Crude Polyphenol	136
5.2.3	Cocoa Butter Extraction	136
5.2.4	Crude Polyphenol Extract	136
5.2.5	Determination of Free Amino Acids	136
5.2.6	Determination of Sugars	137
5.2.7	Determination of Alkylpyrazines	137
5.2.8	Lipidic Model System	137
5.2.9	Experimental Design	138
5.2.10	Statistical Analysis	140
5.3	Results and Discussion	
5.3.1	Effect of polyphenol and pH on flavour precursors	140
5.3.2	Effect of Polyphenol Concentration and pH on Cocoa Flavour Compounds (Alklypyrazines)	174
5.3.3	Conclusion	196

6	CONCLUSIONS AND RECOMMENDATION	
6.1	General Conclusions	197
6.2	Recommendations	199
	BIBLIOGRAPHY	202
	APPENDICES	226
	BIODATA OF STUDENT	234
	LIST OF PUBLICATIONS	235



LIST OF TABLES

Table		Page
2.1	Changes in cocoa composition during fermentation	45
2.2	Different types of substrates and reaction conditions of Maillard-model studied by several authors.	56
2.3	Changes in free amino acids and sugars ($\mu\text{M g}^{-1}$) in cocoa during Processing, with the percentage change relative to the previous column in parentheses	61
2.4	Volatile compounds analysis in several samples using Solid Phase Microextraction (SPME) techniques	70
2.5	Different polyphenols identified in cocoa beans and cocoa Products	75
4.1	Concentration of amino acids, sugars, polyphenol, fat and moisture in fermented Malaysian cocoa beans.	101
4.2	Effect of roasting temperature of 120°C on the degradation amino acids, sugars and procyanidins	105
4.3	Effect of polyphenol concentration on amino acid degradation (%)	109
4.4	Changes in procyanidin content (% loss) after 45 mins roasting with amino acids	110
4.5	Correlation coefficient of amino acids degradation and procyanidin degradation during roasting in lipidic model system	114
4.6	Effect of polyphenol concentration on sugars degradation	117
4.7	Changes in procyanidin content (% loss) after 45 mins roasting with sugars.	117
4.8	Correlation coefficient of sugars degradation and procyanidin degradation during roasting in lipidic model system	118
4.9	Effect of polyphenol on the formation of alkylpyrazines (mg g^{-1}) in lipidic model system	126
4.10	Changes in procyanidin content (% loss) after 45 mins roasting with amino acids and sugars	128
4.11	Correlation coefficient of alkylpyrazines formation	

	and procyanidin degradation during roasting in lipidic model system	129
5.1	Independent variables and their levels in central composite design	139
5.2	Central composite design and experimental result for the response function of cocoa flavour precursors	143 144
5.3	Analysis of variance on the independent variables as linear, quadratic and interaction terms on the response variables of cocoa flavour precursors	145 146
5.4	Regression coefficients of the second-order polynomial for the response variables of cocoa flavour precursors	150 151
5.5	Regression coefficients and lack-of-fit for the reduced response surface models of cocoa flavour precursors	152 153
5.6	Central composite design and experimental result for the response function of alkylpyrazines	177
5.7	Analysis of variance on the independent variables as linear, quadratic and interaction terms on the response variables of alkylpyrazines	178
5.8	Regression coefficients of the second-order polynomial for the response variables of alkylpyrazines	180
5.9	Regression coefficients and lack-of-fit for the reduced response surface models of alkylpyrazines	181

LIST OF FIGURES

Figure		Page
2.1	Simplified flow diagram of cocoa fermentation	37
2.2	General overview of the Maillard reaction showing flavour compounds as end products	54
2.3	Structure of common polyphenols found in cocoa and cocoa products	75
4.1	Experimental flow for the study of polyphenol effect on cocoa flavour precursors and alkylpyrazine in lipidic model system	98
4.2	Effect of type of SPME fiber and adsorption time on the peak area of alkylpyrazines.	121 122
5.1	Surface plot of sucrose concentration as affected by the polyphenol concentration and pH variable	148
5.2	Surface plot of asparagine concentration as affected by the polyphenol concentration and pH variable	154
5.3	Surface plot of arginine concentration as affected by the polyphenol concentration and pH variable	155
5.4	Surface plot of histidine concentration as affected by the polyphenol concentration and pH variable	156
5.5	Surface plot of alanine concentration as affected by the polyphenol concentration and pH variable	157
5.6	Surface plot of iso-leucine concentration as affected by the polyphenol concentration and pH variable	158
5.7	Surface plot of tyrosine concentration as affected by the polyphenol concentration and pH variable	159
5.8	Surface plot of valine concentration as affected by the polyphenol concentration and pH variable	160

5.9	Surface plot of glycine concentration as affected by the polyphenol concentration and pH variable	162
5.10	Surface plot of methionine concentration as affected by the polyphenol concentration and pH variable	165
5.11	Surface plot of serine concentration as affected by the polyphenol concentration and pH variable	163
5.12	Surface plot of threonine concentration as affected by the polyphenol concentration and pH variable	166
5.13	Superimposed contour plot of significant variables for flavour precursors	173
5.14	Surface plot of 2-Methylpyrazine concentration as affected by the polyphenol concentration and pH variable	182
5.15	Surface plot of 2,5-Dimethylpyrazine concentration as affected by the polyphenol concentration and pH variable	183
5.16	Surface plot of 2,3,5-Trimethylpyrazine concentration as affected by the polyphenol concentration and pH variable	184
5.17	Surface plot of 2,3,5,6-Tetramethylpyrazine concentration as affected by the polyphenol concentration and pH variable	184
5.18	Surface plot of standard 2-Methylpyrazine concentration as affected by the polyphenol concentration and pH variable	185
5.19	Surface plot of standard 2,5-Dimethylpyrazine concentration as affected by the polyphenol concentration and pH variable	186
5.20	Surface plot of standard 2,6-Dimethylpyrazine concentration as affected by the polyphenol concentration and pH variable	187
5.21	Surface plot of standard Ethylpyrazine concentration as affected by the polyphenol concentration and pH variable	188
5.22	Surface plot of standard 2,3-Dimethylpyrazine concentration as affected by the polyphenol concentration and pH variable	188
5.23	Surface plot of standard 2,3,5-Trimethylpyrazine concentration as affected by the polyphenol concentration and pH variable	189

5.24	Superimposed contour plot of significant variables for generated alkylpyrazines	195
5.25	Superimposed contour plot of significant variables for standard alkylpyrazines	195

LIST OF ABBREVIATIONS

%	: percentage	mg	: milligram
⁰ C	: degree Celsius	ml	: mililitre
µg	: microgram	mm	: milimetre
µl	: microlitre		
bp	: boiling point		
cm	: centimeter	mM	: millimole
Da	: Dalton	N	: normality
g	: gram	nm	: nanometer
g	: gravity	PVP	: Polyvinylpyrrolidone
GC	: Gas Chromatography	rpm	: rotation per minute
HPLC	: High Performance Liquid Chromatography	RSM	: response surface methodology
hr	: hour	s	: second
i.d.	: internal diameter	UV	: ultraviolet
l	: litre	v	: volume
M	: molarity	w/w	: weight/weight
min	: minute	wt	: weight

CHAPTER 1

GENERAL INTRODUCTION

1.1 Introduction

The cocoa tree, *Theobroma cacao*, L., is usually found in warm and moist climates in areas about 20⁰ latitude north and south of the equator (Wollgast and Anklam, 2000a; 2000b). The Maya and Aztec Indians of central America cultivated the cocoa tree before Columbus discovered America. When Hernando Cortes and his men conquered the Aztecs, they found the Indians, who consider cocoa to be divine and delivered to them directly by God, using the beans to create a drink name chocolatl, an Aztec word meaning acrid or sour (Minifie, 1980, Ensminger *et al.*, 1994, Dillinger *et al.*, 2000). The Swedish botanist, Carolus Linnaeus, must have been familiar with the pleasant flavour of cocoa or chocolate when he classified it in the first edition of his 'Species Plantarum' published in 1753, since the Latin term *theobroma* means 'food of the gods'. This pleasant chocolate flavour builds up during fermentation and roasting of cocoa beans (Woods and Lass, 1985, Ensminger *et al.*, 1994). In Malaysia, there is report stating that cocoa plant was found planted in Sabah in the 1700 after having been brought in by the Spaniards from Latin America to the Philippines and then to Indonesia (Wood and Lass, 1985).

Freshly harvested cocoa beans do not have any chocolate flavour. The unique chocolate flavour derived from cocoa beans will only be produced after series of processing involving fermentation and roasting. A correct fermentation process is crucial in order

to produce suitable cocoa flavour precursors. The flavour precursors produced by fermentation are resulted from the breakdown of protein and carbohydrate into reducing sugars, amino acids and peptides (Foster, 1978).

Reducing sugars, free amino acids and oligopeptides have been identified as important (non volatile) precursors of hundreds of volatile compounds assembling the characteristic of cocoa aroma (Ziegleder and Biehl, 1988; Voigt and Biehl, 1995). Incorrect fermentation process affect the flavour quality badly, this is because unfermented or partly fermented cocoa beans that is further roasted does not develop any chocolate flavour and the resulting flavour was found to be extremely astringent and bitter (Lopez, 1986; Biehl and Voigt, 1996; Puziah *et al.*, 1998a, 1998b).

Sucrose was almost completely hydrolysed to fructose and glucose by invertase present in the beans during the fermentation (Puziah *et al.*, 1998a, 1998b). However, the reducing sugars can also be formed from the hydrolysis of anthocyanins to arabinose and galactose by glycosidase enzyme (Mamot, 1989; Hoskin and Dimick, 1994).

Free amino acids, namely hydrophobic amino acids, are aroma precursors of cocoa beans produces during cocoa fermentation, which will develop into cocoa-specific aroma upon roasting (Barel *et al.*, 1985; Voigt *et al.*, 1994a; 1994b). During cocoa

fermentation, proteolysis, catalyzed by aspartic proteinase and carboxypeptidase, increased the content of free amino acids and oligopeptides (Biehl *et al.*, 1982b; Ziegleder and Biehl, 1988; Puziah *et al.*, 1998a; 1998b). The aspartic proteinase is an endopeptidase that breakdowns proteins at hydrophobic sites to produce hydrophobic amino acids and oligopeptides. The carboxypeptidase is an exopeptidase that breakdowns proteins at the carboxy terminals producing hydrophilic amino acids and oligopeptides (Biehl *et al.*, 1993; Voigt *et al.*, 1993; 1994a).

Roasting of the whole cocoa bean or nib is an essential step to further develop the chocolate flavour, which should already present in the precursors form arising from correct fermentation. Cocoa roasting is a process that depend on time-temperature relationship, where the time can ranges from 5 to 120 min and the temperature ranges from 120 to 150 °C (Wollgast and Anklam, 2000a). All the aroma precursors which include free amino acids, peptides and reducing sugars will involved in heat-induced Maillard non-enzymatic browning reaction to produce subsequent cocoa-specific flavour compounds (Barel *et al.*, 1985; Hoskin and Dimick, 1984a). Maillard non-enzymatic browning is a reaction between the carbonyl group of reducing sugars and a nitrogen source from free, uncharged amine group of amino acids or protein with the loss of one mole of water (Davies *et al.*, 1997; Hofmann and Schieberle, 2000; Jinap *et al.*, 1998; Labuza and Braisier, 1992).