



UNIVERSITI PUTRA MALAYSIA

**STUDIES ON LOCAL ISOLATES OF INFECTIOUS
BURSAL DISEASE VIRUS**

THU ZAR THAN @ HAFIZA HJ HASHIM

SF 995

STUDIES ON LOCAL ISOLATES OF INFECTIOUS BURSAL DISEASE VIRUS

By

THU ZAR THAN @ HAFIZA HJ HASHIM

**Thesis Submitted in Fulfilment of the Requirements for
the Degree of Master of Science in the Faculty of
Veterinary Medicine and Animal Science,
Universiti Pertanian Malaysia**

May 1996



ACKNOWLEDGEMENTS

I am exceedingly grateful and obliged to my chairman, Dr Mohd Hair Bejo for his dedicated efforts, active participation, untiring guidance, stimulating discussion, and constant encouragement throughout the planning and execution of this study, without these helps it could not be completed. His sacrifices and preserverance will ever be remembered.

Sincere thanks are also conveyed to Prof. Dr. Abdul Latiff Ibrahim and Assoc. Prof. Dr. Aini Ideris who provided their advice, helpful discussion and enlightening comments that improved this study.

It pleases me to have this opportunity of expressing my gratitude to Prof. Dr. Tengku Azmi Tengku Ibrahim, Dean of the Faculty of Veterinary Medicine and Animal Science, UPM, for his kind encouragement and granting me to pursue this study. Subsequently, thanks also go to Prof. Dr. Sheikh Omar Abdul Rahman, Head of the Department of Veterinary Pathology and Microbiology, UPM, for his kind support.

I would like to acknowledge my thanks to Dr. Ko Ko Gyi, Dr. Tin Tun Oo, Dr. Min Soe and Daw Yi Yi, University of Animal Husbandry and Veterinary Science, Burma, who initiated in doing this study.

I also acknowledge my deep appreciation and gratitude to the following who have been of great help in doing this thesis: Puan Rodiah Hussein,



Dr. Moonafizad Mohammad, En. Abdul Aziz Bahsir, Dr. Karim Sadun, En. Md. Kamaruddin Awang, En. Abdul Rahim Osman, En. Md. Jamil Samad, En. Noraziman Sulaiman, Pn. Salina Mustapha, Pn. Aminah Jusoh, Mr. Ho Oi Kuan, Mrs. Josephine Low, Pn. Jamilah Abdul Rahman and all other staff members of the Faculty for sharing their technical skills and for their ever available assistances.

To my most beloved parents, U Than and Daw Khin Yi Soe, words are not enough for the sacrifices they have made in laying the foundation since my early life, for their love, full support and constant prayers for me. I tender my appreciation to all my sisters and brothers who motivated and rendered patience and understanding to meet my priorities. It is impossible to list all of my friends' names here but the one who deserves special mention is Moe Moe.



TABLE OF CONTENTS

ACKNOWLEDGEMENTS.....	ii
LIST OF TABLES.....	vii
LIST OF FIGURES.....	viii
LIST OF PLATES.....	ix
LIST OF ABBREVIATIONS.....	xi
ABSTRACT.....	xii
ABSTRAK.....	xv
CHAPTER	
1 INTRODUCTION.....	1
2 LITERATURE REVIEW.....	5
Background.....	5
Infectious Bursal Disease Virus.....	6
Classification.....	6
Morphology.....	6
Resistance to Chemical and Physical Agents.....	7
Propagation.....	8
Transmission.....	9
Susceptibility.....	10
Pathogenesis.....	11
Clinical Signs.....	12
Immunosuppression.....	13
Prevention and Control.....	14
Diagnosis.....	15



3	CHARACTERIZATION OF THE LOCAL ISOLATES OF INFECTIOUS BURSAL DISEASE VIRUS	
	Introduction.....	17
	Materials and Methods.....	19
	Case History.....	19
	Gross Pathology.....	22
	Histopathology.....	22
	Transmission Electron Microscopy.....	22
	Virus Isolation and Identification.....	23
	Results.....	26
	Gross Lesions.....	26
	Histopathology.....	31
	Transmission Electron Microscopy.....	34
	Virus Isolation and Identification.....	34
	Discussion.....	43
4	PATHOGENICITY AND IMMUNOGENICITY STUDIES OF LOCAL ISOLATE OF INFECTIOUS BURSAL DISEASE VIRUS IN SPECIFIC PATHOGEN FREE CHICKENS.	
	Introduction.....	47
	Materials and Methods.....	49
	Experimental Chickens.....	49
	IBDV Inoculum.....	50
	Bursa to Body Weight Ratio.....	51
	Histopathology.....	52
	Transmission Electron Microscopy.....	53
	Virus Isolation and Identification.....	54



Statistical Analysis.....	55
Results.....	56
Clinical Signs.....	56
Bursa to Body Weight Ratio.....	58
Gross Pathology.....	61
Histopathology.....	67
Transmission Electron Microscopy.....	74
Virus Isolation and Identification.....	77
Discussion.....	83
5 GENERAL DISCUSSION AND CONCLUSION.....	87
BIBLIOGRAPHY.....	91
PUBLICATIONS.....	103
APPENDICES.....	104
VITA.....	116



LISTS OF TABLES

Table		Page
1	Case history of seven local isolates of infectious bursal disease virus.	20
2	Mean body weight, bursal weight and bursal to body weight ratio of chickens inoculated with IBDV local isolates.	41
3	Body weight, bursal weight and bursal to body weight ratio of the control and infectious bursal disease groups.	59
4	The occurrences of haemorrhages in the bursa of Fabricius, skeletal muscles and at the junction of proventriculus and gizzard of the sacrificed and dead chickens at days 2 and 3 post inoculation of IBDV.	64
5	Histological lesion scoring of the bursa of Fabricius in the (a) control and (b) IBD groups throughout the trial.	68
6	Antibody titre of infectious bursal disease and Newcastle disease in the control and IBD groups.	81
7	Fifty Percent Tissue Culture Infective Dose (TCID ₅₀ /ml).	113
8	Fifty Percent Embryo Infective Dose (EID ₅₀ /ml).	114
9	Fifty Percent Embryo Lethal Dose (ELD ₅₀ /ml).	115



LIST OF FIGURES

Figure		Page
1	(a) Body and (b) bursal weight of the control and IBD groups throughout the trial.	60
2	The ratio of bursa to body weight of the control and IBD groups throughout the trial.	63
3	IBD antibody titre of the control and IBD groups throughout the trial.	82



LIST OF PLATES

Plate		Page
1	IBD field outbreak, UPM 93273. (a) Haemorrhages of the leg muscles. (b) Haemorrhages of the bursa of Fabricius. Yellowish exudate was present in the bursal lumen.	28
2	IBD field outbreak, UPM 94274. (a) Haemorrhages of the breast muscles. (b) Haemorrhages at the junction of proventriculus and gizzard.	29
3	IBD field outbreak. (a) UPM 94274, enlargement and atrophy of the bursa of Fabricius. (b) UPM 94283, haemorrhages at the bursa of Fabricius.	30
4	IBD field outbreak. (a) UPM 93273, necrosis and haemorrhages of the bursal follicles and infiltration of inflammatory cells in the interstitial connective tissue (x4). (b) UPM 94230, severe necrosis of bursal follicles and lymphoid cells depletion with haemorrhages, and infiltration of inflammatory cells in the interstitial connective tissue (x4).	33
5	IBD field outbreak, UPM 94274, IBDV particles in crystalloid arrays measured about 50-55 nm in diameter either with or without membrane presence in (a) the macrophage (x 115 500), and (b) the necrotic cell (x 32 250).	36
6	IBD field outbreak, embryonated chicken egg. (a) Control embryo. (b) Dead embryo with severe haemorrhages and oedema at day 3 post inoculation with UPM 93273.	37
7	IBDV (UPM 93273) chick inoculation. (a) Atrophied bursa of Fabricius covered with transudates (arrow) and the control. (b) Severe lymphoid cells necrosis and presence of cystic follicles (x4).	42
8	IBD in SPF chickens. (a) Severe depression, ruffled feathers and lateral recumbency with coarse tremor shortly before death at day 3 post inoculation. (b) Watery diarrhoea at day 10 post inoculation.	57



9	IBD in SPF chickens. Day 2 post inoculation. (a) Haemorrhages of the leg muscles. (b) Haemorrhagic line at the junction of proventriculus and gizzard.	65
10	IBD in SPF chickens. (a) Day 2 post inoculation. haemorrhages and enlargement of the bursa of Fabricius. (b) Day 4 post inoculation, atrophy (arrow) of the bursa of Fabricius and the control.	66
11	IBD in SPF chickens. Day 2 post inoculation. (a) The bursa of Fabricius from control bird (Score 1) (x4). (b) Necrosis of the lymphoid follicles and infiltration of inflammatory cells in the interstitial connective tissue (Score 5) (x4).	69
12	IBD in SPF chickens. (a) Day 6 post inoculation, lymphoid follicles undergone atrophy and vacuolation, and interstitial connective tissue were infiltrated with inflammatory cells (Score 5) (x4). (b) Day 10 post inoculation, severe atrophy and vacuolation of follicles infiltrated with inflammatory cells and fibrous tissues (Score 5) (x4).	72
13	IBD in SPF chickens. (a) Day 21 post inoculation, atrophic bursa of Fabricius with recovery of the follicles (Score 4) (x4). (b) Day 3 post inoculation, severe haemorrhages and necrosis of the lymphoid follicles of the dead bird (Score 5) (x10).	73
14	IBD in SPF chickens. Day 1 post inoculation. (a) Normal lymphocytes seen in control bird (x 7 500). (b) Undistinguishable IBDV particles were observed as highly electron dense materials in the cytoplasm of lymphoid cell (x 15 000).	75
15	IBDV in SPF chickens. (a) Day 2 post inoculation, a cluster of IBDV particles (arrow) and undistinguishable electron dense IBDV particles observed in the cytoplasm of lymphoid cell (x 24 375). (b) Day 3 post inoculation, membrane enclosed mature IBDV particles in crystalline array clearly definable in lymphoid cell (x 30 000).	76
16	IBD in SPF embryonated chicken eggs Day 3 post inoculation. (a) Haemorrhages of chorioallantoic membrane, and (b) haemorrhages and oedema of the dead embryo.	78



LIST OF ABBREVIATIONS

BF	- Bursa of Fabricius
CAM	- Chorioallantoic Membrane
CEF	- Chicken Embryo Fibroblast
CPE	- Cytopathic Effect
EM	- Electron Microscope
EID ₅₀	- Embryo Infective Dose 50%
ELISA	- Enzyme Linked Immunosorbent Assay
HA	- Haemagglutination
HE	- Hematoxylin and Eosin
HI	- Haemagglutination-Inhibition
IBD	- Infectious Bursal Disease
IBDV	- Infectious Bursal Disease Virus
ND	- Newcastle Disease
NDV	- Newcastle Disease Virus
PBS	- Phosphate Buffer Saline
pH	- Hydrogen-ion Concentration
pi	- Post Inoculation
RNA	- Ribonucleic Acid
TCID ₅₀	- Fifty percent Tissue Culture Infective Dose
TEM	- Transmission Electron Microscopy
UPM	- Universiti Pertanian Malaysia



Abstract of the thesis presented to the Senate of
Universiti Pertanian Malaysia in fulfilment of the requirements for
the degree of Master of Science

**STUDIES ON LOCAL ISOLATES OF
INFECTIOUS BURSAL DISEASE VIRUS**

By

THU ZAR THAN @ HAFIZA HJ HASHIM

May 1996

Chairman: Dr. Mohd Hair Bin Bejo

Faculty: Veterinary Medicine and Animal Science

Infectious bursal disease (IBD) is one of the most important viral diseases in chickens. IBD virus (IBDV) from seven local field outbreaks in layer, broiler and village chickens were isolated, propagated and identified. The pathogenicity and immunogenicity of one of the selected isolates were determined in specific pathogen free (SPF) chickens.

The study showed that the IBDV can cause sudden onset of mortality ranging from 15% to 90% during the outbreaks. It occurred in both the vaccinated and non-vaccinated chickens with the age group ranging from 18 days to 14 weeks. Layers appeared to be more susceptible to the virus than the broilers, whilst the highest mortality (90%) was observed in village chickens.

Haemorrhages of the bursa of Fabricius, muscles and mucosal layers at the junction of the proventriculus and gizzard were the typical gross lesions



caused by the virus. The bursal lesions vary from moderate enlargement and oedematous to severe atrophy. Histologically, the enlarged bursa showed acute necrotizing bursitis, whilst chronic necrotizing bursitis was seen in the atrophic organ. The virus particles were detected in the cytoplasm of the lymphoid cells, macrophages and necrotic cells of the bursa of Fabricius under transmission electron microscopy (TEM). Inoculation of the IBDV isolates obtained from the bursa of Fabricius into embryonated chicken eggs caused embryonic death, haemorrhages, oedema and hepatic necrosis. The lesions were more severe in SPF eggs than the commercial eggs. Inoculation of the virus into susceptible chickens caused some variation in the mortality and lesions. All isolates of the IBDV from the outbreaks showed distinct curve of agar gel diffusion precipitation lines between the positive sample and reference serum against IBDV.

Studies on one of the isolates (UPM 93273), inoculated orally at a dose of $10^{6.8} \text{EID}_{50}$ per ml in 28-day-old SPF chickens showed that the virus was highly pathogenic and immunogenic. Sudden onset of mortality (60%) to about four times than the field outbreak occurred at days 2 and 3 post inoculation. The gross and histological lesions of the affected chickens were consistent to those of the field outbreak except that more severe lesions were observed in the SPF chickens. Acute necrotizing bursitis was observed during the early stage of the infection at days 2 to 4, while at the late stage, chronic necrotizing bursitis was recorded. The virus was detected from the bursa of Fabricius from days 1 to 10 and days 1 to 12 using embryonated chicken eggs and TEM respectively. Antibody titre against the virus was first detected at day 6 post inoculation (1271 ± 75) and reached the highest level by day 21 (4490 ± 735).



It was concluded that the study has successfully isolated and propagated a highly pathogenic strains of serotype 1 IBDV from seven field outbreaks in the country. Studies on one of the isolates demonstrated that the virus was also highly immunogenic, and thus, it can be useful as seed virus in the production of local IBD vaccine, which is safe and efficient in the prevention and control of the disease in the country.



Abstrak tesis yang dikemukakan kepada Senat
Universiti Pertanian Malaysia bagi memenuhi syarat untuk
Ijazah Master Sains

**KAJIAN KE ATAS ISOLAT VIRUS TEMPATAN
PENYAKIT BURSA BERJANGKIT**

Oleh

THU ZAR THAN @ HAFIZA HJ HASHIM

Mei 1996

Pengerusi: Dr. Mohd Hair Bin Bejo

Fakulti: Kedokteran Veterinar dan Sains Peternakan

Penyakit bursa berjangkit (IBD) adalah merupakan salah satu penyakit virus terpenting pada ayam. Virus IBD (IBDV) daripada tujuh wabak penyakit tempatan yang berlaku pada ayam penelur, pedaging dan ayam kampung diasingkan, dibiakkan dan dikenalpasti. Kepatogenan dan keimmunogenan satu isolat terpilih dikenalpasti dalam patogen khusus bebas (SPF) ayam.

Kajian menunjukkan IBDV boleh menyebabkan mortaliti secara tiba-tiba dalam lingkungan 15% hingga 90% semasa wabak penyakit. Ia berlaku pada ayam yang telah dan belum menerima pemvaksinan, berumur di antara 18 hari hingga 14 minggu. Ayam penelur didapati lebih mudah dijangkiti virus daripada ayam pedaging, manakala mortaliti (90%) tertinggi terdapat pada ayam kampung.

Pendarahan pada bursa Fabricius, otot, lapisan mukosa diantara proventikulus dan hempedal adalah lesi matakasar khusus jangkitan IBDV.



Lesi bursa berbeza dari sederhana membesar dan edema kepada atrofi yang teruk. Pemeriksaan histologi pada bursa yang membesar menunjukkan bursitis nekrosis akut, manakala bursitis nekrosis kronik dilihat dalam atrofik bursa. Butiran IBDV didapati dalam sitoplasma sel-sel limfoid, makrofaj dan sel yang mengalami nekrosis dalam bursa Fabricius melalui mikroskop elektron pancaran (TEM). Penginokulatan IBDV isolat yang diperolehi dari bursa Fabricius ke dalam telur ayam berembrio menyebabkan kematian, pendarahan, edema dan nekrosis hati pada embrio. Lesi embrio di telur SPF lebih teruk dibandingkan dengan di telur komersial. Inokulasi IBDV ke dalam ayam menunjukkan sedikit perbezaan mortaliti dan lesi di antara isolat. Semua isolat IBDV tempatan menunjukkan keluk jelas ujian sebaran pemendakan gel agar (AGPT) di antara sampel positif dan sampel serum rujukan IBDV.

Kajian pada satu isolat (UPM 93273) yang diinokulat melalui mulut pada dos $10^{6.8}$ EID₅₀ setiap ml pada ayam SPF berumur 28 hari menunjukkan IBDV isolat tersebut sangat patogenik dan imunogenik. Mortaliti (60%) secara tiba-tiba sehingga mencapai empat kali ganda dari kejadian wabak di ladang berlaku pada hari kedua dan ketiga post inokulasi. Lesi matakasar dan histologi pada ayam yang terlibat adalah seperti lesi sewaktu wabak penyakit, kecuali lesi yang lebih teruk terdapat dalam ayam SPF. Bursitis nekrosis akut dilihat di peringkat awal jangkitan virus pada hari kedua hingga keempat, manakala di peringkat akhir, bursitis nekrosis kronik berlaku. Virus IBD dapat dikesan di bursa Fabricius dari hari pertama sehingga hari kedua dengan menggunakan TEM dan hari pertama sehingga hari

kesepuluh melalui telur ayam berembrio. Titre antibodi IBDV mula dikesan pada hari keenam post inokulasi (1271 ± 75) dan mencapai tahap tertinggi pada hari ke 21 (4490 ± 735).

Kesimpulannya, kajian ini telah berjaya mengasing dan membiakkan serotaip 1 IBDV strain yang sangat patogenik daripada tujuh wabak penyakit yang berlaku di negara ini. Kajian pada salah satu isolat tersebut menunjukkan ianya juga sangat imunogenik. Isolat ini mungkin boleh digunakan sebagai benih virus dalam pengeluaran vaksin IBD tempatan yang selamat dan dapat memberi perlindungan sepenuhnya dalam pencegahan dan kawalan wabak IBD di negara ini.

CHAPTER 1

INTRODUCTION

Infectious bursal disease (IBD) or Gumboro disease, an extremely important viral disease in poultry industry worldwide, was first reported in the district of Delaware in the U.S.A in 1957 (Cosgrove, 1962). However, the clinical outbreaks of the disease was only reported in Malaysia in 1991 (Hair-Bejo, 1992; Loganathan *et al.*, 1992). Since then, acute IBD outbreaks were diagnosed in both layer and broiler chickens in the country. The disease causes intense economic losses due to sudden onset of high mortality and high condemnation of the carcasses or resulting from high mortality associated with secondary infections and failure to the vaccination programmes against other highly virulent diseases (Hair-Bejo, 1994).

Infectious bursal disease virus (IBDV) is a bisegmented double-stranded RNA (dsRNA) virus with icosahedral symmetry and a diameter of about 50 to 55 nm (Muller *et al.*, 1979b; Dobos *et al.*, 1979; Becht, 1980; Jackwood *et al.*, 1984). Two distinct serotypes of IBDV designated as serotype 1 within that occurred antigenic variants, and serotype 2 were recognized (McFerran *et al.*, 1980; Jackwood *et al.*, 1982; McNulty and Saif, 1988; Wood *et al.*, 1988). Only serotype 1 IBDVs have been known to cause naturally occurring disease in chickens. It occurs as a clinical or subclinical IBDV infection (Rosenberger and Gelb, 1978). The clinical IBD usually occurs in chickens



infected with either standard or highly virulent strains of serotype 1 IBDV in birds more than 2-3 weeks old (Cosgrove, 1962; Parkhurst, 1962; Winterfield and Hitchner, 1964; Luthgen, 1969), whilst infection in the younger age could lead to subclinical IBD. The subclinical IBD is also observed in chicks infected with variant strains of the virus (Snyder, 1990).

IBDV replicates primarily in the lymphoid cells of the bursa of Fabricius (Ismail *et al.*, 1987; Okoye and Uzoukwu, 1990). It induces severe necrosis of the lymphocytes and atrophy of the organ (Hirai *et al.*, 1974). The availability of the large number of bursal cells (B cells) has been shown to be an essential factor in development of IBD infection (Ismail *et al.*, 1987).

Isolation of IBDV from the field outbreaks may be difficult, although the virus was successfully isolated in embryonated chicken eggs (Winterfield *et al.*, 1962) with the chorioallantoic membrane (CAM) route performed the best for the virus isolation. Embryo (Hitchner, 1978). The virus also has been isolated using chicken embryo fibroblast cell cultures (Lukert and Davis, 1974; McNulty *et al.*, 1979; Lee and Lukert, 1986), bursa B-cell derived lymphoblastoid cell line (Hirai and Calnek, 1979) and BGM-70 cell line (Jackwood *et al.*, 1987).

IBDV is very stable to chemical and physical agents and it can remain for long in a contaminated environment (Benton, 1967b). Once the infection set

up on a farm, it recurs in several subsequent flocks despite the most thorough cleansing and disinfection (Winterfield and Hitchner, 1964; Edgar and Cho, 1973; Hair-Bejo *et al.*, 1995a). There is no therapeutic measures for IBDV infection and the disease can only be prevented by proper immunization programmes in both the parent stock and their progeny.

To date, about 46 types of IBD vaccines were imported for use in West Malaysia (Chin, 1993). Several vaccination programmes against the disease has been carried out, but their success are unpredictable (Hair-Bejo, 1993). Outbreaks of IBDV continued to occur in both the vaccinated and non-vaccinated chickens in the country (Hair-Bejo, 1994). The safety and efficacy of the vaccines under local conditions is little understood. Studies on some of the vaccines showed that the IBD vaccines can induce lesion similar to those observed in the field infection (Hair-Bejo *et al.*, 1994). The failure of IBD vaccines to induce antibody against IBDV was also reported (Hair-Bejo *et al.*, 1995c). Furthermore, the emergence of different strains of IBDV in the country may complicate the immunization programme of the disease.

Thus, it is an urgent need to study on local isolates of IBDV for the prevention and control of the disease in the country. The highly pathogenic IBDV isolates can be used for challenge virus in the study of the safety and efficacy of IBD vaccines, whilst the highly immunogenic viruses is needed for virus seed in the production of local IBD vaccine which is safe and provides

full protection against the IBDV field challenge. Therefore, the objectives of this study are:

1. to isolate propagate and characterize the recent field isolates of IBDV, and
2. to determine the pathogenicity and immunogenicity of a selected IBDV local field isolate.

CHAPTER 2

LITERATURE REVIEW

Background

Infectious bursal disease (IBD) was first reported in a broiler flock in the small village, Delaware, USA in 1957 (Cosgrove, 1962). The common synonyms of the disease are avian nephrosis (Cosgrove, 1962), Gumboro disease (Faragher, 1972), infectious bursitis (Ibragimov, 1976) and avian infectious bursitis (Rodon, 1982). Since the first outbreak of IBD, the disease was then reported from most major poultry producing areas around the world such as in Europe (Faragher, 1972), Australia (Firth, 1974), India (Mohanty *et al.*, 1971) and Japan (Shimizu *et al.*, 1971).

Serious outbreaks of clinical IBD with high mortality, up to 90% due to highly virulent strains of serotype 1 IBD virus (IBDV) occurred in late 1980's throughout Europe (Chettle *et al.*, 1989; Van den Berg *et al.*, 1991). The disease had spread worldwide and was described in Asia in 1990's (Nunoya *et al.*, 1992) including Malaysia in 1991 (Hair-Bejo, 1992, Loganathan *et al.*, 1992). It occurred in both the vaccinated and non-vaccinated chickens and the mortality increased with the presence of concurrent infections (Hair-Bejo, 1994).

Infectious Bursal Disease Virus

Classification

Infectious bursal disease virus was first classified as a picornavirus (Cho and Edgar, 1969). To date, IBDV is referred as a Diplorna virus, a member of the newly established genus birna virus under family Birnaviridae with a genome consisting of a bisegmented double-stranded RNA (dsRNA) (Dobo *et al.*, 1979; Brown, 1984). Two serotypes of IBDV, serotypes 1 and 2, have been recognized. Serotype 1 IBDV is pathogenic to chicken whereas serotype 2 virus is isolated from turkey (McFerran *et al.*, 1980; Jackwood *et al.*, 1982; Cummings *et al.*, 1986) and non pathogenic in chickens (Ismail *et al.*, 1987). The strains of serotype 1 IBDV has been tentatively classified into three groups based on the pathogenicity and antigenicity; very virulent strains of serotype 1 IBDV (vvIBDV) (Chettle *et al.*, 1989; Van den Berg *et al.*, 1991; Nunoya *et al.*, 1992; Tsukamoto *et al.*, 1992), standard strains of serotype 1 IBDV (stIBDV) (Hirai *et al.*, 1973; Sharma and Lee, 1984; Chettle *et al.*, 1989; Craft *et al.*, 1990), and variant strains of serotype 1 IBDV (vaIBDV) (Saif, 1984; Rosenberger and Cloud, 1985; Sharma *et al.*, 1989; Craft *et al.*, 1990; Snyder, 1990).

Morphology

Negatively stained IBDV particles examined under transmission electron microscopy (TEM) demonstrated that the virus is a non-enveloped, single capsid



shell of icosahedral symmetry varying from 55 to 65 nm (Hirai and Shimakura, 1974; Hirai *et al.*, 1979; Dobo *et al.*, 1979; Brown, 1984). The virus varies somewhat in appearance depending upon the purification procedure employed. Local pH and ionic concentrations affect the state of hydration of the virion and this has the morphological consequences described (Harkness *et al.*, 1975).

IBDV particles arrayed in a crystalline pattern measured about 50-55 nm was detected in the cytoplasm of lymphoid cells, macrophages, heterophils and reticular epithelial cells of the bursa of Fabricius examined directly under TEM (Kaufer and Weiss, 1976; Hair-Bejo, 1993). Immature forms of IBDV was first found in the cytoplasmic inclusion bodies at 24 hours post inoculation, whilst relatively large amounts of mature viruses are present in macrophage inclusions by day 3 (Kaufer and Weiss, 1976).

Buoyant density of IBDV particles in cesium chloride (CsCl) gradients has been reported to be 1.33 g/cm³ (Nick *et al.*, 1976; Becht, 1980; Muller and Becht, 1982; Fahey *et al.*, 1985). A higher buoyant density of 1.34 g/ml (Hirai *et al.*, 1979) and lower densities of 1.31 g/ml (Pattison *et al.*, 1975; Todd and McNulty, 1979) and 1.32 g/ml (Jackwood *et al.*, 1982, 1984) have also been reported. Immature (incomplete) virus particles have buoyant densities lower than 1.33 g/ml in CsCl gradients.

Resistance to Chemical and Physical Agents

IBDV is relatively refractory to heat, ultraviolet irradiation and photodynamic inactivation (Petek *et al.*, 1973). It is inactivated at pH 12.0, but

