



UNIVERSITI PUTRA MALAYSIA

**BACTERIAL AND SHORT-CHAIN FATTY ACID PROFILING OF FAECAL
SAMPLES**

NUR HUDA BINTI FAUJAN

FSTM 2011 18

**BACTERIAL AND SHORT-CHAIN FATTY ACID PROFILING
OF FAECAL SAMPLES**



NUR HUDA BINTI FAUJAN

**DOCTOR OF PHILOSOPHY
UNIVERSITI PUTRA MALAYSIA**

2011

**BACTERIAL AND SHORT-CHAIN FATTY ACID PROFILING OF FAECAL
SAMPLES**

by

NUR HUDA BINTI FAUJAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirement for the degree of Doctor of Philosophy**

December 2011



This thesis is dedicated to

- *teachers who teach people for a better tomorrow.*
- *my parents, my husband and my daughter, all my siblings and family-in-laws for their moral support, love, patience and understand*

Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

BACTERIAL AND SHORT-CHAIN FATTY ACID PROFILING OF FAECAL SAMPLES

By

NUR HUDA BINTI FAUJAN

December 2011

Chairman : Professor Fatimah Abu Bakar, PhD

Faculty : Food Science and Technology

Human gastrointestinal tract consists of an extremely complex microbiota mainly the anaerobic organisms. Many studies have been done to investigate the human intestinal microbiota. The use of molecular techniques based on DNA for microbial studies is of interest due its reliability, rapid and sensitivity method. Real-time polymerase chain reaction (PCR) has become one of the most widely used methods and an important tool for analysis of bacterial quantification as well as for detection and identification of bacterial species.

In this study, real-time PCR has been used to investigate and quantify bacterial species in human faecal samples. Prior to the analysis, the extraction of DNA from human faeces had been optimised to produce a high quality and yield of DNA. Based on this purpose, DNA from faecal samples was extracted using QIAamp[®] DNA Stool Mini Kit. However, to give maximum DNA yields, four different procedures of extraction using this kit have been modified by adding 1 mL of 1x PBS buffer in

preheating step or 600 μL of 1x PBS buffer incorporated with InhibitEx (inhibitor-adsorption) tablet. Two different amounts of 20 mg/mL proteinase K (15 μL or 25 μL) were used to digest protein and remove protein contamination. From this study, results showed that the addition of 600 μL of 1x PBS buffer incorporated with the InhibitEX tablet increase removal of inhibitors in faecal samples. Addition of 25 μL of proteinase K also produced higher DNA yield compared to the addition of 15 μL of proteinase K for protein digestion. Optimisation of DNA extraction is a crucial factor in molecular analysis of faecal samples due to the presence of many inhibitors in faecal samples that can interfere with the accuracy of real-time PCR analysis.

Next, specific primer based on 16S rRNA gene was used to analyse and amplify bacterial DNA from faecal samples. The specificity of all primers used were tested by conventional polymerase chain reaction (PCR). Results showed that all primers were highly specific with the target bacteria and amplified DNA with the expected PCR products size. In order to apply real-time PCR for quantitative analysis, the real-time PCR conditions have been optimised to provide an accurate analysis. Standard calibration curve for quantitative analysis was constructed using plasmid DNA. All standard calibration curves plotted from plasmid DNA produced slope between -3.84 and -3.29, which highlighted the efficiency of reaction assay between 81.8% and 101.4%. Efficiency of real-time PCR assay is a crucial factor in order to provide accurate and precise result for quantification.

In order to apply real-time PCR for analysis of intestinal microbiota, faecal samples were obtained from subjects with inflammatory bowel disease (IBD) and also healthy condition. From this study, results showed that *Bacteroides* and *B. fragilis* in faecal samples of IBD patients were significantly higher ($P < 0.05$) than that of healthy persons. The mean for *Bacteroides* in faecal samples of IBD patients and healthy persons were $10.9 \log_{10}$ cells/g and $10.3 \log_{10}$ cells/g, respectively. The mean for *B. fragilis* in faecal samples of IBD patients and healthy person were $5.9 \log_{10}$ cells/g and $4.6 \log_{10}$ cells/g, respectively. *B. fragilis* was not even detected in some faecal samples of healthy persons. Other bacteria such as *Desulfovibrios*, *Clostridium coccooides*, *Escherichia coli* and *Enterococcus faecalis* did not show any significant difference ($P > 0.05$) in faecal samples between two groups of subjects.

The concentration of short-chain fatty acids (SCFAs) of faecal samples in both groups was also analysed using high performance liquid chromatography (HPLC) method. The SCFAs in human colon are produced by fermentation of carbohydrates, proteins and peptides by human colonic bacteria. Results showed that acetic, butyric and propionic acids concentration was lower in faecal samples of IBD patients as compared to that of healthy persons. However, only butyric and propionic acids were significantly lower ($P < 0.05$) in faecal samples of IBD patients as compared to that of healthy persons. The mean concentration of butyric acid in faecal samples of IBD patients was $66.2 \mu\text{mol/g}$ and in healthy person was $175.0 \mu\text{mol/g}$. The mean concentration of propionic acid in faecal samples of IBD patients was $52.8 \mu\text{mol/g}$ and in healthy persons was $93.3 \mu\text{mol/g}$. Results also showed that concentration of lactic acid in faecal samples of IBD patients was higher than that of healthy persons. The concentration of formic, pyruvic and isobutyric acids in faecal samples were also

analysed. In faecal samples of IBD patients, the concentration of pyruvic and isobutyric acids were higher than that of healthy person but not significant ($P>0.05$).

In conclusion, DNA extraction from faecal samples using QIAamp[®] DNA Stool Mini Kit with addition of 600 mL of 1x PBS buffer in removal inhibitors step and higher amount of 20 mg/mL proteinase K in eliminating contaminated protein resulted in increased quality and yield of faecal DNA. By using real-time PCR, this study showed that the composition of certain bacterial species in IBD faecal samples differed from that in faecal samples of healthy persons. The concentration of certain SCFAs also changed in faecal samples of IBD patients and which might be attributed to difference in certain bacterial species. Furthermore, this study also showed that real-time PCR assay was an efficient and rapid method for studying bacterial profile in faecal samples.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan Ijazah Doktor Falsafah

**PEMPROFILAN BAKTERIA DAN ASID LEMAK BERANTAI-PENDEK
DALAM SAMPEL NAJIS**

Oleh

NUR HUDA BINTI FAUJAN

Disember 2011

Pengerusi : Profesor Fatimah Abu Bakar, PhD

Fakulti : Sains dan Teknologi Makanan

Saluran pencernaan makanan manusia mengandungi mikrobiota yang sangat kompleks terutamanya daripada organisma anaerob. Banyak kajian telah dilakukan untuk menyelidik mikrobiota usus manusia. Penggunaan teknik molekular yang berdasarkan penggunaan DNA untuk kajian mikrob telah menjadi perhatian kerana keputusan yang lebih dipercayai, cepat dan sensitif. Kaedah tindak balas rantaian polimerase (PCR) masa-nyata telah menjadi salah satu kaedah yang paling banyak digunakan dengan meluas dan menjadi satu alat yang penting untuk analisis pengiraan bakteria serta pengesanan dan pengenalpastian bakteria.

Dalam kajian ini, PCR masa-nyata telah digunakan untuk menyelidik dan mengira spesies bakteria dalam sampel najis manusia. Sebelum analisis dijalankan, pengekstrakan DNA daripada najis manusia telah dioptimakan untuk menghasilkan perolehan DNA yang banyak dan berkualiti. Berdasarkan tujuan ini, DNA telah diekstrak menggunakan Kit Mini QIAmp[®] DNA Stool. Bagaimanapun, untuk

memberikan perolehan DNA yang maksimum, empat prosedur pengekstrakan yang berbeza telah diubahsuai semasa penggunaan kit ini dengan penambahan 1 mL larutan penimbal PBS pada langkah prapemanasan sampel atau dengan menambah 600 μ L larutan penimbal 1x PBS yang digabungkan dengan tablet InhibitEX (penjerapan-perencat). Dua jumlah 20 mg/mL proteinase K yang berbeza (15 μ L atau 25 μ L) juga telah digunakan untuk mencernakan protein dan menyingkirkan pencemaran protein. Dari kajian ini, keputusan menunjukkan bahawa penambahan 600 μ L larutan penimbal 1x PBS yang digabungkan dengan tablet InhibitEX telah meningkatkan penyingkiran bahan perencat dalam sampel najis. Penambahan 25 μ L proteinase K juga telah menghasilkan DNA yang lebih banyak berbanding dengan penambahan hanya 15 μ L proteinase K bagi pencernaan protein. Pengoptimuman pengekstrakan DNA merupakan faktor yang sangat penting dalam analisis molekular sampel najis kerana kehadiran banyak bahan perencat dalam sampel najis yang boleh mengganggu ketepatan analisis PCR masa-nyata.

Seterusnya, primer spesifik yang berdasarkan gen 16S rRNA telah digunakan untuk menganalisis dan mengandakan DNA bakteria daripada sampel najis. Kespesifikan semua primer yang digunakan telah diuji melalui PCR konvensional. Keputusan kajian menunjukkan bahawa semua primer adalah spesifik dengan bakteria yang disasarkan dan mengandakan DNA dengan saiz produk yang telah dijangkakan. Untuk mengaplikasikan penggunaan PCR masa-nyata sebagai analisis kuantitatif, keadaan PCR masa-nyata telah dioptimumkan untuk memberikan analisis yang tepat. Lengkok kalibrasi piawai yang digunakan untuk analisis kuantitatif telah dihasilkan dengan menggunakan DNA plasmid. Semua lengkok kalibrasi piawai yang telah diplot daripada DNA plasmid menghasilkan kecerunan diantara -3.84 hingga -3.29,

yang mana ini telah memberikan kecekapan asai tindak balas di antara 81.8% hingga 101.4%. Kecekapan asai tindak balas rantaian polymerase masa-nyata merupakan faktor yang sangat penting untuk memberikan keputusan pengiraan yang tepat dan jitu.

Untuk mengaplikasikan penggunaan PCR masa-nyata untuk analisis mikrobiota usus, sampel najis telah diperolehi daripada subjek dengan penyakit radang usus (IBD) dan juga yang sihat. Dari kajian ini, hasil kajian menunjukkan bahawa *Bacteroides* dan *B. fragilis* dalam sampel najis pesakit IBD adalah lebih tinggi secara signifikan ($P < 0.05$) daripada sampel najis manusia sihat. Purata bacaan *Bacteroides* dalam sampel najis pesakit IBD dan manusia sihat adalah $10.9 \log_{10}$ cells/g dan $10.3 \log_{10}$ cells/g. Purata bacaan *B. fragilis* dalam sampel najis pesakit IBD dan manusia sihat adalah masing-masing $5.9 \log_{10}$ cells/g dan $4.6 \log_{10}$ cells/g. Pada beberapa sampel najis manusia sihat, *B. fragilis* juga tidak dikesan. Bakteria lain seperti *Desulfovibrios*, *Clostridium coccooides*, *Escherichia coli* dan *Enterococcus faecalis* tidak menunjukkan perbezaan yang signifikan ($P > 0.05$) antara dua kumpulan subjek.

Kepekatan asid lemak rantai pendek (SCFAs) dalam sampel najis pada kedua-dua kumpulan juga telah dianalisis menggunakan kaedah kromatografi cecair berprestasi tinggi (HPLC). SCFAs dalam kolon manusia adalah merupakan produk hasil penapaian karbohidrat, protein dan peptida oleh bakteria usus manusia. Hasil kajian menunjukkan bahawa kandungan asid asetik, asid butirik dan asid propionik pada sampel najis pesakit IBD adalah lebih rendah jika dibandingkan dengan sampel najis manusia sihat. Namun, hanya kepekatan asid butirik dan asid propionik sahaja yang lebih rendah secara signifikan ($P < 0.05$) dalam sampel najis pesakit IBD berbanding

dalam najis manusia sihat. Purata kepekatan asid butirik dalam sampel najis pesakit IBD adalah 66.2 $\mu\text{mol/g}$ manakala pada manusia sihat adalah 175.0 $\mu\text{mol/g}$. Purata kepekatan asid propionik pada najis pesakit IBD adalah 52.8 $\mu\text{mol/g}$ manakala pada manusia sihat adalah 93.3 $\mu\text{mol/g}$. Keputusan juga menunjukkan bahawa kepekatan asid laktik dalam sampel najis pesakit IBD lebih tinggi daripada dalam sampel najis manusia sihat. Kepekatan asid formik, asid piruvik dan asid isobutirik dalam sampel najis juga telah dianalisis. Dalam sampel najis pesakit IBD, kepekatan asid piruvik dan asid isobutirik adalah lebih tinggi daripada manusia sihat tetapi perbezaan antara dua sampel ini tidak berbeza secara signifikan ($P>0.05$).

Sebagai kesimpulan, pengestrakan DNA dari sampel najis menggunakan Kit Mini QIAmp[®] DNA Stool dengan penambahan 600 mL larutan penimbang 1x PBS pada langkah penyingkiran bahan perencat dan penggunaan jumlah 20 mg/mL proteinase K yang tinggi telah menghasilkan DNA yang lebih berkualiti dan banyak. Dengan menggunakan PCR masa-nyata, kajian ini telah menunjukkan bahawa komposisi bakteria tertentu dalam sampel najis pesakit IBD berbeza berbanding dalam sampel najis manusia sihat. Kepekatan SCFAs tertentu juga telah berubah dalam sampel najis pesakit IBD dan menunjukkan ada perkaitan dengan perubahan komposisi beberapa spesies bakteria tertentu. Seterusnya, kajian ini juga telah menunjukkan bahawa penggunaan kaedah asai PCR masa-nyata dalam mengkaji pemprofilan bakteria dalam sampel najis adalah cekap dan cepat.

ACKNOWLEDGEMENT

In the name of Allah, Most Gracious, Most Merciful, Alhamdulillah I have completed this thesis.

It is unimaginable that an academic effort of this magnitude could successfully come to fruition without the help of others. The participation and contribution of the individuals and institutions throughout the planning and completion of this thesis are gratefully appreciated.

First of all, I am deeply thankful to my university, Universiti Sains Islam Malaysia, Nilai for giving me a scholarship under tutor scheme to pursue my study in Doctor of Philosophy. My sincere thanks are also extended to the Ministry of Science, Technology and Innovation (MOSTI) for the financial support through Science Fund grant (Project Number: 02-01-04-SF0078) for this research, and Universiti Putra Malaysia for giving me place to pursue my study.

Next, I extend my heartfelt gratitude to my supervisor Professor Dr. Fatimah binti Abu Bakar for her guidance, assistance and encouragement with patience during this research. My deepest gratitude also goes to Professor Dr. Mohd Yazid bin Abdul Manap, Associate Professor Dr. Shuhaimi bin Mustafa, and Associate Professor Dr. Loong Yik Yee for their huge contribution in my supervisory committee and giving me invaluable guidance and academic advices throughout the course of my study here.

Also special thanks to all laboratory assistants especially Mr. Arifen bin Kamarudzaman in Basic Biotechnology Laboratory and Mr. Abdul Halim bin Abdul Rahman in HPLC room. Appreciation also goes to my friends, Muhammad Anas Othaman, Dr. Khalilah Abdul Khalil, Raba'atun Adawiyah Shamsuddin, Mohd Hamim Hamdan Mustafa, Mohd Yamin Shaari, Dr. Norlelawati Arifin, Dr. Nor Hayati Ibrahim, Dr. Marina Abdul Manaf, Khairunizah Hazila Khalid, and Nik Iskandar Putra Samsudin. Thank you for their help, guidance, and encouragement. I cannot imagine completing this task without their support.

Finally, my deepest appreciation to my husband, Mr. Mohd Farisan bin Zakaria, and to all family members especially my parents Professor Dr. Hj. Faujan bin Hj. Ahmad @ Amat and Pn. Hjh. Samilah binti Hj. Kutim, and all my siblings Nur Hana, Nur Hadi, Nur Hani, Nur Hafizah, Nur Hidayah, Nur Hakim, Nur Hariz and Nur Haziqah for their prayer, moral support, understanding, and continuous encouragement, in order to make this thesis comes true. This work is also dedicated to my daughter, Nur Qaisarah binti Mohd Farisan.

Last but not least, I would like to express my thanks and gratitude to all patients and volunteers who provided me with their samples which were a huge trigger for the research.

Thank you.

I certify that a Thesis Examination Committee has met on 15 December 2011 to conduct the final examination of Nur Huda binti Faujan on his (or her) thesis entitled "Bacterial and Short-Chain Fatty Acid Profiling of Faecal Samples" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Farinazleen binti Mohd Ghazali, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Son Radu, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Internal Examiner)

Mariana Nor binti Shamsudin, PhD

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Internal Examiner)

Chia-Yin Lee, PhD

Professor
National Taiwan University
Taiwan
(External Examiner)



SEOW HENG FONG, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 2 March 2012

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of Supervisory Committee were as follows:

Fatimah Abu Bakar, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Mohd Yazid Abdul Manap, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Shuhaimi Mustafa, PhD

Associate Professor
Faculty of Biotechnology and Molecular Sciences
Universiti Putra Malaysia
(Member)

Loong Yik Yee, MD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:


DECLARATION

I declare that the thesis is my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.



NUR HUDA BINTI FAUJAN

Date: 15 December 2011



© COPYRIGHT

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vii
ACKNOWLEDGEMENTS	xi
APPROVAL	xiii
DECLARATION	xv
LIST OF TABLES	xix
LIST OF FIGURES	xx
LIST OF ABBREVIATIONS	xxiv
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	6
2.1 Human Gastrointestinal Tract as an Ecosystem	6
2.2 Human Gut Microbiota	8
2.2.1 Classification of Human Gut Microbiota	8
2.2.2 Influence of Age on Human Gut Microbiota	12
2.2.3 The Importance of Human Gut Microbiota to Human Health	15
2.2.4 Role of Gut Microbiota in Gut Disorders and Diseases	18
2.3 Faecal Samples for Molecular Analysis	20
2.3.1 DNA Extraction from Faecal Samples for Molecular Analysis	21
2.4 Real-time Polymerase Chain Reaction	22
2.4.1 Application of Real-time Polymerase Chain Reaction	26
2.4.2 Quantification Methods in Real-time Polymerase Chain Reaction	27
2.4.3 Real-time Polymerase Chain Reaction as Tool for Monitoring Faecal Bacteria in Human Faeces	28
2.5 Characteristics of 16S rRNA Gene	30
2.6 Inflammatory Bowel Disease	31
2.6.1 Definition of Inflammatory Bowel Disease	31
2.6.2 Types of Inflammatory Bowel Disease	32
2.6.3 The Incidence of Inflammatory Bowel Disease in Malaysia	34
2.6.4 Intestinal Microbiota and Inflammatory Bowel Disease	37
2.7 Short-Chain Fatty Acids in Human Intestine	45
3 MATERIALS AND METHODS	48
3.1 Standard Bacterial Strains	48
3.2 Growth Conditions of Bacterial Strains	49
3.3 Subjects and Specimens	49
3.3.1 Characteristics of Inflammatory Bowel Disease Subjects	50
3.3.2 Faecal Samples Collection	51

3.4	Preparation of Template DNA	51
3.4.1	Extraction of Template DNA from Bacterial Cultures	51
3.4.2	Extraction of Template DNA from Faecal Samples	52
3.4.3	Determination of Purity and Concentration of DNA Extraction	54
3.5	Primers	54
3.6	Polymerase Chain Reaction Analysis	56
3.6.1	Amplification of DNA in Conventional Polymerase Chain Reaction for Primers Specificity Test	56
3.6.2	Gel Electrophoresis	56
3.6.3	Purification of PCR Products	57
3.7	Cloning of PCR Products	58
3.7.1	Ligation of PCR Products	58
3.7.2	Transformation of Competent Cells	58
3.7.3	Screening for Plasmid Insert	59
3.8	Isolation and Purification of Plasmid DNA from <i>E. coli</i> Competent Cells	60
3.8.1	Cultivation of Bacterial Strains	60
3.8.2	Extraction and Purification of Plasmid DNA	60
3.8.3	Determination of Plasmid DNA Concentration	61
3.8.4	Sequencing and Analysis of Plasmid DNA	62
3.9	Real-time Polymerase Chain Reaction	62
3.9.1	Preparation of Plasmid DNA Standard for Real-time Polymerase Chain Reaction Quantification	62
3.9.2	Preparation of Standard Calibration Curves from Plasmid DNA for Real-time Polymerase Chain Reaction Quantification	63
3.9.3	Real-time Polymerase Chain Reaction Assay	65
3.9.4	Amplification of Plasmid DNA Standard for Real-time Polymerase Chain Reaction Quantification	66
3.9.5	Control for Each Assay	66
3.9.6	Calculation of 16S rRNA Gene Copy Numbers	67
3.9.7	Copy Number Calculation for Real-time Polymerase Chain Reaction Quantification	68
3.10	Evaluation of DNA Extraction Efficiency from Faecal Samples by Real-time Polymerase Chain Reaction	69
3.11	Determination of Short-Chain Fatty Acids in Faecal Samples	69
3.11.1	Chemicals and Reagents	69
3.11.2	Preparation of Samples for Analysis	70
3.11.3	High Performance Liquid Chromatography for Short-Chain Fatty Acids Analysis	70
3.11.4	Standard Preparation for the Calibration Curve for Reference Standard	71
3.12	Statistical Analysis	71

4	RESULTS AND DISCUSSION	72
4.1	Enrichment of Bacterial Strains for DNA Extraction	72
4.2	Faecal Samples	72
4.3	Evaluation of DNA Extraction and Purification	73
4.3.1	Genomic DNA from Bacterial Cultures	73
4.3.2	Genomic DNA from Faecal Samples	74
4.4	Specificity of Primers Testing using Conventional Polymerase Chain Reaction	79
4.5	Sequence Analysis of Plasmid DNA	86
4.6	Concentration of Plasmid DNA Stock for Real-time Polymerase Chain Reaction Standard Curve Derivation	89
4.7	Real-time Polymerase Chain Reaction	90
4.7.1	Real-time Polymerase Chain Reaction Standard Curves for Quantification of Faecal Bacteria	90
4.7.2	Melting Curves Analysis for Primer Specificity	102
4.8	Efficiency of DNA Extraction from Faecal Samples	107
4.9	Real-time Polymerase Chain Reaction for Bacterial Quantification in Faecal Samples of Inflammatory Bowel Diseases Patients and Healthy Persons	109
4.10	Short-Chain Fatty Acids in Faecal Samples using High Performance Liquid Chromatography	114
4.10.1	Validation of Retention Time for Analytical Methods	114
4.10.2	Short-Chain Fatty Acids in Faecal Samples of Inflammatory Bowel Disease Patients and Healthy Persons	117
4.10.3	Bacterial Population and Short-Chain Fatty Acids in Faecal Samples	122
5	CONCLUSION	125
	REFERENCES	127
	APPENDICES	156
	BIODATA OF STUDENT	171
	LIST OF PUBLICATIONS	172