

## Mycelial Growth and Germanium Uptake by Four Species of *Ganoderma*

CHOW-CHIN TONG and PEI-JOO CHONG  
Center for Research in Enzyme and Microbial Technology (CREAM)  
Faculty of Science and Environmental Studies  
Universiti Pertanian Malaysia  
43400 UPM Serdang, Selangor, Malaysia

**Keywords:** *Ganoderma tsugae*, *G. subamboinense* var. *laevisporium*, *G. tropicum*, *G. lucidum*, germanium

### ABSTRAK

Miselial dari empat spesies *Ganoderma* (*G. tsugae*, *G. subamboinense* var. *laevisporium* ATCC 52419, *G. tropicum* dan *G. lucidum*) telah ditumbuhkan dalam medium cecair yang mempunyai kepekatan germanium (*Ge*) yang berlainan selama 20 hari pada suhu 28°C. Didapati bahawa kadar pertumbuhan dan berat kering miselial keempat-empat spesies semakin menurun dengan penambahan kepekatan *Ge* dalam medium walaupun peratusan pengambilan *Ge* oleh miselial semakin bertambah. Tahap pertahanan miselial terhadap *Ge* di dalam medium pertumbuhan berbeza untuk keempat-empat spesies *Ganoderma*. Untuk setiap spesies, kepekatan optima *Ge* di dalam medium telah ditentukan sebagai 100mg/l untuk mendapatkan keadaan optima bagi pengambilan *Ge* serta pertumbuhan miselial yang optimum.

### ABSTRACT

Four *Ganoderma* species (*G. tsugae*, *G. subamboinense* var. *laevisporium*, ATCC 52419, *G. tropicum* and *G. lucidum*) were incubated in liquid medium containing different concentrations of germanium (*Ge*) for up to 20 days at 28°C. Increasing the *Ge* concentration of the medium resulted in a gradual decrease in the growth of the fungal mycelium. However, the *Ge* content in the mycelium increased with increasing *Ge* concentration. Different species recorded different levels of tolerance towards the *Ge*. In each case, the optimum concentration of the incorporated *Ge* in the medium was established as 100 mg/l for both optimal uptake of *Ge* by the fungal mycelium and optimal mycelial growth.

### INTRODUCTION

Traditionally in the Orient, *Ganoderma* has been considered an "elixir of life". Today, there is a considerable body of contemporary research that shows that certain species of *Ganoderma* are highly effective medicinal agents as a popular remedy to treat hepatopathy, chronic hepatitis, nephritis, hypertension, hyperlipidemia, arthritis, neurasthenia, insomnia, bronchitis, asthma, gastric ulcer, arteriosclerosis, leukopenia, diabetes, anorexia, mushroom poisoning and debility due to prolonged illness (Willard 1990). The medicinal value

of *G. lucidum* is closely linked to the presence of the following compounds: organic-*Ge*, polysaccharides, triterpenoids and adenosine (Tong 1995).

In Malaysia, studies on cultivation techniques of a suitable strain of *G. lucidum* well adapted to the local climatic conditions (Tong and Chen 1990), its growth characteristics and *Ge* uptake by the mycelium (Tong *et al.* 1994a) as well as the fruiting bodies of this fungus (Tong *et al.* 1994b) have been carried out. Because of the paucity of information on the uptake of *Ge* by other species of *Ganoderma* of com-

mercial value, the present study was undertaken to ascertain differences, if any, in their uptake of Ge.

## MATERIALS AND METHODS

### Cultures

Four species, *G. tsugae*, *G. subamboinense* var. *laevisporium* (ATCC 52419), *G. tropicum* and *G. lucidum* were obtained from Taiwan and maintained on potato dextrose agar (PDA) at 28°C and subcultured every 3 weeks.

### Uptake of Ge by Fungal Mycelium Grown in Liquid Medium

Samples (100 ml) of potato dextrose broth containing each of the following concentrations (50, 100, 200, 300, 400 mg/l) of GeO<sub>2</sub> were poured into 1-l flat bottles and sterilized at 121°C, 15 psi for 15 min. Each bottle was then inoculated with three agar mycelium discs (1.3 cm diam) of a ten-day-old culture and incubated in the dark at 28°C for 20 days. The mycelium was later harvested through pre-weighed Whatman No. 1 filter paper and washed with several changes of double distilled deionised water (200 ml). The filter paper, together with the washed mycelium, was dried at 60°C for 48 h. The dry weight of the mycelium was recorded.

### Analysis of Germanium

Fungal mycelium samples were ashed at 700-800°C for 1 – 3 hours. The ash was then dissolved in 5M HCl. Germanium in the solution was determined by hydride generation – inductively coupled plasma atomic emission spectrometry (ICP-AES) using the Labtest Plasmascan 710 instrument. The method of analysis was adapted from the method for arsenic analysis (Lee and Low 1987).

A spectropure (Aldrich Chemical Co. Inc.) Ge atomic absorption standard solution of 990 mg/l was used as the stock solution. The standard solutions used in the

analysis were prepared by sequential dilution from the stock solution. All reagents used were analytical grade. A solution of 2% sodium borohydride in 0.1% sodium hydroxide was prepared daily from NaBH<sub>4</sub> pellets.

The rate of uptake (%) of Ge is calculated as:

$$\frac{\text{Amount of Ge in known dry weight of mycelium}}{\text{Amount of Ge in 100 ml medium}} \times 100$$

## RESULTS AND DISCUSSION

### Effect of Ge on Mycelial Growth in Liquid Medium

With increasing amounts of GeO<sub>2</sub> incorporated into the medium, there was a corresponding decrease in the dry weight of the mycelium (Fig 1). The effect was most obvious for *G. tsugae*, which experienced a continuous sharp drop in the mycelial dry weight at concentrations above 50 mg/l of GeO<sub>2</sub>. Similarly for *G. lucidum*, the growth of the mycelium was




Fig 1. Mycelial growth of *Ganoderma* species in liquid medium containing different concentrations of GeO<sub>2</sub> incubated for 20 days at 28°C. Results represent average of triplicates.

● *G. tsugae* ▲ *G. tropicum* ○ *G. lucidum*  
■ *G. subamboinense* var. *laevisporium*

TABLE 1  
Effect of GeO<sub>2</sub> in liquid medium on the uptake of Ge by mycelium after 20 days of incubation at 28°C

Species	Conc. of GeO <sub>2</sub> in 100 ml liquid medium (mg/l)	Mycelial dry weight (g)	Amount of Ge in mycelium (mg/kg)	Uptake of Ge (%)
<i>G. tsugae</i>	50	0.79	752 + 5	11.8
	100	0.70	1892 + 15	13.3
	200	0.68	2456 + 22	8.3
	300	0.54	3855 + 127	6.9
	400	0.38	4312 + 90	4.0
<i>G. subamboinense</i> var. <i>laevisporium</i>	50	1.43	34 + 4	1.0
	100	1.10	99 + 3	1.1
	200	0.93	175 + 11	0.8
	300	0.87	260 + 9	0.7
	400	0.82	310 + 8	0.6
<i>G. tropicum</i>	50	1.53	235 + 7	7.2
	100	0.94	1080 + 22	10.1
	200	0.52	2072 + 15	5.4
	300	0.46	2715 + 43	4.2
	400	0.37	4002 + 90	4.3
<i>G. lucidum</i>	50	0.12	687 + 18	6.4
	100	0.11	1796 + 14	8.0
	200	0.10	2845 + 31	6.0
	300	0.05	4251 + 112	3.2
	400	0.04	5431 + 87	2.6

Results represent average of triplicates

retarded considerably. This may be due to toxicity at high concentrations. Only extremely sparse growth occurred at concentrations of 250 mg/l of GeO<sub>2</sub> and at concentrations above 300 mg/l of GeO<sub>2</sub> there was no growth of the mycelium at all until the tenth day of incubation (visual observation). The limited growth which commenced after this lag period may probably be due to the induction of some detoxification mechanism.

The growth of the mycelium for *G. subamboinense* var. *laevisporium* was least affected by the amount of GeO<sub>2</sub> in the medium and there was a more gradual decline in the mycelial dry weight than in the other species (Fig 1). Of the four species

tested, *G. tropicum* produced the most abundant mycelial growth at Day 20 and the effect of GeO<sub>2</sub> was more gradual above 200 mg/l of GeO<sub>2</sub> but significantly decreased between 50 and 200 mg/l.

#### *Uptake of Ge by Fungal Mycelium in Liquid Medium*

Table 1 shows that in general, Ge uptake by the mycelium increased with increasing amounts of GeO<sub>2</sub> in the growth medium. Of the four species tested, *G. subamboinense* var. *laevisporium* assimilated least Ge into the mycelium, which may explain why its mycelial growth is least affected by increasing quantities of GeO<sub>2</sub> in the medium. At 400 mg/l of the GeO<sub>2</sub> in the medium, *G.*

*tsugae* and *G. tropicum* took up almost 13 times, while *G. lucidum* took up 17 times more Ge than *G. subamboinense* var. *laevisporium*. However, it was noted that the rate of uptake (%) of Ge by the mycelium increased to a maximum at 100 mg/l of GeO<sub>2</sub> in all species and then decreased with increasing concentration of GeO<sub>2</sub> in the medium. The highest rate of uptake recorded was 13.3, 10.1, 8.0 and 1.1% for *G. tsugae*, *G. tropicum*, *G. lucidum* and *G. subamboinense* var. *laevisporium*, respectively.

Thus, under the conditions studied, it was deduced that for maximum production of fungal mycelium with optimal Ge content, the optimum concentration of GeO<sub>2</sub> to be added to the medium was 100 mg/l.

## REFERENCES

- LEE, C.K. and K.S. LOW. 1987. Determination of arsenic in cocoa beans by hydride generation with inductively coupled plasma atomic emission spectrometry. *Pertanika* **10**: 69-73.
- TONG, C.C. 1995. Nutraceutical properties of the mushroom *Ganoderma*. *Buletin Maklumat Pertanian Malaysia* **15(3)**: 1-13.
- TONG, C.C. and Z.C. CHEN. 1990. Cultivation of *Ganoderma lucidum* in Malaysia. *Mushroom Journal for the Tropics* **10**: 27-30.
- TONG, C.C., S.L. KHOONG and C.K. LEE. 1994a. Growth characteristics and germanium content of *Ganoderma lucidum* growing in different substrates. *Asia-Pacific Journal of Molecular Biology and Biotechnology* **2(4)**: 327-333.
- TONG, C.C., S.L. KHOONG and C.K. LEE. 1994b. Germanium uptake by the fruiting bodies and mycelium of the fungus *Ganoderma lucidum*. *Asean Food Journal* **9(2)**: 69-72.
- WILLARD, T. 1990. *Reishi Mushroom-Herb of Spiritual Potency and Medical Wonder*. Issaquah, Washington: Sylvan Press.

(Received 2 May 1996)

(Accepted 20 August 1996)