

**EXPLORING BIODIVERSITY
& FISHERIES BIOLOGY**
A Fundamental Knowledge for
**Sustainable Fish
Production**



PROFESSOR DR. AZIZ ARSHAD

**EXPLORING BIODIVERSITY
& FISHERIES BIOLOGY**
A Fundamental Knowledge for
**Sustainable Fish
Production**

Prof. Dr. Aziz Arshad

PhD (Plymouth), MS (Wales), BSc (Fisheries, UPM)

24 Januari 2014

Dewan Pertanian
Fakulti Pertanian
Universiti Putra Malaysia



Universiti Putra Malaysia Press

Serdang • 2014

<http://www.penerbit.upm.edu.my>

© **Universiti Putra Malaysia Press**

First Print 2014

All rights reserved. No part of this book may be reproduced in any form without permission in writing from the publisher, except by a reviewer who wishes to quote brief passages in a review written for inclusion in a magazine or newspaper.

UPM Press is a member of the Malaysian Book Publishers Association
(MABOPA)

Membership No.: 9802

Typesetting : Sahariah Abdol Rahim @ Ibrahim

Cover Design : Md Fairus Ahmad

Design, layout and printed by

Penerbit Universiti Putra Malaysia

43400 UPM Serdang

Selangor Darul Ehsan

Tel: 03-8946 8855 / 8854

Fax: 03-8941 6172

<http://www.penerbit.upm.edu.my>

Contents

ABSTRACT	1
INTRODUCTION	3
FISHERIES BIOLOGY RESEARCH	4
BIOLOGY AND POPULATION STUDIES OF SERGESTID SHRIMPS OF <i>ACETES SPP.</i> (DECAPODA:SERGESTIDAE)	5
POPULATION PARAMETERS AND FEEDING BIOLOGY OF SERGESTID SHRIMP, <i>Lucifer hanseni</i>	11
BIOLOGY AND ECOLOGY OF DOG CONCH (<i>Strombus canarium</i> LINNAEUS, 1758) (GASTROPODA: STROMBIDAE)	13
TEMPORAL VARIATIONS IN REPRODUCTION, GROWTH, CONDITION INDEX AND SOME POPULATION PARAMETERS OF GREEN-LIPPED MUSSEL, <i>Perna viridis</i> (LINNAEUS 1758)	20
BIOLOGY AND DISTRIBUTION OF PEN SHELLS (BIVALVIA: PINNIDAE)	24
DISTRIBUTION, LARVAL ABUNDANCE, SPAWNING AND EARLY DEVELOPMENT OF SADDLE OYSTER <i>Placuna ephippium</i> (PHILIPSSON, 1788)	28
BREEDING, SEED PRODUCTION, CULTURE OF THE COMMERCIALY IMPORTANCE SEA URCHINS	30
FISH LARVAL COMPOSITION, DISTRIBUTION AND DIETS IN THE SEAGRASS-MANGROVE ECOSYSTEM	35

REFERENCES	38
BIOGRAPHY	49
ACKNOWLEDGEMENTS	53
LIST OF INAUGURAL LECTURES	55

ABSTRACT

Research on fisheries biology receives less attention due to several underlying factors. However, the opportunity is still great as the scope under fisheries biology is very wide and furthermore, our waters is rich in fish species that are of importance to the fisheries industry. Malaysia dependence on capture fisheries is still strong, as aquaculture production stays at around 30% of the total fish production. Research on wide range of marine organisms that help to enhance the fish production ought to be encouraged. In comparison to finfish, molluscan shells and crustaceans are lagging behind in term of their biological information and have long been receiving less interest from the researchers. This is despite seafood demands for shrimps, crabs and shells are consistently high and on the increase. Taking the matter as an important issue, our research team in UPM has for the past two decades focused and engaged on biological research of this group of fish resources. Knowledge and output on the biological aspects of these invertebrate species is important in many ways; primarily it might help to propel the development of new species for shellfish aquaculture whether as a culture species itself or alternatively acting as a source of live foods for hatchery use. Acting along this objective and tagging up with national interest on increasing fish production, works on the reproductive and feeding biology as well as diversity and ecology are undertaken. The lecture will take a closer look at the research being assumed by us and will discuss the results and findings on several invertebrate species that we studied.



INTRODUCTION

The definition of fish according to Macmillan is an animal that lives in water and swims. It breathes by using its gills and moves by using its tail and fins. However, when we mention fisheries resources, it is now well recognized that this would also include other aquatic resources such as shellfish and seaweeds. When we refer fisheries as a term, we also include the fish populations that are of commercial value encompassing marine, brackishwater and freshwater fisheries. Fisheries biology is a term to describe a scientific discipline that emphasizes the study of fisheries. The scope for fisheries biologists is quite diverse, and ranging from study on fish stocks or populations, its ecological surroundings and on the conservation and sustainable management of the resources. Research on the topic of fisheries biology in Malaysia is still not extensively implemented. We still lack published information on the topic. This could possibly be due to a limited pool of resource persons, lack of interest on the subject amongst fishery researchers on the topic, and perhaps greater priority is given to aquaculture related research.

In terms of application, the knowledge on fisheries biology extends beyond the management measures of the populations. It also encompasses the biology and ecology of the fish species. As a benefit, extensive fisheries biology knowledge of wild but potential aquaculture species would enhance the opportunity for the introduction of perhaps new aquaculture species. In addition, information on the varieties of live foods inhabiting the aquatic ecosystem would allow emergence of nutritious but cost effective live feed in the fish hatchery operation. Established facts on some of the important biological findings of the stocks give specific accounts on their behaviors and these would reciprocally consolidate the management policies been developed by the authority.

Fisheries production in Malaysian is still being dominated by the capture fisheries or catches that come from the sea. This has been going for many decades despite various measures and projects been implemented by the government to boost the aquaculture of the country. Amongst the problem faced by the aquaculture farmers are shortage of good quality and specific pathogen free seeds and spiraling costs of formulated fish feeds. At present aquaculture only contributes about 30% of the total fish production and we have to double the effort to enhance aquaculture sector through research and development as well as high technology input.

With greater concern hailed by fishery managers and established reports made on the overfishing state of capture fisheries, alarming increase in pollution status of the aquatic water body, lost of fish habitat through coastal development and uncertainties due to global climate change, its becoming more of a great apprehension that many of the existing commercial fisheries might be further threatened and the resources would one day be diminished. This would certainly require greater efforts to be channeled towards developing the aquaculture sector as this is the logical alternative for the continuation of the fish supply to the increasing human population. One of the remedial efforts is through the intensive research and development where researches are tailored and focused on the development of new technology and enhanced culture systems with the prime objective of increased production.

FISHERIES BIOLOGY RESEARCH

The overall mission of the UPM Fisheries Biology Research Group is to provide our stakeholders with the knowledge required to ensure the conservation of fish and aquatic resources that are faced with increasing societal demands and increasing loss or degradation of habitat. Our group has undertaken several research projects that

are of significant contribution to the fisheries sector of the country. Most of the emphasis is confined to the study on the biological aspects of the marine organisms mainly marine invertebrates. Both the crustaceans and molluscs are the two important taxa group of fish resources that we prioritized in our research activities. The contents and outputs of the research are given below.

BIOLOGY AND POPULATION STUDIES OF SERGESTID SHRIMPS OF *ACETES SPP.* (DECAPODA: SERGESTIDAE)

Acetes shrimp or locally known as geragau, udang baring, udang bubuk, udang siring is a minor planktonic crustacean group represented by a small number of species but formed one of the economically important organisms in Asia region. (Arshad *et al.*, 2007; Amin *et al.*, 2011). Although *Acetes* shrimp does not belonged to the highly priced shrimp commodity group, it is still important in term of its use as a the fish resources as these shrimps are continuously being exploited by small scale coastal fishermen for the making of shrimp paste, cencalok and several other products. A great number of small income fishermen thrive on this shrimp for their living income although in some fishing areas *Acetes* fishing is seasonal. In view of this social and economical importance, an extensive studies encompassing the taxonomy, morphometric variation, population genetics, reproductive cycle, sex ratio, fecundity, feeding habits, seasonal abundance, growth, mortality, recruitment, yield-per-recruit and status of the stock of *Acetes* spp. were undertaken between 2005 to 2007. Most of the important *Acetes* fishing and landing areas in Peninsular Malaysia are included in the research. Due to limited funds and logistics, we are unable to implement our extensive coverage in Sabah and Sarawak.



Fig . *Acetes indicus*. a, male (x60) and b, female (x60)




Fig a, Procured tooth present in male (x40); b, One clasping spine in male (x40); c, petasma without pars astringens (x40); d, telson of female almost triangular (x40); e, third thoracic sternite is deeply channelled in female (x40).

(Morphological characteristics of *Acetes indicus*, with figures showing diagnostic features that are used in *Acetes* taxonomy)

Earlier works by Pathansali (1966) reported the presence of six *Acetes* species and our review on the species diversity of Peninsular Malaysia and East Malaysia listed five species of *Acetes* viz. *Acetes indicus*, *A. japonicus*, *A. intermedius*, *A. vulgaris* and *A. serrulatus*. *Acetes japonicus* was recorded from samples taken from Klebang Besar (Malacca) and Kuala Gula (Perak). The widely distributed *A. japonicus* was recorded from Klebang Besar (Malacca), Kuala Gula (Perak), Bagan Ajam (Penang), Kuala Sala (Kedah) and Sg Berembang (Perlis). *Acetes intermedius* were identified of samples collected from Seberang Takir (Terengganu) and Bintulu (Sarawak). Both *Acetes vulgaris* and *A. serrulatus* were confined to samples from Pontian and Kukup in Johor (Amin *et al.*, 2011).

We also examined the morphological difference of *A japonicus* specimens at four different fishing sites on the western coast of Peninsular Malaysia and found there are significant difference in term of their total length, standard length and carapace length (Arshad *et al.*, 2013). This perhaps explained the different sizes

of the *Acetes* shrimps being landed at different localities along the western coast of Peninsular Malaysia. Most of the *Acetes* resources is traditionally being exploited by coastal fishermen using push nets and set bag net although trawl net has also been adopted for the exploitation of this shrimp in the east coast of Peninsular Malaysia and Sarawak. We went on to examine the efficiency of using set bag net for trapping *Acetes* shrimps and reported that 90% of the catch of this traditional gear are comprised of *Acetes* shrimps (Oh *et al.*, 2010; Amani *et al.*, 2011). The results label set bag net as non-destructive fishing gear and very selective for *Acetes* fishing.

Our research team also examined the population genetic variation of *Acetes japonicus* of different localities using the Random Amplified Polymorphic DNA (RAPD) marker. A total of 90 samples of *Acetes japonicus*, comprised of 30 (15 males and 15 females) from Kedah, 30 (15 males and 15 females) from Perak and 30 (15 male and 15 females) from Malacca were used. The percentages of polymorphic bands of the three geographic populations investigated varied from 57.77% to 87.77%. Genetic distances between populations and cluster analysis from UPGMA grouped the populations into two major clusters. The Perak and Malacca populations were in one cluster, while the Kedah population was clustered by itself indicating it was genetically different. The genetic distance was the highest for the Kedah and the Malacca populations while the lowest were for the Perak and the Malacca populations which probably has a closed ancestral relationship (Aziz *et al.*, 2010).

As reproduction is a very important biological characteristic of an aquatic organism, we embarked on the reproductive biology study of several *Acetes* species. Our findings on the sex ratio of *A. indicus* and *A. japonicus* in the coastal waters of Malacca showed results that are in favour of females in most months of the

year. The analysis of the annual variation of gonadosomatic index (GSI) showed the continuous breeding character of *A. indicus* and *A. japonicus* throughout the year. Size at first sexual maturity of female *A. indicus* was observed at 23 mm and that was > 17 mm of total length for female *A. japonicus*. The estimated mean fecundity of *A. indicus* was 1666.28 (\pm 46.32) eggs. The mean monthly GSI for females *A. indicus* showed positive and significant ($P < 0.05$) correlation with conductivity ($r = 0.67$), salinity ($r = 0.65$) and TSS ($r = 0.59$). No significant ($P > 0.05$) correlation was found between the mean monthly GSI and the remaining two variables (temperature and dissolved oxygen). The results are included in Amin *et al.* (2009) and Amin *et al.* (2010). The results could be utilized by the relevant government agencies for the specific management of the resources.

Our biological study also includes the feeding habit and stomach's food items of *Acetes* shrimps. According the Simple Resultant Index (% Rs), the stomach contents of *A. indicus* were comprised of plant matters (22.85%), fine sand and mud (16.19%), crustacean appendages (19.03%), debris (15.46%), unidentified fragments (10.56%), zooplankton (6.78%), phytoplankton (6.47%), algae (3.49%), shrimp nauplii (1.25%) and mollusc larvae (0.91%). Similarly, diet compositions of *A. japonicus* were made up of plant matters (31.82%), debris (20.06%), phytoplankton (18.45%), fine sand and mud (11.75%), appendages of decapods (6%), unidentified fragments (5.86%), algae (4.17%) and zooplankton (1.80%). These various compositions of food items proved that the two shrimps are bottom feeder omnivore (refer Amin, 2008; Amani *et al.*, 2011). Details feeding works and diets of *A. serrulatus*, *A. japonicus* and *A. intermedius* are reported in Oh (2011), Amani *et al.* (2011b) and Mashitah (2013) respectively. The finding of these researches also

shed information on the type of food items consumed by *Acetes* and this could be applied for the culture of a particular species.

Our research also looked at the exploitation status of the sergestid shrimps in the fishing area in Klebang Besar Malacca. This fishing ground is being threatened with rapid coastal development and risks of habitat degradation and habitat lost is very high and in some stretches of the coastal belt, *Acetes* fishing is no more practiced. The average monthly catch per unit effort (CPUE) of the estuarine push net (EPN) was estimated at 2.50 (\pm 3.42) kg/fisherman/hr. The total catch comprised of three major categories those were *Acetes* shrimps (90%), followed by fish juveniles (9%) and other shrimps (1%). The annual percent composition of *A. indicus*, *A. japonicas* and *A. intermedius* were found to be 57%, 41% and 2%, respectively. The peak catch was observed in the month of October to December. There was no significant correlation ($P > 0.05$) between monthly catches and environmental parameters (temperature, dissolved oxygen, salinity, conductivity and total suspended solid). The results are documented in Amin (2008) PhD thesis, and later published by Amin *et al.* (2008). The population parameter of *A. intermedius* of Bintulu Sarawak is reported in Mashitah *et al.* (2011) and Mashitah (2013).

Length and weight data are a useful and standard result of fish sampling programs. These data are needed to estimate growth rates, length and age structures, and other components of fish population dynamics. The length-weight relationship of *A. indicus*, *A. japonicus* and *A. intermedius* is reported in Amin *et al.* (2009). The length frequency distribution for *A. indicus* suggested that the population consisted of 'two dominant age group' with mean values of 20.80 (\pm 0.07) mm and 29.85 (\pm 0.09) mm of the total length, respectively. And the population of *A. japonicus* consisted of maximum two age groups, with means of 15.18 (\pm 0.90) mm and

21.56 (± 1.03) mm of total length. The population of *A. intermedius* also consisted of maximum two age groups, with means of 19.18 (± 0.05) mm and 26.92 (± 0.06) mm of the total length. The positive allometric nature of growth for *A. indicus* was observed. However, isometric nature of growth was found in combined sexes of *A. japonicus*. The positive allometric nature of growth was also observed in female and both sexes of *A. intermedius*. There were significant difference between males and females size-frequency distribution of *A. indicus* (Kolmogorov-Smirnov test: $d_{\max} = 0.42$, $P < 0.05$), *A. japonicus* (Kolmogorov-Smirnov test: $d_{\max} = 0.39$, $P < 0.05$) and *A. intermedius* (Kolmogorov-Smirnov test: $d_{\max} = 0.40$, $P < 0.05$). (Amin, 2008; Amin *et al.*, 2008; Arshad *et al.*, 2012).

The growth, mortality, recruitment and relative yield per recruit of *Acetes* spp. were investigated based on monthly length-frequency data, using FiSAT software. Higher natural mortalities of male *A. indicus* and *A. japonicus* versus the fishing mortalities observed from the study indicated the unbalance position in the stock. Exploitation level (E) of female was higher than males in *A. japonicus* population. This study indicated two major recruitment events per year where two cohorts were produced per year for *A. indicus* and *A. japonicus* populations. The recruitment pattern of *A. intermedius* was continuous with one major cohort per year. Results from the analysis of the exploitation rate (E) based on the fishing mortality estimates, and from the relative yield-per-recruit (Y/R), indicate that the *Acetes japonicus* fishery is over exploited although *A. indicus* and *A. intermedius* fishery were slightly below the optimum level of exploitation. This implies that any further unrestrained increase in fishing effort might overshoot the level giving maximum sustainable yield, thus driving the stock down and leading to economic losses. (Amin *et al.*, 2009; Amin *et al.*, 2010; Amin *et al.*, 2012)

POPULATION PARAMETERS AND FEEDING BIOLOGY OF SERGESTID SHRIMP, *Lucifer hanseni*

Another crustacean of fishery importance is another sergestid shrimp called *Lucifer*. It is sometimes known as ghost shrimp. This type of shrimp is particularly abundant in the shallow coastal waters of where it can often create problem with zooplankton sorting due to its mass seasonal occurrence. We embark into *Lucifer* research to gather biological information on its morphological characteristics and feeding habit, to finally determine its suitability as a live feed for the fish hatchery. Growth, mortality, recruitment, status of the stock and feeding habits of *Lucifer hanseni* were examined during November 2009 to October 2010 from five stations along the Tebrau Straits, Tg Pelepas, Johor. We examined the spatial distribution of this shrimp in the Straits and found the population to be highest in density in the seagrass ecosystem as compared to the mangroves.



(*Lucifer hanseni*, one of two species known to occur in Malaysian waters)

The growth, mortality and recruitment of *L. hanseni* were investigated based on monthly length-frequency data, using FiSAT software. The average natural mortality (2.84 yr^{-1}) was higher than the average value of fishing mortality (1.20 yr^{-1}) indicates the imbalance position in the stock. The value for exploitation level (E) was 0.30. This indicates that the fishery of *L. hanseni* in the Tebrau Straits is under exploited. This is based on the assumption that a stock is optimally exploited when fishing mortality (F) equals natural mortality (M), or $E = (F/Z) = 0.5$. The annual recruitment of *L. hanseni* showed continuous recruitment with three major peaks in March, July and October. The peaks were affected by environmental factors and the availability of food source (see Arshad *et al.*, 2010)


Feeding habits and diet composition of *L. hanseni* were also investigated. A total of 600 guts were examined. Diet composition was categorized into five main groups which were debris, phytoplankton, zooplankton, macroalgae and unidentified materials. According to simple resultant index (%SRI), the highest percentage of food items was debris (42.76%) followed by phytoplankton (34.96%), zooplankton (12.56%), unidentified materials (5.80%) and macroalgae (3.93%). The various composition of food items proved that *L. hanseni* is opportunistic omnivorous. *Lucifer hanseni* can switch their food preference depending on the abundance of certain types of food during the time (Norhafizah, 2013).

Based on the stomach content, it can be derived that this species also has the potential to be cultured in controlled environment and used them to feed the fish during their early development. The next option is to try and culture them in tank and then scale it up, we haven't had any trials to this extent yet.

BIOLOGY AND ECOLOGY OF DOG CONCH (*Strombus canarium* Linnaeus, 1758) (GASTROPODA: STROMBIDAE)

This research is in particular the first most extensive works ever carried out on this species of gastropod anywhere in the region. The dog conch, *Strombus canarium* is an important species within the seagrass bed ecosystem of the Merambong Shoal and its contribution to the productivity of the seagrass ecosystem is important (Cob *et al.*, 2009a). Being the largest and most dominant herbivorous gastropod, they certainly might contribute significant role in the wellbeing and maintenance of the seagrass ecosystem. The species is also important economically, as people have been utilizing them for ages. It provides important artisanal, subsistence fisheries for the locals especially those who live along the shoreline.

The conch-collecting season at Merambong shoal in Tg Pelepas Johor was, however seasonal and restricted to only a few months, which was well anticipated by the locals (Cob *et al.*, 2009d). Previously the conch was harvested for family consumption only, but now most collectors sold them to local retailers. Considering the importance, abundance and economic value of the species, it was surprising that the population was not being thoroughly studied before. The main objectives of this study were to investigate, analyzed and document the life history, biology and ecology of the species.



(Siput Gonggong, *Strombus canarium* (Cob, 2008))

The life history characteristics of the species, from eggs to adult, have successfully described (Cob *et al.*, 2009e). *Strombus canarium* produced a long gelatinous tube of egg strand, in the form of an egg mass. The egg mass was spawned gradually, from one end to the other by dorso-ventral looping movements of the propodium, and a single female could lay up to 2 egg masses each time. The egg masses were mostly found securely attached to the seagrass. The eggs took between 4.5 to 5.5 days of incubation time before hatch, producing highly phototactic veliger larvae that swim actively towards the surface.

Our team then went on to examine the larval development. The planktotrophic stage ended for about 3 weeks, before metamorphosed and settled as early as day-18, with short period of metamorphic competence (5 - 6 days). Sequence of behavioral pattern related to metamorphosis was also observed and described. This behavior resulted in completely camouflaged juveniles that

physically blend into the surrounding sediments. At this stage the juvenile shell was very fragile, and they still unable to bury and hide into the sediment therefore camouflaging is important as their defense mechanism. This is why small juveniles are extremely difficult to sample in the field (Cob *et al.*, 2009e).

At the time of settlement, specific metamorphic inducers must be in contact. Specific metamorphic cue is needed as no spontaneous metamorphosis was observed. In this study, metamorphic cue associated with sediment from nursery site was found to be the strongest. It was also important to note that sediment taken from other than the nursery ground was unable to induce metamorphosis. That is why, in their natural habitat the juveniles are found congregating at only specific areas within the shoal. The larvae might have specifically settled to this location. The nursery habitats were characterized by low to medium density of *Halophila* bed that mixed with low density of *Thalassia* and/or *Halodule* grasses, and most importantly has high sediment organic content, high sorting coefficient and low in mean particle grain size. Throughout the study period the conch nursery sites were consistently found located within the microhabitat described above. This type of habitat (high sediment organic content) actually provides important source of food for the newly settled juveniles, as was observed during the laboratory culture experiment (Cob *et al.*, 2010).

The growth rate of the species studied was relatively high compared with reported value for other *Strombus* species. In this study growth parameters were determined using different estimation method, i.e. Electronic Length Frequency Analysis (ELEFAN I) and Length-based Fishery Stock Assessment (LFSA) protocol in FiSAT software package. Both results were comparable to each other, and more importantly were also comparable to the actual growth rates derived from tagging experiment and laboratory culture of

the juvenile. There was much debate on the application of these methods for organisms other than fish, but this study proved that it was also useful and applicable to the gastropods (Cob *et al.*, 2008a; Cob *et al.*, 2008c; Cob *et al.*, 2009b).

Although the juveniles were normally limited to areas within the nursery habitat, adult populations on the other hand were more widely distributed within the shoal. They however normally present in patches of local colonies and were more abundant in sheltered areas compared with those facing open seas. Multivariate analysis found that adult population was highly correlated with area dominated by *Halophila* sp., mixed with low density of *Thalassia* sp. and *Enhalus acoroides*. The distribution / conch density was also found highly correlated with sediment organic content and negatively correlated with macroalgae (*Ulva*) density. This explains the general distribution pattern of the adults described in above paragraph. The result also suggests that food is one of the important factors, apart from reproduction, that determined their distribution along the shoal. Most of conch feeding session was spent on probing and selecting food materials within the top most layer of sediment and detritus surface, with very little time spend grazing on fresh macrophyte stands (Cob, 2008; Cob *et al.*, 2012).

In many instances, *Strombus* are referred as detritivore that feeds directly on seagrass detritus. In this study *S. canarium* was however not consuming the detritus itself but rather grazed on the epibiota that live on its surface. Although stomach content analysis showed large amount of detrital materials, its low calorific value, high fiber content and low digestibility make its role as major source of nutrient uncertain. The detrital material might have accidentally been eaten and low digestibility accumulates them in the conch stomach (Cob, 2008).

The *S. canarium* population studied was highly seasonal in abundance. This explain why the fishery activity only viable during certain period of the year. Monthly length frequency distribution indicates high variation in conch densities, which was mostly due to variations in adult populations. There was significantly higher number of adults during wet season compared to dry season. The adults most probably have migrated to deeper areas during dry season (April to September) and migrate back into the shoal area to copulate and spawn during breeding season (October to March). Unfortunately, migratory behavior observation cannot be conducted for the population (Cob *et al.*, 2009c).

Results from various experiment and analyses during this study have narrowed down to the conclusion that reproduction might be the main reason behind this fluctuation / migration of adults. There was high correlation between conch density and GSI. Although GSI alone cannot satisfactorily determined breeding season, combined with other biological, ecological and behavioral observations (condition index, condition factor, egg mass density, frequency of copulating pairs and population density), it can adequately be concluded that the species was indeed highly seasonal in reproduction. The reproductive season starts around early November to late March. The species is highly synchronized in their reproduction, which is important for their survivals. Nevertheless reproduction also occurred all year round, particularly around July to August where slight increased in GSI, density, coupling and spawning was observed. A few of the cohorts identified in population dynamic study were found originated (spawned) from this period. However, larval growth and development during this period (dry season) was inferior compared to wet season. This indicates that dry season is not a good season for reproduction (Cob *et al.*, 2008b).

Sexual dimorphism is obvious within the *Strombus* as they are dioecious. In this study, male and female *S. canarium* not only differed morphologically, but also showed differences in growth rates, sexual maturity and longevity. Apart from that, *S. canarium* showed sexual polymorphism where another morphs have been identified i.e. the imposex females. The later have larger, heavier and higher degrees of flaring compared with both normal males and females, and were easily recognized via the present of male genitalia on female reproductive organ. Details on imposex findings are reported in Cob *et al.* (2011).

Based on relative yield per recruit analysis, the population studied was currently still underexploited, which can be attributed to the low technology involved. There was no specific tools used where the conch were only manually collected. The collecting activity is also restricted to only a few days in a month (those with extreme low tides). Apart from that, the important factor that kept the landings low is the migratory behavior of the species, as was described earlier, which make the harvesting only viable within a few months in a year, i.e. during their breeding season.

On extensive data and information gathered in the research, our team has been succesfull in investigating the various aspects of biology and ecology of *Strombus canarium* and got them published. In general, it shares great similarity in terms of anatomy and biology with other *Strombus* species. Nevertheless, various aspects of biology and ecology unique to this tropical species have been established. There is certainly good prospect of introducing *S. canarium* for aquaculture. This includes sea-farming and restocking juveniles to the natural stock for fishery purpose, which has been practiced in the Caribbean for *S. gigas*. Among the important characteristics of *S. canarium* which make it very feasible for introduction into aquaculture are their;

Economic features: High quality and delectable meat, which is readily acceptable by the locals. The species was well distributed within the region thus good for market expansion. It has shell of high quality that can be sold as souvenir, collectible and other purpose (refer Cob, 2008)

Life History development: They live in colony and normally present in high numbers, therefore easily utilized. Spawning occur year round and egg mass available all season, easily collected, thus there will be no issues in terms of seed supply. Able to breed in captivity, with high fecundity and high percentage of egg-mass hatchability. They are herbivorous and detritivorous, which is a favorable trophic level for culture.

Nevertheless, more studies needed in order to assess the practicality, cost effective and viability of the production before engaging in any aquaculture project on the species. Both intensive and extensive culture methods for commercial hatchery production of seeds and grow out should thoroughly be investigated and evaluated. Several methods have been proposed for conch grow out: establishing protected areas for ocean ranching by reseeding of seagrass flat, stocking shallow channel-like raceways, or ponds along the shore with water circulation by the tidal fluctuation, and penning or embaying stocked juveniles.

TEMPORAL VARIATIONS IN REPRODUCTION, GROWTH, CONDITION INDEX AND SOME POPULATION PARAMETERS OF GREEN-LIPPED MUSSEL, *Perna viridis* (LINNAEUS 1758)

We embarked on this research primarily not only because green mussel is one of the commercially importance shellfish in the country, but also because there have been very few additional updates especially on the biology of this bivalves since some few earlier works in the 60s and 70s. Mussel farming to the fisheries sector can provide an alternative to the fluctuating supply of wild mussels. Cultivation on suspended ropes facilitates not only maintains continuity of production but also gives faster growth, better product, lower mortality, fuller use of the water column and the use of previously unproductive areas. In Malaysia, suspended rope culture has been conducted on experimental scale in several areas in Penang, Lumut, Perak and in the Straits of Johor. Spat falls in the Straits of Johor have been found to occur throughout the year with peaks from February to May and October to November in the east, and from June to August and November to December in the west (Choo, 1983).

Accurate forecasting of spat fall is critically vital for sustaining a good mussel farming. Therefore, a thorough study on the reproductive biology of *P. viridis* and defining the accurate times of collecting spat is essential to sustain the growing mussel culture industry in Malaysia. If proven to be successful, this will eliminate the need for hatchery technology which is thought to be uneconomical viable in the near future. Most of the previous studies tackling this problem have concluded their findings based on few culture experiments, plankton and spat collection only. In this research study the findings will be based on the more in detail research. The specific aim of this research was to investigate

the temporal reproductive cycle, growth rate, sex ratio, spat fall densities, condition index and population parameters of green-lipped mussel, *Perna viridis* under Malaysian condition (Barwani *et al.*, 2007).

Condition Index (which relates tissue dry weight to shell volume), of the green mussel *P. viridis* was concurrently studied with the histological observations of the gonads. Monthly samples of *P. viridis* were collected from the same culture site. Variations in average monthly Condition Index ranged from 21.06 to 26.72 g/cm³. Rapid declining of the Condition Indices of the mussels was recorded in December, 2003 and January, 2004. This rapid declining indicates the spawning period which agrees with the results from the histological study (Barwani *et al.*, 2007).

Studies on the gametogenesis, size at sexual differentiation, and size at first maturity of *P. viridis* from Sebatu, Malacca were performed using histological sections. More than three hundred specimens of *P. viridis* between 14.96 and 113.52 mm in total shell length were examined. No gonadal development was observed in individuals smaller than 17 mm. Sexual differentiations began at 18 mm with incipient acini formation. Specimens larger than 20.0 mm had well-developed gonads at different stages of maturation. Size at first maturity was smaller than previously reported at other localities. Active gametogenesis was recorded throughout the study period, with all stages of development usually occurring simultaneously within the same sample population (Barwani *et al.*, 2007).

In the determination of gonadal index (GI), over 300 individuals were histologically prepared and analyzed. A clear fall in gonadal index was observed in the months of January and February, at this time most of the female were clearly spent. A second fall in GI with less intense was observed again during the months of September and October. Gonadal index results show that spawning event in *P.*

viridis is year around except during the months of May until August. The release of gamete between sexes was fairly synchronized, no differences in GI were observed between the male and female. 1000 specimen of *P. viridis* were used for sex ratio. The number of visually identified male, female and undetermined by looking at the mantle was 524, 412 and 64, respectively. The male: female ratio was 0.79:1 by visual identification. 316 individuals were histologically identified for sex ratio. The number of identified male, female and undetermined individual was 161, 151 and 4. The histological male: female ratio was 0.94:1. Two hermaphrodites specimen were recorded during the months of June and September, 2004. Both specimens had the occurrence of the male and female gametes on only one side of the mantle. No trace of hermaphroditism was observed neither in the second side of the mantle nor the mesosoma (Barwani *et al.*, 2007; Barwani *et al.*, 2013).

Spat fall study of *P. viridis* was carried out from February, 2004 until December 2004 on collector ropes suspended from the same culture site. Spat fall of the mussels was active during the entire study period with two peaks, one main peak during the months of March to June and a second peak during the months of August to December. Average, maximum and minimum settlement densities were 11,389, 29,788 and 1649 spat per meter respectively. Maximum settlement densities were counted during the month of April, while the minimum settlement densities were counted during the month of August (Barwani *et al.*, 2007).

Population parameters like asymptotic length (L_{∞}), growth co-efficient (K), mortalities, exploitation level (E) and recruitment pattern of green mussel *P. viridis* from the same site have been studied. The study was carried out using the length frequency based analysis of FiSAT software to evaluate the growth parameters, mortality rates and exploitation level. Asymptotic length (L_{∞})

was 102.38 mm and growth co-efficient (K) was estimated at 1.50 yr⁻¹. Total mortality was 1.44 yr⁻¹ for *P. viridis*. Natural mortality (M) and fishing mortality (F) were 1.67 yr⁻¹ and 0.81 yr⁻¹, respectively. Exploitation level (E) of *P. viridis* was 0.32 while the maximum allowable limit of exploitation (E_{max}) value was 0.43. The exploitation level (< 0.50) is below the optimum level which indicates the under fishing condition of *P. viridis* in the coast of Sebatu/ Malacca, Malaysia (Barwani *et al.*, 2007).

Average values recorded for temperature, dissolved oxygen, pH, and salinity were 29.44 °C, 6.06 mg/l, 7.92, 30.12 ppt respectively. Highest values recorded for temperature, dissolved oxygen, pH, and salinity were 32.17 °C, 7.61 mg/l, 8.44, 33.26 ppt respectively. Lowest values recorded for temperature, dissolved oxygen, pH, and salinity were 27.82 °C, 4.94 mg/l, 7.32, 27.93 ppt, respectively. Chlorophyll-a, and total suspended solids (TSS) ranged from 1.50 to 9.40 mg/l, and 0.0253 to 0.0733 g/l, respectively (Barwani *et al.*, 2007) .

We also work on the causal point for spat mortality and found pea crab, a natural parasite to the molluscs. Incidences of pea crab *Arcotheres latissimus* infestation in *P. viridis* at Sebatu was monitored from December 2003 until December 2004. Maximum infestation frequency was observed during the month of June. Size ranged of infested *P. viridis* from 19.26 to 92.18 mm. Wet weight of infested *P. viridis* ranged from 0.17 to 10.41 g. Wet weight of pea crabs ranged from 0.009 to 0.55 g. The information shed some light on the enemy of cultured mussel in our waters (Barwani *et al.*, 2007).

BIOLOGY AND DISTRIBUTION OF PEN SHELLS (BIVALVIA: PINNIDAE)

Research on the taxonomy, biology and ecology of pen shells were conducted in Merambong shoal off the South-western state of Johor, Peninsular Malaysia. Several pen shells specimens were also acquired from other sites in Peninsular Malaysia to allow wider scope of sample collection. The study was conducted from August 2005 to April 2007. The site was chosen due to ample abundance of pen shells that live associatedly with the seagrass. A total of seven species have been recorded from the study areas comprising five species of *Pinna* and two species of *Atrina*. Ten internal and external morphological characteristics have been used for the taxonomic identification of pen shells (Idris *et al.*, 2008b; Idris *et al.*, 2009). *Pinna* species showed that of the ten characteristics analyzed, four characteristics were highly significant ($P < 0.01$) (WL, DPML, PAMPDNL and WS). Similar result of four morphometric characteristics showed a highly significant ($P < 0.01$) between the characters of three closely related species, *Pinna bicolor*, *P. deltodes* and *P. atropurpurea*. (Idris *et al.*, 2008a; ; Idris *et al.*, 2008c; Idris, 2009; Idris *et al.*, 2012).



(*Pinna pectinata*, of the *Pinna* species found in Merambong Shoal, Johor)

Adductor muscle tissue used for the isolation of DNA and RAPD successfully detected polymorphisms in the pen shells populations. The result showed 19 primers have produced various banding patterns and thus provided sufficient information for reliable discrimination of the analyzed samples. The results indicated that the primers used generated a total of 160 fragments with 70% to 100% of polymorphic fragments. The genetic distance among these seven species of pen shells was in the range of 0.0197 to 0.3190. The dendrograms constructed from RAPD markers data were able to reveal the relationships between the pen shells populations (Idris, 2009)..

Enhalus acoroides, *Halophila ovalis*, *Halophila minor*, *Cymodocea serrulata* and *Thalassia hemprichii* were among the seagrasses associated with pen shells habitat in Merambong and Tanjung Adang shoals. However, pen shells can also be found in stony sand area of Merambong Island and live associated with

zoanthid, *Zoanthus pulchellus*. The sediment types from four study areas were classified as sandy loam. Monthly *in situ* physico-chemical seawater parameters recorded showed no significant difference ($P > 0.05$) during the study period. Pen shells were recorded and classified into three classes of distribution i.e. clumped, random and rare.

Higher density was shown by *P. bicolor* in Merambong shoal (1) with the value of 0.83 ind/m² while *P. incurva* showed lower density with 0.03 ind/m² in Tanjung Adang shoal. *Pinna bicolor*, *P. deltodes* and *P. atropurpurea* were consistently found in these four study areas while, *P. deltodes* Menke and *P. incurva* Gmelin were both a new distribution record for Sungai Pulai seagrass beds. Merambong shoal population recorded higher diversity and richness as compared to the Tanjung Adang shoal and Merambong Island, but the value of evenness was similar between Merambong shoal, Tanjung Adang shoal and Merambong Island. Five major phyla comprising 37 species of fouling organisms were recorded. Members from phylum Crustacea and Mollusca were higher in percentage of distribution when compared to the Annelida, Echinodermata and Chordata. Phylum Mollusca showed the highest diversity while phylum Annelida was recorded the highest species richness. A symbiotic adult alpheid shrimp, *Synalpheus carinatus* was recorded inhabiting the mantle cavity of the pen shells (Idris, 2009).

Pinna bicolor reaches sexual maturity at shell length of 170 mm. *Pinna bicolor* is dioecious and no hermaphrodite individual was found during the study period. Five stages of gonad development were observed and clearly been identified. *Pinna bicolor* showed a clear spent phase in the month of October 2006, December 2006 and March 2007 while the developing and spawning phases were

observed throughout the whole study period. Monthly *in situ* physico-chemical parameters and rainfall recorded during the study did not significantly correlated with the reproductive activity of *P. bicolor* in Merambong Shoal. Pearson Correlation analysis also did not show any significant correlation between gonad index (GI) and physico-chemical parameters of seawater in Merambong shoal (Idris, 2009)..

Growth rate in natural habitat has been found to be indeterminate and rapid when compared to *P. bicolor* in culture tank. For the length-weight relationships, the growth coefficient “*b*” was found to be significantly higher than the isometric value (3.111) at 5% level and this is an indication of isometric growth in *P. bicolor* from Merambong shoal. The adductor muscle of *A. vexillum* was found the biggest in size when compared to other species (Idris 2009).

The study on penshells, *Pinna* spp. has generated new information on the biology and ecology characteristics of the bivalves. There has been no earlier works on the species and our research group has managed to carry out comprehensive studies that allow us to further research on the culture of penshells and its seed production.

**DISTRIBUTION, LARVAL ABUNDANCE, SPAWNING
AND EARLY DEVELOPMENT OF SADDLE OYSTER
Placuna ehippium (PHILIPSSON, 1788)**



Placuna ehippium from family Placunidae has never been previously studied in Malaysia although it has long been known to the local coastal communities in Pulai River Estuary, Johor as ‘Senteng’ and has been collected for food for generations by the fishermen there during low tide period. *P. ehippium* has a kind sweet taste and possesses succulent meat and adductor muscle. It is a delicacy among local fishermen communities. This bivalve species was selected as one of the studied species because of its high potential as new aquaculture product with its great taste and durability in high impact area. The research that we embarked in also allowed us to survey other molluscs species of economic importance that are quite abundant living in the Marambong Shoal seagrass beds. Merambong Shoal has been under constant threat of fauna extinction due to habitat lost since intensive development that taken shape for the construction of Port Tanjung Pelepas.

The distribution of this species and other molluscan in Merambong Shoal seagrass bed has been studied, along with the diversity and coverage of marine macrophytes, the sediment texture profile of Merambong Shoal, and their correlations with bivalve distribution. The abundance in density and distribution of 15 epifaunal bivalve species from 10 families have been estimated, and the coverage of 7 seagrass species and 1 genus of macroalgae on the shoal during low tide were calculated. *Placuna ehippium* showed great abundance in certain section of the seagrass beds in particular the sheltered part of the shoal. In general, epifaunal bivalves abundance and macrophyte coverage were more heterogeneous in the southern shoal while more homogenous at the northern shoal.

Temporal larval abundance of Placunidae and other bivalves were explored aiming to discover the natural spat fall season and explore the theory of seagrass meadow being a nursery for non-habitant bivalve larvae. The correlations between temporal bivalve larval abundance, environmental parameters and rainfall were investigated. Placunidae larvae were found from September to January of the study period and none during other months. The highest abundance of Placunidae was recorded in November 2007 with density 226 larvae m⁻³. Major peak on total bivalve larval abundance was noticed between November and January 2008. Through with research, our team has been able to document the larval diversity present on the Merambong seagrass beds. Larvae of six bivalve families were identified by umbonal shell morphology with Mytilidae larvae highest in abundance during November (7345 larvae m⁻³). No significant correlation was found between bivalve larvae abundance and seawater parameters like temperature, salinity, DO, pH and TSS. There are significant correlations between total larvae abundance and total rainfall within a week and two weeks prior to sample collection ($P < 0.01$) (Nur Leena, 2013),

Early development of *P. ehippium* was recorded and described. *Placuna ehippium* broodstocks were collected during low tide in Merambong Shoal seagrass bed. Thermal cycling procedure had successfully induced the spawning of this species. The average size of the spherical eggs before fertilisation was $57.65 \pm 5.97 \mu\text{m}$ in diameter and $90.91 \pm 2.04\%$ eggs were fertilized successfully. Embryos developed into straight-hinged larvae 22 hours after fertilisation. Umbo started to take shape from Day-2. Larval shells are inequivalve with right valve almost flat and left valve inflated. Larvae developed into pediveliger on Day-9 and started to form plantigrade on Day-11. Rapid decreases were observed in survival rate during settlement (Nur Leena, 2013).

The results have shed some lights into the adoption of this species of bivalves as one of the candidates for the bivalves mariculture. Controlled reproduction can be initiated but not to deny more information on the larval rearing of this *Placuna ehippium* needs to be sorted out in order to allow us to practice its culture or restocking to the wild habitats.


BREEDING, SEED PRODUCTION, CULTURE OF THE COMMERCIALLY IMPORTANCE SEA URCHINS

This particular research is amongst the most recent works that we undertake in UPM. Sea urchin is a marine invertebrate that belongs to phylum Echinodermata. Sea urchin has its own commercial importance as the gonad is used to produce sea urchin roe sushi and is considered as a prized delicacy mainly in Japan (Kaneniwa and Takagi, 1986). Sea urchin gonads are rich in valuable bioactive compounds, such as polyunsaturated fatty acids (PUFAs) and β -carotene. PUFAs, especially eicosapentaenoic acid (EPA,

C20:5) (n-3)) and docosahexaenoic acid (DHA C22:6 (n-3)), have significant preventive effects on arrhythmia, cardiovascular diseases and cancer (Pulz and Gross, 2004). On the other hand, the high levels of AA and EPA recently detected in *Diadema setosum* and *Salmacis sphaeroides* supported the development of aquaculture of this urchin (Chen *et al.*, 2009), since PUFAs are important for human nutrition.


Eleven species of sea urchins have been documented in Malaysian waters including *Diadema setosum* and *Salmacis sphaeroides*, (Rahman and Yusoff, 2010). However, very few systematic works have been done on the abundance, distribution and population growth patterns of *D. setosum* and *S. sphaeroides* in Malaysia (Rahman *et al.*, 2012a, 2013) but no published information on their breeding, nursing, seed production and culture techniques are available. Due to the higher nutritional values of sea urchin gonad, it is very important to develop appropriate techniques for breeding and nursing. In view of this, two projects have been undertaken (i) to develop a viable methodology for breeding, seed production and culture of *D. setosum* and *S. sphaeroides* in captivity (ii) to determine the biochemical composition of gonads in a view to develop nutraceutical and pharmaceutical products.

Sexually matured adults of the sea urchins, *D. setosum* (Figure below) and *S. sphaeroides* (Figure below), were collected from Pulau Pangkor, Perak and Merambong shoal, Johor.




(Matured adults of tropical sea urchins: A) *D. setosum* (Rahman et al. 2012a), B) *S. sphaeroides* (Rahman et al. 2012b).

Gametes from both female and male urchins were obtained by injecting 0.5 M KCl into the coelomic cavity. Eggs were collected and fertilization was done at limited sperm concentration and the resulting embryos and larvae were reared. When the larvae attained feeding stage (four-armed pluteus), they were cultured in glass bottles on a rotating roller with a larval density of 1-2 individual/ml. Larvae were supplemented with a cultured phytoplankton, *Chaetoceros calcitrans* at concentrations of 4,000-8,000 cells per ml of medium daily until attaining metamorphic competence within 1 month post-fertilization.




(Larval developmental stages of *S. sphaeroides*: A) 4-arm pluteus, B) 6-arm pluteus, C) 8-arm pluteus, D) Competent larva with complete rudiment growth (Rahman et al. 2012b).




(Larval developmental stages of *D. setosum*: A) 4-arm pluteus, B) POA (Postoral arm)-elongated stage-1, C) POA-elongated stage-2, D) Competent larva with complete rudiment growth and development

Induction of metamorphosis was performed on coralline red algal extracts + *Chaetoceros* diatom (50:50) in petri dishes (9.0 x 3.0 cm) containing FSW. Majority of the competent larvae were metamorphosed to young juvenile within 1 day post-settlement and then cultured on coralline algal stones in aerated aquaria for three months by which time they attained appropriate juvenile sizes (Fig. 3a,b) for stocking in grow out culture aquaria.




(Juveniles of *S. sphaeroides*: A) 1-day-old juvenile, B) 3-month-old juvenile (Rahman *et al.* 2012b).



(Juveniles of *D. setosum*: A) 1-day-old juvenile,
B) 3-month-old juvenile)

In order to develop the appropriate aquaculture techniques, the 3-month old juvenile urchins are reared in aquaria (46 x 30 x 30 cm) at different stocking densities and algal feeding regimes. After two years of rearing in captive condition, all the urchins attained adult sizes with adequate matured gonads for harvesting.



(Sexually matured adult sea urchins after the culture period of two years in captive aquaria-rearing condition: A) *S. sphaeroides*,
B) *D. setosum*)

This study demonstrates the first successful culturing of *S. sphaeroides* and *D. setosum* through the detailed larval development, metamorphosis and juvenile rearing in captivity (Rahman *et al.*, 2012b). Therefore, the results obtained from the designated project will immensely be helpful towards the development of breeding, seed production and culture techniques of commercially important sea urchins and other marine invertebrates, which are yet to be fully determined and explored in the Malaysian coral reef communities. In addition, development of appropriate rearing and culture techniques would immensely be helpful to produce adequate quantities of nutraceutical and pharmaceutical products from these high-valued Echinoderms.

FISH LARVAL COMPOSITION, DISTRIBUTION AND DIETS IN THE SEAGRASS-MANGROVE ECOSYSTEM

In addition to the research done on the invertebrates, our group also embarked on the diversity, ecology and biology of fish larvae. The interest to assume the works came from our integrated research in Merambong Shoal, Johor. Fish larval composition, spatio-temporal distribution, density, family richness, Shannon-Wiener index and feeding habits were determined by analyzing samples collected from the seagrass-mangrove ecosystem of Johor Strait, Peninsular Malaysia between October 2007 and September 2008. Five stations were established to represent the different ecological complexities of the Merambong seagrass ecosystem. In total, 24 families of fish larvae belonging to six Orders were identified from seagrass-mangrove ecosystem of Johor Straits (Ara, 2011; Ara *et al.*, 2013b; Ara *et al.*, 2013c).

Five Common Families



Blenniidae, 50x




Gobiidae, 50x



Sillaginidae, 50x



Clupeidae, 40x



Terapontidae, 40x

(Some important fish larvae collected from the Merambong Shoal seagrass beds)

In total, 24 families were identified from the seagrass ecosystem of the Johor Straits. In comparison to all ecological complex, seagrass ecosystem recorded highest fish larval abundance. Overall, five families viz. Clupeidae, Blenniidae, Terapontidae, Gobiidae and Sillaginidae were the top five dominant taxa in the study area. Shannon-Wiener index varied significantly between monsoon and inter-monsoon seasons and peaking in the months October-January and May-August. The highest density of larval fishes was recorded at seagrass station and the spatial variations in larval density were significant ($p < 0.05$) between seagrass and other four sampling sides. Among the 24 families, 18 families were found to have correlation either positively or negatively with the surrounding water parameters. The highest and significant regression coefficient was observed in Sillaginidae which indicated its abundance was

influenced by the surrounding water parameters (Ara, 2011; Ara *et al.*, 2011b; Ara *et al.*, 2011c).

For this research, 267 Blenniidae, 401 Clupeidae, 126 Gobiidae and 117 Terapontidae stomachs were biologically examined. Analyses of prey in the stomachs identified 24 important items belonging to six major groups viz. phytoplankton, zooplankton, algae, plant like matter, debris and unidentified materials. According to the Simple Resultant Index analysis, the predominant food item in the stomach of all four families (Blenniidae, Clupeidae, Gobiidae and Terapontidae) was phytoplankton (> 60%). This was followed by zooplankton in Blenniidae (18.24%) and Clupeidae (8.60%). On the other hand, the second diet composition was unidentified plant-like matter in Gobiidae (14.73%) and Terapontidae (8.02%).

Among phytoplankton, *Dacytyloccopsis fascicularis* (26.31%) was large in quantity in stomach sacs of Blenniidae and this was followed by *Nitzschia baccata* (23.38%). Conversely, *Nitzschia* sp. showed the highest quantity (26.33%) in the stomach sacs of Clupeidae larvae. In Gobiidae larvae, Chromophyta (28.30%) was observed as the highest quantity. Similarly, Chromophyta (29.12%) was the highest quantity in the stomach sacs of Terapontidae. Overall, two most dominant phytoplankton (*Dacytyloccopsis fascicularis* and *Nitzschia* sp.) was observed among the four larval families. It is revealed that diverse food items were found in the stomachs of Blenniidae, Clupeidae, Gobiidae and Terapontidae larvae and phytoplankton formed more than 60% in the diet composition (Ara *et al.*, 2009; Ara *et al.*, 2010; Ara *et al.*, 2011a; Ara *et al.*, 2013a; Arshad *et al.*, 2013).

The fish larval works give us further information on the importance of seagrass ecosystem for the breeding and nursing of finfish species. This would perhaps highlight the importance of this ecosystem to fisheries as a contrast to the expansion of Port Tg

Pelepas Johor that are seen threatening the existence of seagrass ecosystem in the area.

SUMMARY

The fisheries biology research undertaken by the researchers from the Department of Aquaculture, Universiti Putra Malaysia has managed to establish extensive information on the biology, diversity and ecology of several important crustaceans mainly sergestids *Acetes* spp. and *Lucifer* sp., and molluscs in particular *Strombus* shell, green mussel, pen shell and saddle oyster. Findings on the sergestids would be of importance on the possibility of developing both sergestid species as a source of live food for the fish hatchery. In addition, the knowledge on the specific bivalves and gastropod species could be further utilized in promoting them as an alternative to the existing culture species. Vast amounts of publications have been generated during the implementation of the projects. Some of the new findings from the research are not yet fully known to the relevant authorities and it is our major expectation from the beginning that all our research findings would be of great benefit to the development of fisheries sector of the country.

REFERENCES

- Amani, Alsayed Abdul Aziz, 2011. Feeding Preferences and Population Parameters of Sergestid Shrimps, *Acetes japonicus* in the Coastal Waters of Kedah, Malaysia. Master Thesis. Universiti Putra Malaysia. 83 pp.
- Amani A A, S M Nurul Amin, A Arshad and M. Aminur Rahman, 2011a. Population Dynamics of Sergestid Shrimps *Acetes japonicus* in the Estuary of Tanjung Dawai, Kedah Peninsular Malaysia. *Journal of Fisheries and Aquatic Science*, 6(7), 751 - 760 (ISSN 1816-4927).

- Amani A A, S M N Amin and A Arshad, 2011b. Stomach Contents of Sergestid Shrimp *Acetes japonicus* from the Estuary of Tanjung Dawai Peninsular Malaysia. *Journal of Fisheries and Aquatic Science*, 6(7), 771 - 779, (ISSN 1816-4927).
- Amani A A, A Arshad, S M Nurul Amin and Nor Azwady Abd Aziz, 2011. Catch Composition of a Set Bag Net (SBN) Used for *Acetes* Trapping in the Estuarine Waters of Kedah, Peninsular Malaysia. *Journal of Fisheries and Aquatic Science*, 6(3), 279 – 284 (ISSN 1816-4927).
- Ara, Roushon, 2011. Fish Larval Composition, Distribution, and Diets of the Seagrass-Mangrove Ecosystem of Gelang Patah, Johor, Peninsular Malaysia. PhD Thesis. Universiti Putra Malaysia. 175 pp.
- Ara R, Aziz Arshad, S.M.Nurul Amin, A.G. Mazlan and M. Aminur Rahman, 2013a. Feeding Ecology and Nursery of Marine Larval Fishes. *Asian Journal of Animal and Veterinary Advances*, 8,(2), 266 - 274 (ISSN 1683-9919).
- Ara R, S.M.Nurul Amin, A.G. Mazlan and A. Arshad, 2013b. Morphometric Variation Among Six Families of Larval Fishes in the Seagrass-Mangrove Ecosystem of Gelang Patah, Johor, Malaysia. *Asian Journal of Animal and Veterinary Advances*, 8(2), 247 - 256 (ISSN 1683-9919).
- Ara R, A. Arshad, S.M.N. Amin and A.G. Mazlan, 2013c. Temporal and Spatial Distribution of Larval Fish Assemblage in Different Ecological Habitat in Johor Strait, Peninsular Malaysia. *Asian Journal of Animal and Veterinary Advances*, 8(1), 53 - 62 (ISSN 1683-9919).
- Ara R, A Arshad, L. Musa, S.M.N. Amin and P. Kuppan, 2011a. Feeding Habits of Larval Fishes of the Family Clupeidae (Actinopterygii: Clupeiformes) in the Estuary of River Pendas, Johor, Malaysia. *Journal of Fisheries and Aquatic Science*, 6(7), 816 - 821 (ISSN 1816-4927).
- Ara R, A Arshad, S M N Amin, S K Daud, and Mazlan A G, 2011b. Comparison of Larval Fish Density between Seagrass Beds and Outside Seagrass Beds of the Southwestern Johor, Peninsular Malaysia. *Journal of Fisheries and Aquatic Science*, 6(7), 795 – 801 (ISSN 1816-4927).

- Ara R, Aziz Arshad, S M Nurul Amin, Siti Khalijah Daud and Mazlan Abdul Ghaffar, 2011c. Environment and Diversity of Ichthyoplankton in the Seagrass Beds of Sungai Pulai Estuary, Johor, Peninsular Malaysia. *Journal of Food Agriculture & Environment*, 9(3&4), 733 - 738 (ISSN: 1459-0263).
- Ara R, Aziz Arshad, S. M. Nurul Amin, Siti Khalijah Daud and Mazlan Abdul Ghaffar, 2010. Feeding Habits of Larval Fishes of Family Gobiidae (Perciformes: Gobiidae) in Seagrass Beds of Sungai Pulai Estuary, Johore Strait, Peninsular Malaysia. *Coastal Marine Science*, 34 (1), 123 - 128 (ISSN 1349-3000).
- Ara R, A Arshad, N Amrullah, S M Nurul Amin, S K Daud, A A Nor Azwady and A G Mazlan, 2009. Feeding Habit and Temporal Variation of Diet Composition of Fish Larvae (Osteichthys : Sparidae) in the Sungai Pulai Seagrass Bed, Johore, Peninsular Malaysia. *Journal of Biological Sciences*, 9(5), 445 – 451 (ISSN 1727-3048).
- Arshad A, R. Ara, S.M.N. Amin and A.G. Mazlan, 2013. Diet Composition in Larval Fishes of the Family Terapontidae (Actinopterygii: Perciformes) in the Seagrass-bed of Johor Strait, Malaysia. *Asian Journal of Animal and Veterinary Advances*, 8(2), 325 - 332 (ISSN 1683-9919).
- Arshad A, S.M.N. Amin, Wan Nurul Izzah, D. Aziz and R. Ara, 2013. Morphometric Variation Among the Populations of Planktonic Shrimp, *Acetes japonicus* in the West Coast of Peninsular Malaysia. *Asian Journal of Animal and Veterinary Advances*, 8(2), 194 - 204 (ISSN 1683-9919).
- Arshad A, S M Nurul Amin, Y L Z Nuradiella, Z C Cob, and D Aziz, 2012. Population Characteristics of *Acetes japonicus* from the Kedah Coastal Waters of Peninsular Malaysia. *Journal of Fisheries and Aquatic Science*, 7(2), 162 - 172 (ISSN 1816-4927).
- Arshad A, R Ara, S M N Amin, M Effendi, Zaidi C C, and Mazlan A G, 2011. Influence of Habitat Parameters on Shrimp Post-larvae in the Sungai Pulai Seagrass Beds of Johor Straits, Peninsular Malaysia. *Scientific Research and Essays*, 6(26), 5501 - 5506 (ISSN 1992- 2248).

- Arshad A, S M Nurul Amin, Norhafizah Osman and Che Roos Saad, 2010. Population Parameters of the Planktonic Shrimp *Lucifer intermedius* (Decapoda : Sergestidae) of Sg. Pulai Seagrass Beds, Johore, Peninsular Malaysia. *Sains Malaysiana*, 39(6), 877 – 882 (ISSN 0126-6039).
- Arshad A, S M Nurul Amin, Yu Guk Ting, Oh Siew Yong, Japar Sidik Bujang and Mazlan Abd Ghaffar, 2008. Population Characteristics, Length-Weight and Length-Length Relationships of *Acetes vulgaris* (Decapoda: Sergestidae) of the Coastal Waters of Pontian, Johore, Peninsular Malaysia. *Journal of Biological Sciences*, 8(8), 1298 – 1303 (ISSN 1562-2916).
- Arshad A, S. M. Nurul Amin, S. S. Siraj and S. B. Japar, 2007. New Distribution Records of Sergestid Shrimp, *Acetes intermedius* (Decapoda: Sergestidae) from Peninsular Malaysia with Notes on its Population Characteristics. *Journal of Biological Sciences*, 7(8), 1305 – 1313 (ISSN 1812-5719).
- Barwani, Said Mohamed, 2007. Temporal Variations in Reproduction, Growth, Condition Index and Some Population Parameters of *Perna viridis* (Linnaeus, 1758). PhD Thesis. Universiti Putra Malaysia. 186 pp.
- Barwani S M, A Arshad, S M N Amin and J S Bujang, 2013. Incidence of Hermaphrodite in Green Mussel *Perna viridis* along the West Coast of Peninsular Malaysia. *Asian Journal of Animal and Veterinary Advances*, 8(2), 376 - 382 (ISSN 1683-9919; Impact factor 2011: 0.869)
- Barwani S M, A Arshad, S M Nurul Amin, S. B. Japar, S S Siraj and C K Yap, 2007. Population Dynamics of the Green Mussel, *Perna viridis* from the High Spat-fall Coastal Waters of Malacca, Peninsular Malaysia. *Fisheries Research*, 84(2), 147 – 152 (ISSN 0165-7836).
- Chen, G.-Q., Xian, W.-Z., Lau, C.-C., Peng, J., Qiu, J.-W., Chen, F. and Jiang, Y., 2009. A comparative analysis of lipid and carotenoid composition of the gonads of *Anthocardaris crassisipina*, *Diadema setosum* and *Salinacis sphaeroides*. *Food Chemistry*, 120, 973–977.

- Choo, P.S. 1983. Mussel culture. SAFIS Extension Manual No. 3, Southeast Asian Fisheries Development Center, Bangkok, Thailand. pp.17.
- Cob, Zaidi Che, 2008. Biology and Ecology of Dog Conch (*Strombus canarium* Linnaeus, 1758) (Gastropoda: Strombidae) from Merambong Shoal, Johor Straits, Malaysia. PhD Thesis, Universiti Putra Malaysia. 377 pp.
- Cob Z C, A. Arshad, J. S. Bujang, Y. Bakar, K. D. Simon and A. G. Mazlan, 2012. Habitat Preference and Usage of *Strombus canarium* Linnaeus 1758 (Gastropoda : Strombidae) in Malaysian Seagrass Beds. *Italian Journal of Zoology*, 79(3), 459 – 467 (ISSN 1125-0003).
- Cob Z C, Aziz Arshad, Japar Sidik Bujang and Mazlan Abd. Ghaffar, 2011. Description and Evaluation of Imposex in *Strombus canarium* Linnaeus, 1758 (Gastropoda, Strombidae): A Potential Bio-indicator of Tributyltin Pollution. *Environmental Monitoring and Assessment*, 178(1-4), 393 – 400 (ISSN 1573-2959).
- Cob Z C, A Arshad, J S Bujang, W L W Muda, and M A Ghaffar, 2010. Metamorphosis Induction of the Dog Conch *Strombus canarium* (Gastropoda : Strombidae) Using Cues Associated With Conch Nursery Habitat. *Journal of Applied Sciences*, 10(8), 628 – 635 (ISSN 1812-5654).
- Cob Z C, Aziz Arshad, Japar Sidik Bujang and Mazlan Abd. Ghaffar, 2009a. Exploring the Use of Empirical Methods to Measure the Secondary Production of *Strombus canarium* (Gastropoda: Strombidae) Population in Johor Straits, Malaysia. *Sains Malaysiana*, 38(6), 817 – 825 (ISSN 0126-6039).
- Cob Z C, A. Arshad, J. S. Bujang and M. A. Ghaffar, 2009b. Age, Growth, Mortality and Population Structure of *Strombus canarium* (Gastropoda : Strombidae) : Variation in Male and Female Sub-Populations. *Journal of Applied Sciences*, 9(18), 3287 – 3297 (ISSN 1812-5654).
- Cob Z C, Aziz Arshad, Japar Sidik Bujang and Mazlan Abd Ghaffar, 2009c. Seasonal Variation in Growth and Survival of *Strombus canarium* (Linnaeus 1758) Larvae. *Pakistan Journal of Biological Sciences*, 12(9), 676 – 682 (ISSN 1028-8880).

- Cob Z C, Aziz Arshad, Japar Sidik Bujang and Mazlan Abdul Ghaffar, 2009d. Species Description and Distribution of *Strombus* (Mollusca: Strombidae) in Johor Straits and its Surrounding Areas. *Sains Malaysiana*, 38(1), 39 – 46 (ISSN 0126-6039).
- Cob Z C, Aziz Arshad, Mazlan Abd. Ghaffar, Japar Sidik Bujang, and Wan Lotfi Wan Muda, 2009e. Development and Growth of Larvae of The Dog Conch, *Strombus canarium* (Mollusca : Gastropoda) in Laboratory. *Zoological Studies*, 48(1), 1 – 11 (ISSN 1021-5506).
- Cob Z C, Aziz Arshad, Japar Sidik B., S.M. Nurul Amin and Mazlan Abd Ghaffar, 2008a. Growth, Mortality, Recruitment and Yield-per-Recruit of *Strombus canarium* Linnaeus, 1758 (Mesogastropoda: Strombidae) from the West Johor Straits, Malaysia. *Research Journal of Fisheries and Hydrology*, 3(2), 71 – 77. (ISSN 1816-9112).
- Cob Z C, Aziz Arshad, Japar Sidik Bujang, and Mazlan Abd. Ghaffar, 2008b. Sexual Maturity and Sex Determination in *Strombus canarium* Linnaeus, 1758 (Gastropoda : Strombidae). *Journal of Biological Sciences*, 8(3), 616 – 621 (ISSN 1812-5719).
- Cob Z C, Aziz Arshad, Japar Sidik Bujang and Mazlan Abd Ghaffar, 2008c. On the Biology and Basic Characteristics of the Population Dynamics of the Dog Conch *Strombus canarium* Linnaeus 1758 (Strombidae). *Journal of Bioscience*, 19(2), 81 – 97 (ISSN 0128-4541).
- Cob Z C, Aziz Arshad, Mohd Hanafi Idris, Japar Sidik Bujang, and Mazlan Abd. Ghaffar, 2008d. Sexual Polymorphism in a Population of *Strombus canarium* Linnaeus, 1758 (Mollusca: Gastropoda) at Merambong Shoal, Malaysia. *Zoological Studies*, 47(3), 318 – 325 (ISSN 1021-5506).
- Dania A, S. S. Siraj, A. Arshad, S.M. Nurul Amin and S.A. Harmin, 2010. Population Genetic Variation of Planktonic Shrimp, *Acetes japonicus* (Decapoda: Sergestidae) using RAPD Technique. *Journal of Biological Sciences*, 10(4), 355 – 361 (ISSN 1727-3048).
- Idris, Mohd Hanafi, 2009. Biology and Distribution of Pen Shells (Bivalvia : Pinnidae) in Selected Areas of Peninsular Malaysia. Phd Thesis. Universiti Putra Malaysia. 332 pp.

- Idris M H, A Arshad, S M N Amin, Japar S B, S K Daud, Mazlan A G, Azmi M A and M S Zakaria, 2012. Age, Growth and Length-Weight Relationships of *Pinna bicolor* Gmelin (Bivalvia: Pinnidae) in the Seagrass Beds of Sungai Pulai Estuary, Johor, Peninsular Malaysia. *Journal of Applied Ichthyology*, 28, 597 - 500 (ISSN 0175-8659).
- Idris M H, Aziz Arshad, Japar Sidik Bujang, Mazlan Abd Ghaffar and Siti Khalijah Daud, 2009. Morphological Characteristics of *Pinna bicolor* Gmelin and *Pinna deltodes* Menke from the Seagrass Beds of Sungai Pulai Johore Peninsular Malaysia. *Sains Malaysiana*, 38(3), 333 – 339 (ISSN 0126-6039).
- Idris M H, Aziz Arshad, Japar Sidik Bujang, Siti Khalijah Daud and Mazlan Abdul Ghaffar, 2008a. New Distribution Record of Two Pen Shells (Bivalvia : Pinnidae) From the Seagrass Beds of Sg. Pulai, Peninsular Malaysia. *Journal Biological Sciences*, 8(5), 882 – 888 (ISSN 1812-5719).
- Idris M H, Aziz Arshad, Japar Sidik Bujang, Mazlan Abd Ghaffar and Siti Khalijah Daud, 2008b. Morphometric Analysis as an Application Tool to Differentiate Three local Pen Shell Species. *Pertanika Journal Tropical Agricultural Science*, 31(2), 287 – 298 (ISSN – 1511-3701).
- Idris M H, Aziz Arshad, Japar Sidik Bujang, Mazlan Abd. Ghaffar and Siti Khalijah Daud, 2008c. Biodiversity and Distribution of Pen Shells (Bivalvia : Pinnidae) from the Seagrass Beds of Sungai Pulai, Peninsular Malaysia. *Research Journal of Fisheries and Hydrology*, 3(2), 54 -62. (ISSN 1816-9112).
- Kaneniwa, M. and Takagi, T. 1986. Fatty acids in the lipid of food products from sea urchin. *Bull. Jap. Soc. Sci. Fish.*, 52(9): 1681–1685.
- Masitah Saini, 2013. Feeding Habits, Recruitment and Status of Sergestid Shrimps (*Acetes intermedius* Omori) Stock in Bintulu Sarawak. Master Science Thesis. Universiti Putra Malaysia. 74 pp.
- Masitah S, Aziz Arshad, S M N Amin, and Mohd Hanafi Idris, 2011. Growth, Mortality and Recruitment of Planktonic Shrimp, *Acetes intermedius* in the Coastal Waters of Bintulu, Sarawak East Malaysia. *Journal of Fisheries and Aquatic Science*, 6(7), 780 - 787 (ISSN 1816-4927).

- Norhafizah Osman, 2013. Population Parameters and Feeding Biology of Sergestid shrimps, *Lucifer hansenii* from Tebrau Straits, Johor Malaysia. Master Thesis, Universiti Putra Malaysia.
- Nur Leena Wong Wai Sin, 2014. Distribution, Larval Abundance, Spawning and Early Development of Saddle Oyster, *Placuna ephippium* (Philipsson, 1788) From Merambong Shoal. PhD Thesis. Universiti Putra Malaysia. 145 pp.
- Nurul Amin, S M, 2008. Biology and Population of Sergestid Shrimps (*Acetes* spp.) (Decapoda : Sergestidae) from Klebang Besar, Malacca Malaysia. PhD Thesis. Universiti Putra Malaysia. 297 pp.
- Nurul Amin S M, Aziz Arshad, Siti Shapor Siraj and B. Japar Sidik, 2012. Population Biology and Stock Status of Planktonic Shrimp *Acetes indicus* (Decapoda: Sergestidae) in the Coastal Waters of Malacca, Peninsular Malaysia. *Journal of Ecosystem Health and Management* 15(3), 294 - 302 (ISSN 1463-4988).
- Nurul Amin S M, Aziz Arshad, Siti Shapor Siraj, and Japar Sidik Bujang, 2011. Update on the Species Composition and Distribution of Sergestid Shrimps (*Acetes* spp.) in Malaysian Waters. *Journal of Fisheries and Aquatic Science*, 6(7), 761 - 770 (ISSN 1816-4927).
- Nurul Amin S M, Arshad A., Siraj S S, and Bujang J S, 2010a. Reproductive Seasonality and Maturation of the Sergestid Shrimp, *A. japonicus* (Decapoda: Sergestidae) in the Coastal Waters of Malacca, Peninsular Malaysia. *African Journal of Biotechnology*, 9(45), 7747 – 7752 (ISSN 1684-5315).
- Nurul Amin S M, Aziz Arshad, Noor Hashida Ismail, Mohd Hanafi Idris, Japar Sidik Bujang and Siti Shapor Siraj, 2010b. Morphometric Variation Among the Three Species of the Genus *Acetes* (Decapoda : Sergestidae) in the Coastal Waters of Malacca, Peninsular Malaysia. *Pertanika Journal of Tropical Agricultural Science*, 33(2), 341 – 347; ISSN 1511-3701).
- Nurul Amin S M, Aziz Arshad, Japar Sidik Bujang, Siti Shapor Siraj and Stephen Goddard, 2009a. Reproductive Biology of Sergestid Shrimp *Acetes indicus* (Decapoda : Sergestidae) in the Coastal Waters of Malacca, Peninsular Malaysia. *Zoological Studies*, 48(6), 753 - 760 (ISSN 1021 - 5506).

- Nurul Amin S M, A Arshad, S S Siraj and S B Japar, 2009b. Population Structure, Growth and Length-Weight Relationship of Sergestid Shrimps (*Acetes* spp.) from the Coastal Waters of Malacca, Peninsular Malaysia. *Sains Malaysiana*, 38(2), 159 – 169 (ISSN 0126-6039).
- Nurul Amin S M, A Arshad, S S Siraj and S B Japar, 2009c. Population Structure, Growth, Mortality and Yield per Recruit of Sergestid Shrimp, *Acetes japonicus* (Decapoda : Sergestidae) from the Coastal Waters of Malacca, Peninsular Malaysia. *Indian Journal of Marine Sciences*, 38(1), 57-68 (ISSN 0379-5136).
- Nurul Amin S M, A. Arshad, Japar S B and S S Siraj, 2009d. Age Structure, Growth, Mortality and Yield per Recruit of Sergestid Shrimps *Acetes indicus* (Decapoda: Sergestidae) from the Coastal Waters of Malacca, Peninsular Malaysia. *Journal of Applied Sciences*, 9(5): 801 -814 (ISSN 1812-5654).
- Nurul Amin S M, A Arshad, Sarikha Shamsudin, Japar S B and S S Siraj, 2008a. Catch per Unit Effort of Estuarine Push Net with Emphasis on Occurrence and Abundance of *Acetes* Shrimps in the Coastal Waters of Malacca, Peninsular Malaysia. *Pertanika J. Sci. Technol.*, 16(2), 281 – 289 (ISSN 0128-7680).
- Nurul Amin S M, A Arshad, Japar S, B. and S S Siraj, 2008b. Growth, Mortality and Yield-Per-Recruit of Sergestid Shrimp, *Acetes intermedius* Omori, 1975 (Decapoda: Sergestidae) from Length Frequency Analysis in the Coastal Waters of Malacca, Peninsular Malaysia. *Pertanika Journal Tropical Agricultural Science*, 31(1), 95 – 106 (ISSN 1511-3701).
- Nurul Amin S M, A Arshad, Z Zainal, M H Idris, S S Siraj and S B Japar, 2008c. First Distribution Records of *Acetes intermedius* (Decapoda : Sergestidae) in Bintulu Coastal Waters: Population Structure, Length-Weight and Length-Length Relationship. *Journal of Sustainability Science and Management*, 3(1), 74 – 83 (ISSN 1985-3629).
- Nurul Amin S M, Aziz Arshad, Siti Shapor Siraj and Japar Sidik Bujang, 2007. The Biology and Life History of Sergestid Shrimp *Acetes indicus* (Decapoda : Sergestidae) in the Coastal Waters of Malacca, Peninsular Malaysia. *The Bioscientist*, 5(2), 9 – 17 (ISSN 1675-5618).

- Oh S Y, A Arshad, S B Japar, A A Nor Azwady and S M N Amin, 2011. Diet Composition of Sergestid Shrimp *Acetes serrulatus* from the Coastal Waters of Kukup, Johor Peninsular Malaysia. *Journal of Fisheries and Aquatic Science*, 6(7), 809 - 815 (ISSN 1816-4927).
- Oh S Y, A Arshad, S P Pang, and S M Nurul Amin, 2010. Catch Composition of Estuarine Set Bag Net Fishery in the Coastal Area of Pontian, Johor Peninsular Malaysia. *Journal Biological Sciences*, 10(3), 247 – 250 (ISSN 1727-3048).
- Pulz, O. and Gross, W. 2004. Valuable products from biotechnology of macroalgae. *Appl. Microbiol. Biotech.*, 65(6), 635–648.
- Rahman M A, Fatimah Md. Yusoff, A. Arshad, S.M.N. Amin and Mariana Nor Shamsudin, 2013. Population Characteristics and Fecundity Estimates of Short-spined White Sea Urchin, *Salmacis sphaeroides* (Linnaeus, 1758) from the Coastal Waters of Johor, Malaysia. *Asian Journal of Animal and Veterinary Advances*, 8(2), 301 - 308 (ISSN 1683-9919).
- Rahman M A, S M Nurul Amin, Perumal Kuppan, Aziz Arshad, Mariana Nor Shamsudin and Fatimah Md. Yusoff, 2012a. Length-Weight Relationships and Fecundity Estimates of Long-Spined Sea Urchin, *Diadema setosum* from Pulau Pangkor Waters, Peninsular Malaysia. *Journal of Ecosystem Health and Management*, 15(3), 311 – 315 (ISSN 1463-4988).
- Rahman M A, Fatimah Md. Yusoff, A. Arshad, Mariana Nor Shamsudin and S. M. Nurul Amin, 2012b. Embryonic, Larval and Early Juvenile Development of the Tropical Sea Urchin, *Salmacis sphaeroides* (Echinodermata: Echinoidea). *The Scientific World Journal*, Vol. 2012, Article ID 938482, 9 pp)
- Rahman, M.A. and Yusoff, F.M. 2010. Sea urchins in Malaysian coastal waters. *The Oceanographer*, 4, 20–21.



BIOGRAPHY

Aziz Arshad [PhD, Prof.] was born on the 9th of October 1959 in Arau, Perlis. He began his earlier primary education in Paya Primary School, Perlis from 1966 to 1972 and finished his secondary education at Derma Secondary School, Kangar Perlis in 1977. He graduated in Fisheries Science as the pioneer batch from the Universiti Pertanian Malaysia in March 1983, gained his MS degree from University of Wales Bangor in 1987 and obtained his doctorate in Marine Invertebrate Ecology in April 1999. He is currently serving as Professor of Fisheries Biology in the Department of Aquaculture, Faculty of Agriculture and filling up the role as an expert and resource person in Fisheries Biology.

To date, he has reached his 30 years of service with UPM. He has been fulfilling almost all the major responsibilities laid on him as a lecturer in UPM. He started to develop his teaching skill as an invertebrate biologist and invertebrate culture when first attached to the Faculty of Fisheries and Marine Science, UPM. At present faculty, he is entrusted to conduct courses such as Biology of Aquatic Invertebrates, and Fisheries Science for the undergraduate program. Besides teaching, he also supervised final year projects and to date more than 150 undergraduate students has completed their final year projects under his supervision. He has managed to graduate 9 PhD and 5 Master students as their principal supervisor.

As a researcher, fisheries biology is his main subject of interest. Typically, he tailored all his research projects to include fish as the main research subject. His focus of research for the last decade has been on the biology and ecology of crustacean and molluscs. His untiring efforts have been reflected by the intensive publications accumulated by him for the past five years. The research team where he is part of it has close linkages and networking with other

international scientists and had been invited in past to present the research findings at both local and international meetings.

In total, he has handled and completed 16 research projects during his entire career with UPM with majority of them are on the topic of fisheries biology. His involvement in research has further extended by several collaborative works with other research colleagues from outside country. His authority in fisheries biology and marine ecology subject has received some degree of recognition from other universities through their appointment as external examiner for postgraduate students and reviewer for academic curriculum. Throughout his engagement in academic activities, he has written and co-authored 150 papers in peer reviewed journals. He has attended various local and international conferences and seminars as well as participating in exhibition on research products that portrayed the highlight of his achievements in research. His expertise and interest has also been generously shared with the public particularly to fish farmers as well as extension workers on the culture protocols, seed production and larval management aspects of the invertebrate culture.

His proficiency and competence in the subject area is academically recognised and been entrusted to referee several papers for both national and international journals. He was also an assessor for MOSTI research proposal application under Agriculture (Aquaculture) and Sea to Space category at university and national levels respectively, a national panel member for the curriculum development for Higher Secondary School Examination (STPM) Biology Subject for the Ministry of Education.

On the contribution to the nation and public, he is a life member for Malaysian Fisheries Society (MFS) and Malaysian Society of Applied Biology (MSAB), an honorary treasurer of the Asian Fisheries Society till present. He indulged himself with

Aziz Arshad

non-governmental societies and contributed his expertise and effort to propel the beneficial fishery agenda forward. Throughout his involvement, he was assuming the post of treasurer, secretary and vice president of the Malaysian Fisheries Society from 1992 – 2006. He was also appointed to serve as chairperson and member of organising committee of both national and international meetings.

In 2006, he received the Excellent Service Award from the university. He has been fairly prudent on his work performance. Since 2001, he has consistently been achieving an excellent status performance for his services to the department, faculty and university. In 2007, he won the Vice Chancellor University Excellence Award for Best in Professional Services. Despite all the past achievements, he is aspired towards superior contribution and achievement in the academic excellence of his career.



ACKNOWLEDGEMENTS

The author would like to take the opportunity to thank Universiti Putra Malaysia for providing him with the platform to channel all his research interests and deep attraction on the subject of fisheries biology. This has been made feasible through the provision of infrastructures, equipments, manpower and grant support. Throughout his 30 years of service, he has managed to dedicate his time and efforts to explore and investigate various aspects of biology and ecology of the marine invertebrates and fish that are of importance to the fishery sector of the country.

He would also like to extend his deep appreciation and gratitude to all the co-researchers in UPM from the Department of Aquaculture (Fatimah Md Yusoff, Muta Harah Zakaria, S M Nurul Amin, and Aminur Rahman) and UPMKB (Japar Sidik Bujang, Mohd Hanafi Idris) and in UKM's Faculty of Science and Technology (Mazlan Abd Ghaffar and Zaidi Che Cob) who have been together with him during the research period, contributing in term of innovative ideas and suggestive comments on the implementation of the projects. Without their cooperation and supports, it would be impossible for him to carry forward the research agendas with great success.

He would also like to thank his entire postgraduate researchers who had graduated and worked alongside him with full dedication and untiring efforts to ensure the smooth delivery of the research outputs. To them all, every single person has shown tremendous sacrifice and utmost dedication with his/her publication efforts and words alone cannot express their contribution to his research career and productivity in UPM.

There will be no success without the pillar of the family supports. To that, the author would like to honour his wife, Norhan Abu Bakar for her moral supports, patience, encouragement and continuous shower of loves and understanding. Their five children

(Dania, Ainan, Iylia, Azmil and Azim) have always being the source of constant inspiration to his career advancement in the university. Last but not least, in memory of both parents who are not able to share the moments with him but whose contribution has never been out of his thought and appreciation.

LIST OF INAUGURAL LECTURES

1. Prof. Dr. Sulaiman M. Yassin
The Challenge to Communication Research in Extension
22 July 1989
2. Prof. Ir. Abang Abdullah Abang Ali
Indigenous Materials and Technology for Low Cost Housing
30 August 1990
3. Prof. Dr. Abdul Rahman Abdul Razak
Plant Parasitic Nematodes, Lesser Known Pests of Agricultural Crops
30 January 1993
4. Prof. Dr. Mohamed Suleiman
Numerical Solution of Ordinary Differential Equations: A Historical Perspective
11 December 1993
5. Prof. Dr. Mohd. Ariff Hussein
Changing Roles of Agricultural Economics
5 March 1994
6. Prof. Dr. Mohd. Ismail Ahmad
Marketing Management: Prospects and Challenges for Agriculture
6 April 1994
7. Prof. Dr. Mohamed Mahyuddin Mohd. Dahan
The Changing Demand for Livestock Products
20 April 1994
8. Prof. Dr. Ruth Kiew
Plant Taxonomy, Biodiversity and Conservation
11 May 1994
9. Prof. Ir. Dr. Mohd. Zohadie Bardaie
Engineering Technological Developments Propelling Agriculture into the 21st Century
28 May 1994
10. Prof. Dr. Shamsuddin Jusop
Rock, Mineral and Soil
18 June 1994

Exploring Biodiversity and Fisheries Biology

11. Prof. Dr. Abdul Salam Abdullah
Natural Toxicants Affecting Animal Health and Production
29 June 1994
12. Prof. Dr. Mohd. Yusof Hussein
Pest Control: A Challenge in Applied Ecology
9 July 1994
13. Prof. Dr. Kapt. Mohd. Ibrahim Haji Mohamed
Managing Challenges in Fisheries Development through Science and Technology
23 July 1994
14. Prof. Dr. Hj. Amat Juhari Moain
Sejarah Keagungan Bahasa Melayu
6 Ogos 1994
15. Prof. Dr. Law Ah Theem
Oil Pollution in the Malaysian Seas
24 September 1994
16. Prof. Dr. Md. Nordin Hj. Lajis
Fine Chemicals from Biological Resources: The Wealth from Nature
21 January 1995
17. Prof. Dr. Sheikh Omar Abdul Rahman
Health, Disease and Death in Creatures Great and Small
25 February 1995
18. Prof. Dr. Mohamed Shariff Mohamed Din
Fish Health: An Odyssey through the Asia - Pacific Region
25 March 1995
19. Prof. Dr. Tengku Azmi Tengku Ibrahim
Chromosome Distribution and Production Performance of Water Buffaloes
6 May 1995
20. Prof. Dr. Abdul Hamid Mahmood
Bahasa Melayu sebagai Bahasa Ilmu- Cabaran dan Harapan
10 Jun 1995

Aziz Arshad

21. Prof. Dr. Rahim Md. Sail
Extension Education for Industrialising Malaysia: Trends, Priorities and Emerging Issues
22 July 1995
22. Prof. Dr. Nik Muhammad Nik Abd. Majid
The Diminishing Tropical Rain Forest: Causes, Symptoms and Cure
19 August 1995
23. Prof. Dr. Ang Kok Jee
The Evolution of an Environmentally Friendly Hatchery Technology for Udang Galah, the King of Freshwater Prawns and a Glimpse into the Future of Aquaculture in the 21st Century
14 October 1995
24. Prof. Dr. Sharifuddin Haji Abdul Hamid
Management of Highly Weathered Acid Soils for Sustainable Crop Production
28 October 1995
25. Prof. Dr. Yu Swee Yean
Fish Processing and Preservation: Recent Advances and Future Directions
9 December 1995
26. Prof. Dr. Rosli Mohamad
Pesticide Usage: Concern and Options
10 February 1996
27. Prof. Dr. Mohamed Ismail Abdul Karim
Microbial Fermentation and Utilization of Agricultural Bioresources and Wastes in Malaysia
2 March 1996
28. Prof. Dr. Wan Sulaiman Wan Harun
Soil Physics: From Glass Beads to Precision Agriculture
16 March 1996
29. Prof. Dr. Abdul Aziz Abdul Rahman
Sustained Growth and Sustainable Development: Is there a Trade-Off 1 or Malaysia
13 April 1996

Exploring Biodiversity and Fisheries Biology

30. Prof. Dr. Chew Tek Ann
Sharecropping in Perfectly Competitive Markets: A Contradiction in Terms
27 April 1996
31. Prof. Dr. Mohd. Yusuf Sulaiman
Back to the Future with the Sun
18 May 1996
32. Prof. Dr. Abu Bakar Salleh
Enzyme Technology: The Basis for Biotechnological Development
8 June 1996
33. Prof. Dr. Kamel Ariffin Mohd. Atan
The Fascinating Numbers
29 June 1996
34. Prof. Dr. Ho Yin Wan
Fungi: Friends or Foes
27 July 1996
35. Prof. Dr. Tan Soon Guan
Genetic Diversity of Some Southeast Asian Animals: Of Buffaloes and Goats and Fishes Too
10 August 1996
36. Prof. Dr. Nazaruddin Mohd. Jali
Will Rural Sociology Remain Relevant in the 21st Century?
21 September 1996
37. Prof. Dr. Abdul Rani Bahaman
Leptospirosis-A Model for Epidemiology, Diagnosis and Control of Infectious Diseases
16 November 1996
38. Prof. Dr. Marziah Mahmood
Plant Biotechnology - Strategies for Commercialization
21 December 1996
39. Prof. Dr. Ishak Hj. Omar
Market Relationships in the Malaysian Fish Trade: Theory and Application
22 March 1997

Aziz Arshad

40. Prof. Dr. Suhaila Mohamad
Food and Its Healing Power
12 April 1997
41. Prof. Dr. Malay Raj Mukerjee
A Distributed Collaborative Environment for Distance Learning Applications
17 June 1998
42. Prof. Dr. Wong Kai Choo
Advancing the Fruit Industry in Malaysia: A Need to Shift Research Emphasis
15 May 1999
43. Prof. Dr. Aini Ideris
Avian Respiratory and Immunosuppressive Diseases- A Fatal Attraction
10 July 1999
44. Prof. Dr. Sariah Meon
Biological Control of Plant Pathogens: Harnessing the Richness of Microbial Diversity
14 August 1999
45. Prof. Dr. Azizah Hashim
The Endomycorrhiza: A Futile Investment?
23 Oktober 1999
46. Prof. Dr. Noraini Abdul Samad
Molecular Plant Virology: The Way Forward
2 February 2000
47. Prof. Dr. Muhamad Awang
Do We Have Enough Clean Air to Breathe?
7 April 2000
48. Prof. Dr. Lee Chnoong Kheng
Green Environment, Clean Power
24 June 2000
49. Prof. Dr. Mohd. Ghazali Mohayidin
Managing Change in the Agriculture Sector: The Need for Innovative Educational Initiatives
12 January 2002

Exploring Biodiversity and Fisheries Biology

50. Prof. Dr. Fatimah Mohd. Arshad
*Analisis Pemasaran Pertanian di Malaysia: Keperluan Agenda
Pembaharuan*
26 Januari 2002
51. Prof. Dr. Nik Mustapha R. Abdullah
*Fisheries Co-Management: An Institutional Innovation Towards
Sustainable Fisheries Industry*
28 February 2002
52. Prof. Dr. Gulam Rusul Rahmat Ali
Food Safety: Perspectives and Challenges
23 March 2002
53. Prof. Dr. Zaharah A. Rahman
*Nutrient Management Strategies for Sustainable Crop Production in Acid
Soils: The Role of Research Using Isotopes*
13 April 2002
54. Prof. Dr. Maisom Abdullah
Productivity Driven Growth: Problems & Possibilities
27 April 2002
55. Prof. Dr. Wan Omar Abdullah
*Immunodiagnosis and Vaccination for Brugian Filariasis: Direct Rewards
from Research Investments*
6 June 2002
56. Prof. Dr. Syed Tajuddin Syed Hassan
Agro-ento Bioinformation: Towards the Edge of Reality
22 June 2002
57. Prof. Dr. Dahlan Ismail
*Sustainability of Tropical Animal-Agricultural Production Systems:
Integration of Dynamic Complex Systems*
27 June 2002
58. Prof. Dr. Ahmad Zubaidi Baharumshah
The Economics of Exchange Rates in the East Asian Countries
26 October 2002
59. Prof. Dr. Shaik Md. Noor Alam S.M. Hussain
Contractual Justice in Asean: A Comparative View of Coercion
31 October 2002

Aziz Arshad

60. Prof. Dr. Wan Md. Zin Wan Yunus
Chemical Modification of Polymers: Current and Future Routes for Synthesizing New Polymeric Compounds
9 November 2002
61. Prof. Dr. Annuar Md. Nassir
Is the KLSE Efficient? Efficient Market Hypothesis vs Behavioural Finance
23 November 2002
62. Prof. Ir. Dr. Radin Umar Radin Sohadi
Road Safety Interventions in Malaysia: How Effective Are They?
21 February 2003
63. Prof. Dr. Shamsher Mohamad
The New Shares Market: Regulatory Intervention, Forecast Errors and Challenges
26 April 2003
64. Prof. Dr. Han Chun Kwong
Blueprint for Transformation or Business as Usual? A Structural Perspective of the Knowledge-Based Economy in Malaysia
31 May 2003
65. Prof. Dr. Mawardi Rahmani
Chemical Diversity of Malaysian Flora: Potential Source of Rich Therapeutic Chemicals
26 July 2003
66. Prof. Dr. Fatimah Md. Yusoff
An Ecological Approach: A Viable Option for Aquaculture Industry in Malaysia
9 August 2003
67. Prof. Dr. Mohamed Ali Rajion
The Essential Fatty Acids-Revisited
23 August 2003
68. Prof. Dr. Azhar Md. Zain
Psychotherapy for Rural Malays - Does it Work?
13 September 2003

Exploring Biodiversity and Fisheries Biology

69. Prof. Dr. Mohd. Zamri Saad
Respiratory Tract Infection: Establishment and Control
27 September 2003
70. Prof. Dr. Jinap Selamat
Cocoa-Wonders for Chocolate Lovers
14 February 2004
71. Prof. Dr. Abdul Halim Shaari
High Temperature Superconductivity: Puzzle & Promises
13 March 2004
72. Prof. Dr. Yaakob Che Man
Oils and Fats Analysis - Recent Advances and Future Prospects
27 March 2004
73. Prof. Dr. Kaida Khalid
Microwave Aquametry: A Growing Technology
24 April 2004
74. Prof. Dr. Hasanah Mohd. Ghazali
Tapping the Power of Enzymes- Greening the Food Industry
11 May 2004
75. Prof. Dr. Yusof Ibrahim
The Spider Mite Saga: Quest for Biorational Management Strategies
22 May 2004
76. Prof. Datin Dr. Sharifah Md. Nor
The Education of At-Risk Children: The Challenges Ahead
26 June 2004
77. Prof. Dr. Ir. Wan Ishak Wan Ismail
Agricultural Robot: A New Technology Development for Agro-Based Industry
14 August 2004
78. Prof. Dr. Ahmad Said Sajap
Insect Diseases: Resources for Biopesticide Development
28 August 2004

Aziz Arshad

79. Prof. Dr. Aminah Ahmad
The Interface of Work and Family Roles: A Quest for Balanced Lives
11 March 2005
80. Prof. Dr. Abdul Razak Alimon
Challenges in Feeding Livestock: From Wastes to Feed
23 April 2005
81. Prof. Dr. Haji Azimi Hj. Hamzah
Helping Malaysian Youth Move Forward: Unleashing the Prime Enablers
29 April 2005
82. Prof. Dr. Rasedee Abdullah
In Search of An Early Indicator of Kidney Disease
27 May 2005
83. Prof. Dr. Zulkifli Hj. Shamsuddin
Smart Partnership: Plant-Rhizobacteria Associations
17 June 2005
84. Prof. Dr. Mohd Khanif Yusop
From the Soil to the Table
1 July 2005
85. Prof. Dr. Annuar Kassim
Materials Science and Technology: Past, Present and the Future
8 July 2005
86. Prof. Dr. Othman Mohamed
Enhancing Career Development Counselling and the Beauty of Career Games
12 August 2005
87. Prof. Ir. Dr. Mohd Amin Mohd Soom
Engineering Agricultural Water Management Towards Precision Framing
26 August 2005
88. Prof. Dr. Mohd Arif Syed
Bioremediation-A Hope Yet for the Environment?
9 September 2005

Exploring Biodiversity and Fisheries Biology

89. Prof. Dr. Abdul Hamid Abdul Rashid
The Wonder of Our Neuromotor System and the Technological Challenges They Pose
23 December 2005
90. Prof. Dr. Norhani Abdullah
Rumen Microbes and Some of Their Biotechnological Applications
27 January 2006
91. Prof. Dr. Abdul Aziz Saharee
Haemorrhagic Septicaemia in Cattle and Buffaloes: Are We Ready for Freedom?
24 February 2006
92. Prof. Dr. Kamariah Abu Bakar
Activating Teachers' Knowledge and Lifelong Journey in Their Professional Development
3 March 2006
93. Prof. Dr. Borhanuddin Mohd. Ali
Internet Unwired
24 March 2006
94. Prof. Dr. Sundararajan Thilagar
Development and Innovation in the Fracture Management of Animals
31 March 2006
95. Prof. Dr. Zainal Aznam Md. Jelani
Strategic Feeding for a Sustainable Ruminant Farming
19 May 2006
96. Prof. Dr. Mahiran Basri
Green Organic Chemistry: Enzyme at Work
14 July 2006
97. Prof. Dr. Malik Hj. Abu Hassan
Towards Large Scale Unconstrained Optimization
20 April 2007
98. Prof. Dr. Khalid Abdul Rahim
Trade and Sustainable Development: Lessons from Malaysia's Experience
22 Jun 2007

Aziz Arshad

99. Prof. Dr. Mad Nasir Shamsudin
Econometric Modelling for Agricultural Policy Analysis and Forecasting: Between Theory and Reality
13 July 2007
100. Prof. Dr. Zainal Abidin Mohamed
Managing Change - The Fads and The Realities: A Look at Process Reengineering, Knowledge Management and Blue Ocean Strategy
9 November 2007
101. Prof. Ir. Dr. Mohamed Daud
Expert Systems for Environmental Impacts and Ecotourism Assessments
23 November 2007
102. Prof. Dr. Saleha Abdul Aziz
Pathogens and Residues; How Safe is Our Meat?
30 November 2007
103. Prof. Dr. Jayum A. Jawan
Hubungan Sesama Manusia
7 Disember 2007
104. Prof. Dr. Zakariah Abdul Rashid
Planning for Equal Income Distribution in Malaysia: A General Equilibrium Approach
28 December 2007
105. Prof. Datin Paduka Dr. Khatijah Yusoff
Newcastle Disease virus: A Journey from Poultry to Cancer
11 January 2008
106. Prof. Dr. Dzulkefly Kuang Abdullah
Palm Oil: Still the Best Choice
1 February 2008
107. Prof. Dr. Elias Saion
Probing the Microscopic Worlds by Ionizing Radiation
22 February 2008
108. Prof. Dr. Mohd Ali Hassan
Waste-to-Wealth Through Biotechnology: For Profit, People and Planet
28 March 2008

Exploring Biodiversity and Fisheries Biology

109. Prof. Dr. Mohd Maarof H. A. Moxsin
Metrology at Nanoscale: Thermal Wave Probe Made It Simple
11 April 2008
110. Prof. Dr. Dzolkhifli Omar
The Future of Pesticides Technology in Agriculture: Maximum Target Kill with Minimum Collateral Damage
25 April 2008
111. Prof. Dr. Mohd. Yazid Abd. Manap
Probiotics: Your Friendly Gut Bacteria
9 May 2008
112. Prof. Dr. Hamami Sahri
Sustainable Supply of Wood and Fibre: Does Malaysia have Enough?
23 May 2008
113. Prof. Dato' Dr. Makhdzir Mardan
Connecting the Bee Dots
20 June 2008
114. Prof. Dr. Maimunah Ismail
Gender & Career: Realities and Challenges
25 July 2008
115. Prof. Dr. Nor Aripin Shamaan
Biochemistry of Xenobiotics: Towards a Healthy Lifestyle and Safe Environment
1 August 2008
116. Prof. Dr. Mohd Yunus Abdullah
Penjagaan Kesihatan Primer di Malaysia: Cabaran Prospek dan Implikasi dalam Latihan dan Penyelidikan Perubatan serta Sains Kesihatan di Universiti Putra Malaysia
8 Ogos 2008
117. Prof. Dr. Musa Abu Hassan
Memfaatkan Teknologi Maklumat & Komunikasi ICT untuk Semua
15 Ogos 2008
118. Prof. Dr. Md. Salleh Hj. Hassan
Role of Media in Development: Strategies, Issues & Challenges
22 August 2008

Aziz Arshad

119. Prof. Dr. Jariah Masud
Gender in Everyday Life
10 October 2008
120. Prof. Dr. Mohd Shahwahid Haji Othman
Mainstreaming Environment: Incorporating Economic Valuation and Market-Based Instruments in Decision Making
24 October 2008
121. Prof. Dr. Son Radu
Big Questions Small Worlds: Following Diverse Vistas
31 Oktober 2008
122. Prof. Dr. Russly Abdul Rahman
Responding to Changing Lifestyles: Engineering the Convenience Foods
28 November 2008
123. Prof. Dr. Mustafa Kamal Mohd Shariff
Aesthetics in the Environment an Exploration of Environmental: Perception Through Landscape Preference
9 January 2009
124. Prof. Dr. Abu Daud Silong
Leadership Theories, Research & Practices: Farming Future Leadership Thinking
16 January 2009
125. Prof. Dr. Azni Idris
Waste Management, What is the Choice: Land Disposal or Biofuel?
23 January 2009
126. Prof. Dr. Jamilah Bakar
Freshwater Fish: The Overlooked Alternative
30 January 2009
127. Prof. Dr. Mohd. Zobir Hussein
The Chemistry of Nanomaterial and Nanobiomaterial
6 February 2009
128. Prof. Ir. Dr. Lee Teang Shui
Engineering Agricultural: Water Resources
20 February 2009

Exploring Biodiversity and Fisheries Biology

129. Prof. Dr. Ghizan Saleh
Crop Breeding: Exploiting Genes for Food and Feed
6 March 2009
130. Prof. Dr. Muzafar Shah Habibullah
Money Demand
27 March 2009
131. Prof. Dr. Karen Anne Crouse
In Search of Small Active Molecules
3 April 2009
132. Prof. Dr. Turiman Suandi
Volunteerism: Expanding the Frontiers of Youth Development
17 April 2009
133. Prof. Dr. Arbakariya Ariff
Industrializing Biotechnology: Roles of Fermentation and Bioprocess Technology
8 Mei 2009
134. Prof. Ir. Dr. Desa Ahmad
Mechanics of Tillage Implements
12 Jun 2009
135. Prof. Dr. W. Mahmood Mat Yunus
Photothermal and Photoacoustic: From Basic Research to Industrial Applications
10 Julai 2009
136. Prof. Dr. Taufiq Yap Yun Hin
Catalysis for a Sustainable World
7 August 2009
137. Prof. Dr. Raja Noor Zaliha Raja Abd. Rahman
Microbial Enzymes: From Earth to Space
9 Oktober 2009
138. Prof. Ir. Dr. Barkawi Sahari
Materials, Energy and CNGDI Vehicle Engineering
6 November 2009

Aziz Arshad

139. Prof. Dr. Zulkifli Idrus
Poultry Welfare in Modern Agriculture: Opportunity or Threat?
13 November 2009
140. Prof. Dr. Mohamed Hanafi Musa
Managing Phosphorus: Under Acid Soils Environment
8 January 2010
141. Prof. Dr. Abdul Manan Mat Jais
Haruan Channa striatus a Drug Discovery in an Agro-Industry Setting
12 March 2010
142. Prof. Dr. Bujang bin Kim Huat
Problematic Soils: In Search for Solution
19 March 2010
143. Prof. Dr. Samsinar Md Sidin
Family Purchase Decision Making: Current Issues & Future Challenges
16 April 2010
144. Prof. Dr. Mohd Adzir Mahdi
Lightspeed: Catch Me If You Can
4 June 2010
145. Prof. Dr. Raha Hj. Abdul Rahim
Designer Genes: Fashioning Mission Purposed Microbes
18 June 2010
146. Prof. Dr. Hj. Hamidon Hj. Basri
A Stroke of Hope, A New Beginning
2 July 2010
147. Prof. Dr. Hj. Kamaruzaman Jusoff
Going Hyperspectral: The "Unseen" Captured?
16 July 2010
148. Prof. Dr. Mohd Sapuan Salit
Concurrent Engineering for Composites
30 July 2010
149. Prof. Dr. Shattri Mansor
Google the Earth: What's Next?
15 October 2010

Exploring Biodiversity and Fisheries Biology

150. Prof. Dr. Mohd Basyaruddin Abdul Rahman
Haute Couture: Molecules & Biocatalysts
29 October 2010
151. Prof. Dr. Mohd. Hair Bejo
Poultry Vaccines: An Innovation for Food Safety and Security
12 November 2010
152. Prof. Dr. Umi Kalsom Yusuf
Fern of Malaysian Rain Forest
3 December 2010
153. Prof. Dr. Ab. Rahim Bakar
*Preparing Malaysian Youths for The World of Work: Roles of Technical
and Vocational Education and Training (TVET)*
14 January 2011
154. Prof. Dr. Seow Heng Fong
Are there "Magic Bullets" for Cancer Therapy?
11 February 2011
155. Prof. Dr. Mohd Azmi Mohd Lila
Biopharmaceuticals: Protection, Cure and the Real Winner
18 February 2011
156. Prof. Dr. Siti Shapor Siraj
Genetic Manipulation in Farmed Fish: Enhancing Aquaculture Production
25 March 2011
157. Prof. Dr. Ahmad Ismail
Coastal Biodiversity and Pollution: A Continuous Conflict
22 April 2011
158. Prof. Ir. Dr. Norman Mariun
Energy Crisis 2050? Global Scenario and Way Forward for Malaysia
10 June 2011
159. Prof. Dr. Mohd Razi Ismail
Managing Plant Under Stress: A Challenge for Food Security
15 July 2011

Aziz Arshad

160. Prof. Dr. Patimah Ismail
Does Genetic Polymorphisms Affect Health?
23 September 2011
161. Prof. Dr. Sidek Ab. Aziz
Wonders of Glass: Synthesis, Elasticity and Application
7 October 2011
162. Prof. Dr. Azizah Osman
Fruits: Nutritious, Colourful, Yet Fragile Gifts of Nature
14 October 2011
163. Prof. Dr. Mohd. Fauzi Ramlan
Climate Change: Crop Performance and Potential
11 November 2011
164. Prof. Dr. Adem Kiliçman
Mathematical Modeling with Generalized Function
25 November 2011
165. Prof. Dr. Fauziah Othman
My Small World: In Biomedical Research
23 December 2011
166. Prof. Dr. Japar Sidik Bujang
The Marine Angiosperms, Seagrass
23 March 2012
166. Prof. Dr. Zailina Hashim
Air Quality and Children's Environmental Health: Is Our Future Generation at Risk?
30 March 2012
167. Prof. Dr. Zainal Abidin Mohamed
Where is the Beef? Vantage Point form the Livestock Supply Chain
27 April 2012
168. Prof. Dr. Jothi Malar Panandam
Genetic Characterisation of Animal Genetic Resources for Sustainable Utilisation and Development
30 November 2012

Exploring Biodiversity and Fisheries Biology

169. Prof. Dr. Fatimah Abu Bakar
The Good The Bad & Ugly of Food Safety: From Molecules to Microbes
7 Desember 2012
170. Prof. Dr. Abdul Jalil Nordin
My Colourful Sketches from Scratch: Molecular Imaging
5 April 2013
171. Prof. Dr. Norlijah Othman
*Lower Respiratory Infections in Children: New Pathogens, Old Pathogens
and The Way Forward*
19 April 2013
172. Prof. Dr. Jayakaran Mukundan
Steroid-like Prescriptions English Language Teaching Can Ill-afford
26 April 2013
173. Prof. Dr. Azmi Zakaria
Photothermals Affect Our Lives
7 Jun 2013
174. Prof. Dr. Rahinah Ibrahim
Design Informatics
21 Jun 2103
175. Prof. Dr. Gwendoline Ee Cheng
Natural Products from Malaysian Rainforests
1 November 2013
176. Prof. Dr. Noor Akma Ibrahim
The Many Facets of Statistical Modeling
22 November 2013
177. Prof. Dr. Paridah Md. Tahir
Bonding with Natural Fibres
6 Desember 2013
178. Prof. Dr. Abd. Wahid Haron
Livestock Breeding: The Past, The Present and The Future
9 Desember 2013