



UNIVERSITI PUTRA MALAYSIA

**IDENTIFICATION OF INFECTIOUS BRONCHITIS VIRUS ISOLATES
FROM MALAYSIA**

BALKIS BINTI HAJI A. TALIP

FPV 2007 4



**IDENTIFICATION OF INFECTIOUS BRONCHITIS VIRUS
ISOLATES FROM MALAYSIA**

By

BALKIS BINTI HAJI A. TALIP

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

2007



DEDICATED TO.....

My Father,

HAJI A TALIP BIN MD NOH

My Mother,

HAJJAH ZAHARAH@ROHANI BINTI JA'AMAT@YA'AMAT

My sisters,

BAHIYAH

BAKHREZA

BAZILAH

BAIYINAH

My very best friend,

LATIFAH BTE MD ARIFFIN



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

**IDENTIFICATION OF INFECTIOUS BRONCHITIS VIRUS
ISOLATES FROM MALAYSIA**

By

BALKIS BINTI HAJI A. TALIP

December 2006

Chairman: Associate Professor Dr. Siti Suri Arshad, PhD

Faculty: Veterinary Medicine

Infectious bronchitis (IB) is a highly contagious respiratory, urogenital and reproduction disease of chickens and it is distributed worldwide. The disease is caused by infectious bronchitis virus (IBV). IBV is a member of the genus *Coronavirus*, family *Coronaviridae* and it has a single-stranded, positive-sense RNA genome of 27.6 kb. It possesses prominent surface spikes and has three major structural proteins; the spike (S) glycoprotein, the small integral membrane (M) glycoprotein and nucleocapsid (N) protein. In the commercial poultry industry, vaccination is used to control the disease. Despite vaccination program the disease continues to occur because IBV can exist in many serotypes. In many incidences, the existing vaccines are not able to provide full protection to the chickens against infectious bronchitis disease. The immune response stimulated to one serotype does not offer cross protection to another serotype. Moreover, the avian coronavirus capable of mutating and many IBV variants has been reported in many countries. Thus, it is crucial to develop a fast, sensitive and specific diagnostic technique to diagnose and identify the causative agent in order to control the disease



spread. In recent years the reverse transcriptase-polymerase chain reaction (RT-PCR), reverse transcriptase-polymerase chain reaction-restriction fragment length polymorphism (RT-PCR-RFLP), cloning and genes sequencing had been used increasingly to detect and analyses IBV isolates. The objectives of these studies were to compare and optimize reverse transcriptase-polymerase chain reaction (RT-PCR) to diagnose IBV, to differentiate the Mass strain, and to characterize variant IBV isolated from this study. In this study, one-step and two-step RT-PCR techniques were used to amplify the conserved gene region of IBV by using universal and designed primers. This study was conducted on IBV isolates from year 1991 until 2003. In differentiation studies, isolates were group into serotype using reverse transcriptase-polymerase chain reaction (RT-PCR). Isolates recognized to be non Mass was further amplified their hypervariable region of S gene by using RT-PCR followed by RFLP technique to screen for nephropathogenic strain. Out of 31 IBV isolates, nine Mass strain were found. Following RT-PCR-RFLP, only one isolate showed different fragment pattern compared to nephropathogenic origin (MH5365/95). This particular isolates designated as V9/03 was neither Mass nor non-nephropathogenic serotype. The S1 region of V9/03 was further cloned, sequenced and its nucleotide and amino acid were compared to nephropathogenic MH5365/95 and Mass derivatives as well other references strains obtained from Gene bank. The V9/03 showed sequence homology to Taiwan (AY606321) and Korean (AY257060) strain with 82.5% and 81.6% identities respectively. The V9/03 has lower sequences homology (less than 80%) with nephropathogenic (MH5365/95) and Mass derivatives. The phylogenetic studies indicate that the strain V9/03 could be a local variant IBV which is different from local



nephropathogenic MH5365/95 and vaccine strain. This study showed that variant IBV is circulating in the field as result of mutation of IBV due to the prolonged use of live virus vaccines and the immunological pressure of the virus to keep on survival in immune birds.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGENALPASTIAN ISOLAT-ISOLAT VIRUS BRONKITIS BERJANGKIT
DI MALAYSIA**

Oleh

BALKIS BINTI HAJI A TALIP

December 2006

Pengerusi: Profesor Madya Dr.Siti Suri Arshad, PhD

Fakulti: Perubatan Veterinar

Bronkitis berjangkit merupakan penyakit pernafasan ayam yang menular ke serata dunia. Ianya disebabkan oleh virus bronkitis berjangkit (IBV). Virus bronkitis berjangkit (IB) merupakan ahli genera *Coronavirus*, famili *Coronaviridae*. Virus ini merupakan virus bebenang tunggal, RNA genom berpancaindera-positif bersaiz 27.6 kb. Ianya mempunyai permukaan tajam yang menonjol dan mempunyai tiga protein berstruktur yang utama; glikoprotein pepaku (S), glikoprotein membran kecil perlu (M) dan protein nukleokapsid (N). Dalam industri penternakan ayam, vaksinasi digunakan untuk mengawal penyakit. Walaupun program vaksinasi, penyakit terus berlaku kerana IBV boleh wujud di dalam pelbagai serotip. Dalam kebanyakan kejadian, kewujudan vaksin sahaja tidak mencukupi untuk menyediakan perlindungan sepenuhnya kepada ayam-ayam bagi menentang IBV. Tindakbalas imuniti terhadap satu serotip tidak boleh menawarkan perlindungan bersilang kepada serotip yang lain. Lagipun, coronavirus avian berkebolehan untuk mutasi dan banyak varian IBV telah dilaporkan di beberapa



negara. Demikian, adalah sangat genting agen penyebab dikenalpasti untuk mengawal penyebaran penyakit. Sejak akhir-akhir ini, RT-PCR, RT-PCR-RFLP, pengklonan dan penjujukan gen telah banyak digunakan untuk mengesan dan analisis isolat IBV. Objektif-objektif dalam kajian ini adalah untuk membina dan optimasi RT-PCR untuk diagnosa IBV, mengasingkan strain Mass dan untuk mengenalpasti IBV varian terpicil dalam kajian ini. Dalam kajian ini, teknik satu langkah dan 2-langkah RT-PCR telah digunakan untuk menggandakan gen yang majmuk bagi IBV dengan menggunakan primer universal dan direka. Kajian ini telah dijalankan ke atas isolat-isolat IBV dari tahun 1991-2003. Dalam kajian pengasingan, isolat-isolat telah dibahagikan kepada strain Mass dan bukan Mass. Isolat bukan Mass telah digandakan gennya dan dilakukan RFLP untuk mengenalpasti strain nefropatogenik. Sembilan daripada 30 isolat IBV adalah strain Mass, selebihnya dijangkakan nefropatogenik (MH5365/95) kecuali satu isolat. Isolat tersebut adalah V9/03. Isolat ini telah dijujukan dan mempunyai sedikit kesamaan dengan isolat dari Taiwan (AY) dan Korean (AY257060) with 82.5% dan 81.6%. The V9/03 mempunyai persamaan yang rendah (kurang dari 80% dengan nefropatogenik MH5365/95 dan terbitan Mass. Analisis filogenetik menunjukkan V9/03 adalah varian berbeza daripada nefropatogenik MH5365/95 and strain vaksin. Kajian ini menunjukkan bahawa varian IBV berkitaran di ladang, sama seperti mutasi IBV disebabkan penggunaan vaksin virus hidup dalam tempoh yang berpanjangan dan tekanan imunologi virus untuk terus kekal dalam ayam-ayam yang imun.



Acknowledgement

I am grateful to the Almighty Allah for with His blessings, I am able to complete this Molecular Characterization of Infectious Bronchitis Virus thesis. I am forever thankful to my parents, Tuan Haji A Talip Md Noh and Puan Hajjah Zaharah Ya'amat, for without their moral and financial support, I would not be able to complete my Master project.

I would also like to show my highest appreciation to my supervisor, Prof. Madya Dr. Siti Suri Arshad and to all the co-supervisors, Prof. Madya Dr. Abd Rahman Omar and Dr. Sharifah Syed Hassan, who had helped me through the completion of this project.

Many thanks to Mrs. Rodiah Husin, Mr. Kamaruddin Awang Isa and Mrs. Fariza Zainul Abidin who have continuously extended their help in technical works of the research. I would also like to extend my appreciation to Dr. Azri Adzhar, Dr. Faizah Abd Hamid and Mrs. Maizan Haji Mohamed who had given me a lot of ideas and references.

Thank you to Yap May Ling who had guided me throughout the research. To Tan Sheau Wei, Kong Lih Ling and Wan Kei Fei, thank you for your moral support.

Those who have directly and indirectly helped me through this research, Haryati Shila, Dr. Hafiza, Hairul Aini, Thapa, Zulkifli, Tan Do Yew, Kenny, Hazalina, Ernie Zuraida and Zarila, thank you very much.



I am forever grateful to my best friend, Latifah Md Ariffin, for her support, financially and morally, for her patience, ideas and understanding. Not to forget thank you to Khadijah Md Ariffin who helped me a lot in writing this thesis. Lastly, I would like to thank Comel, Cumi and Comot for their presence made this research less stressful and fun.



I certify that an Examination Committee met on the 19th December 2006 to conduct the final examination of Balkis Binti Haji A Talip on her Master of Science thesis entitled “Molecular Characterization of Infectious Bronchitis Virus Isolates from Malaysia” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Dr. Hassan Hj. Mohd Daud, Ph.D.

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Dr. Mohd Hair Bejo, Ph.D.

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Dr. Raha Abdul Rahim, Ph.D.

Associate Profesor
Faculty of Biotechnology and Biomolecular Science
Universiti Putra Malaysia
(Internal Examiner)

Dr. Sazaly Abu Bakar, Ph.D.

Professor
Faculty of Medicine
Universiti of Malaya
(External Examiner)

HASANAH MOHD GHAZALI, Ph.D.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date :



This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirements for the degree of **Master of Science**. The members of the Supervisory Committee were as follows:

Siti Suri Arshad, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Abd Rahman Omar, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Sharifah Syed Hassan, PhD

Director
Veterinary Research Institute
(Member)

AINI IDERIS, PhD

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date :



DECLARATION

I hereby declare that the thesis is based on my original work except for quotation and citations which have been duly acknowledged. I also declare that it has not been previously or is not concurrently, submitted for any other degree at UPM or other institutions.

BALKIS BINTI HAJI A TALIP

Date : -2-2007



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	viii
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF ABBREVIATION	xxi
CHAPTER	
1 INTRODUCTION	1.1
2 LITERATURE REVIEW	
2.1 The aetiology	2.1
2.1.1 Prevalence of Infectious bronchitis disease worldwide	2.1
2.1.2 Prevalence of Infectious bronchitis disease in Malaysia	2.3
2.2 Infectious bronchitis virus	2.4
2.2.1 Classification	2.4
2.2.2 Genome structure	2.4
2.2.3 Gene properties	2.9
2.2.3.1 S1 glycoprotein	2.9
2.2.3.2 sM protein	2.10
2.2.3.3 M protein	2.10
2.2.3.4 N protein	2.12
2.3 Virus replication	2.12
2.4 Antigenic and immunogenic properties of IBV	2.14
2.4.1 Spike protein involves in the induction of protection	2.14
2.4.2 Location of antigenic site	2.16
2.4.3 Sequence variation among IBV isolates	2.17
2.4.3.1 S gene	2.17
2.4.3.2 Gene 3 and M	2.18
2.4.3.3 N gene	2.19
2.4.3.4 Untranslated region (3' UTR)	2.21
2.4.4 Antigenic variation in the IB virus population	2.21
2.5 Pathogenesis of IBV	2.24



2.6	Isolation of IBV (<i>in-vitro</i>)	2.26
2.6.1	Virus isolation using embryonated SPF eggs	2.26
2.6.2	Virus isolation using cell cultures	2.27
2.6.3	Virus isolation using chicken organ cultures	2.27
2.7	Strain classification	2.28
2.7.1	Immunotype or protectotype	2.29
2.7.2	Antigenic types	2.30
2.7.3	Genotypes	2.31
2.8	Methods of differentiating IBV variant	2.32
2.8.1	Reverse transcriptase-polymerase chain reaction (RT-PCR)	2.32
2.8.2	Reverse Transcriptase-Polymerase Chain Reaction-Restriction Fragment Length Polymorphism (RT-PCR-RFLP)	2.37
2.8.3	Enzyme-linked immunosorbent assay (ELISA)	2.38
2.8.4	Virus neutralization (VN)	2.39
2.8.5	Complement fixation (CF) test	2.39
2.8.6	Agar gel precipitin test (AGPT)	2.40
2.8.7	Oligonucleotide fingerprinting	2.40
3	MATERIALS AND METHODS	
3.1	Virus isolates	3.1
3.2	Preparation of tissue samples	3.1
3.3	Propagation of virus stock	3.2
3.4	Primers	3.2
3.5	Small scale preparation of RNA extraction	3.6
3.6	Reverse Transcriptase-Polymerase Chain Reaction	3.6
3.6.1	Two-step RT-PCR	3.6
	3.6.1.1 cDNA synthesis	3.6
	3.6.1.2 Polymerase chain reaction	3.7
3.6.2	One-step RT-PCR	3.8
	3.6.2.1 cDNA synthesis and PCR amplification	3.8
3.7	Nested PCR	3.9
3.8	Restriction enzyme (RE) analysis	3.9
3.9	Agarose gel electrophoresis	3.10
3.10	Purification of DNA fragments	3.10
3.11	Cloning	3.11
3.12	Ligation	3.11
3.13	Transformation of DNA	3.11
3.14	Small scale preparation of DNA plasmid (miniprep)	3.12
3.15	Digestion with restriction enzyme	3.13
3.16	Sequencing of nucleotide	3.13
3.17	Analysis of nucleotides and protein	3.14
4	RESULTS	
4.1	Comparison of two types RT-PCR systems	4.1



	for detection of IBV field isolates	
4.2	Rapid detection of IBV field isolates using universal primer pair UTR1-/UTR2+	4.9
4.3	Differentiation of IBV Massachusetts serotype among IBV local isolates using RT-PCR	4.15
4.4	Characterization of isolates using the S1 region of the S gene using oligonucleotide BD1820R with MHS1E-F and RT-PCR-RFLP	4.21
4.5	Sequence analysis of V9/03 IBV field isolate	4.30
5	DISCUSSION	5.1
6	CONCLUSION AND FUTURE WORK	6.1
	BIBLIOGRAPHY	R.1
	APPENDICES	A.1
	BIODATA OF THE AUTHOR	B.1



LIST OF TABLE

Table		Page
3.1	List of oligonucleotides used for RT-PCR.	3.3
3.2	List of oligonucleotides used for sequencing.	3.5
4.1	Comparison between two-step and one-step RT-PCR in detecting IBV field isolates.	4.7
4.2	List of isolates detected by RT-PCR using universal primer (UTR1-/UTR2+).	4.14
4.3	List of local IBV isolates positive for universal Mass primer and being tested for nested PCR using 2 sets of primer pair.	4.20
4.4	List of IBV isolates negative for Mass strain are for the S1 gene using BD1820R/MHS1E-F.	4.28
4.5	Restriction enzyme analysis of V9/03 compared to local nephropathogenic strain (MH5365/95) and H120 (source : Bioedit).	4.29
4.6	Comparison of amino acid nucleotide identities (%) between S1 sequence of V9/03 and several isolates worldwide.	4.37



LIST OF FIGURES

Figures		Page
2.1	Structure of coronavirus. The viral envelope includes a lipid bilayer derived from intercellular membranes of the host cell and several glycoproteins S and M (Cavanagh <i>et al.</i> , 1994).	2.6
2.2	Molecular organization of coronaviruses (Ackermann, 1995).	2.7
2.3	Genomic organization. The 3' co-terminal nested set of mRNAs is shown. The major structural components of the virion, spike (S), membrane (M) and nucleocapsid (N) protein are shown at the top of the positions for the genes coding. The 5' end of the genomic and subgenomic has a role as leader sequence RNAs. There are conserved sequences i.e. homology regions at the start of each gene (Bournsell <i>et al.</i> , 1985).	2.8
2.4	Antigenic sites in the spike protein of IBV. The numbers represent the differences of amino acid in monoclonal antibody resistant mutants (similar to D207 isolate). The amino acid in location number 63 was derived from strain M41 and most of the others on strains of the D207 group. The shades box indicates the relatedness of VN epitopes and neutralizing antibodies which are sites of A to E shown the association of VN epitopes and sites of F and G are shown the weak of neutralizing antibodies. However, monoclonal antibodies to site D are inhibited hemagglutination. The indicator represented by sp and cp means signal peptide and basic S1 – S2 connecting peptide respectively (Kant <i>et al.</i> , 1992).	2.11
2.5	Coronavirus replication. Coronavirus replication (Sturman and Holmes, 1983). Number of mRNAs and location of nonstructural (NS) proteins may vary, for different coronaviruses virions bind to cell membrane and enter by membrane fusion or endocytosis.	2.15
2.6	The differences of amino acid in S1 subunit of IBV spike protein. The sequences of amino acid residue are depicted by gray boxes, i.e the first amino acid residues from 1 to 130, second amino acid residues from 131 to 260, third amino acid residues from 261 to 391 and fourth amino acid residues from 392 to 520 quarters of S1. However, signal peptides not included. The numbers indicates the differences of amino acid in each quarter for the groups of isolates which are a : among 16 monoclonal antibody resistant mutants IBV strain D207, b : among 13 Mass : like strains, c: among 7 isolates including D207 and UK6/82 with a maximum	2.20



difference of 5% in S1 and d : between UK6/82 and Mass with the total of differences are 20% (Cavanagh, 1995).

- 2.7 Schematic flow digram for reverse transcriptase-polymerase chain reaction (RT-PCR) cycle. In general, reverse transcriptase enzyme will be used coupled with the primer to reverse transcribe RNA to cDNA templates for subsequent PCR. 2.35
- 2.8 Cycle of nested PCR. Nested PCR was performed like standard PCR with modification. 2.36
- 4.1a Agarose gel electrophoresis showing the 300 bp RT-PCR product using two-step procedures obtained with oligonucleotide pair UTR1- and UTR2+, specific for sequence in the 3'UTR. 100 bp DNA ladder was used as marker. Lanes 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14 showed positive product. The lanes 15 and 16 were positive control (Continued). 4.3
- 4.1b Agarose gel electrophoresis showing the 300 bp RT-PCR product using two-step procedures obtained with oligonucleotide pair UTR1- and UTR2+, specific for sequence in the 3'UTR. 100 bp DNA ladder was used as marker. Lanes 19, 20, 21, 22, 23, 24 and 25 showed positive product. The lanes 28 and 29 were positive control. 4.4
- 4.2a Agarose gel electrophoresis of the 300 bp RT-PCR one-step procedures obtained with oligonucleotide pair UTR1- and UTR2+, specific for sequence in the 3'UTR. 100 bp DNA ladder was used as marker. Lanes 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14 and 15 showed positive product. The lane 16 was a positive control (Continued). 4.5
- 4.2b Agarose gel electrophoresis of 300 bp RT-PCR one-step with oligonucleotide pair UTR1- and UTR2+, specific for sequence in the 3'UTR. 100 bp DNA ladder was used as marker. Lanes 19, 20, 22, 24, 25 and 26 showed positive product. The lanes 28 and 29 were positive control. 4.6
- 4.3a Agarose gel electrophoresis of 300 bp selected IBV isolates by using one-step RT-PCR with oligonucleotide pair UTR1-/UTR2+ which is specific for sequence in the 3'UTR. The marker used was 100bp DNA ladder. Lanes 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 and 16 showed positive product (Continued). 4.11
- 4.3b Agarose gel electrophoresis of 300 bp selected IBV isolates by using one-step RT-PCR with oligonucleotide pair UTR1-/UTR2+ which is specific for sequence in the 3'UTR. The marker used 4.12



was 100bp DNA ladder. Lanes 19, 20, 21, 22, 23, 25, 26, 27, 28, 29, 30, 31 and 32 showed positive product. The lane 33 was a positive control (Continued).

- 4.3c Agarose gel electrophoresis of 300 bp selected IBV isolates by using one-step RT-PCR with oligonucleotide pair UTR1-/UTR2+ which is specific for sequence in the 3'UTR. The marker used was 100bp DNA ladder. Lanes 36, 37, 38 and 39 showed positive product. The lane 40 was a positive control. 4.13
- 4.4 Agarose gel electrophoresis of the RT-PCR performed with oligonucleotide pair 640M-/3Az+ to generate approximately 700 bp product derived from gene 3 and gene M sequence. The marker that use for representation size was 100 bp DNA ladder. Lanes 2, 3, 4, 5, 6, 7, 8, 9 and 10 showed positive product. The lanes 11 and 12 were positive control. 4.17
- 4.5 Agarose gel analysis of the nested PCR product revealed the presence of a Massachusetts serotype strain. The nested PCR were performed with Massachusetts-specific oligonucleotide pair 3C+/M- to generate ranging 500-550 bp products. Lanes 2, 3, 4, 5, 6, 7 and 8 showed positive product. The lanes 9 and 10 were positive control. 4.18
- 4.6 Agarose gel analysis of the nested PCR product revealed the presence of a Massachusetts serotype strain. The nested PCR were performed with Massachusetts-specific oligonucleotide pair 3C+/M- to generate 500-550 bp products. Lanes 2, 3, 4, 7 and 8 showed positive product. The lanes 9 and 10 were positive control. 4.19
- 4.7a Agarose gel electrophoresis of the 1.7 kb RT-PCR product obtained with oligonucleotide pair BD1320R and MHSIE-F, specific for sequence in the full-length of S1 and partial of S2 gene sequence. Lanes 2, 3, 4, 5, 6, 7, 8, 9, 10 and 11 showed positive product. The lane 13 was a positive control (Continued). 4.23
- 4.7b Agarose gel electrophoresis of the 1.7 kb RT-PCR product obtained with oligonucleotide pair BD1820R and MHSIE-F, specific for sequence in the full-length of S1 and partial of S2 gene sequence. Lanes 16, 17, 18, 19, 20, 21 and 22 showed positive product (Continued). 4.24
- 4.7c Agarose gel electrophoresis of the 1.7 kb RT-PCR product obtained with oligonucleotide pair BD1820R and MHSIE-F, specific for sequence in the full-length of S1 and partial of S2 gene 4.25



sequence. Lanes 25, 26, 27, 28 and 29 showed positive product.

- 4.8a Restriction enzyme pattern of 1.7 kb PCR product using MboII (Continued). 4.26
- 4.8b Restriction enzyme pattern of 1.7 kb PCR product using MboII. 4.27
- 4.9 Nucleotide sequence of S1 gene obtained from strain V9/03. The DNA consists of four types of nucleotides, adenine (A), thymine (T), cytosine (C) and guanine (G). The sequencing was carried out partially by using walking primer (Table 3.2) and DNA sequencing was performed using an ABI PRISM 3700 DNA Analyzer (Applied Biosystem, USA). 4.35
- 4.10 Amino acid sequences of the S1 gene of isolate V9/03. Alanine (A), Arginine(R), Asparagine (N), Cystine (C), Glutamine (E), Glycine (G), Histadine (H), Isoleucine (I), Leucine (L), Lysine (K), Methionine (M), Phenylalanine (F), Proline (P), Serine (S), Threonine (T), Tryptophan (W), Tyrosine (Y) and Valine (V). The asterisk (*) indicates stop codon. 4.36
- 4.11a The unrooted phylogenetic tree showing the nucleotide sequence relationships of isolate V9/03 to other strains and including Malaysian nephropathogenic strain MH5365/95, JP8443 (AY296745), Shanghai (DQ069317), T-Strain (AY775779), Taiwan (AY606321), Korean (AY257060), Connecticut (L18990), H120 (M21970), M41 (A24863), BEU (Beaudette) (AJ311362), China (DQ07083), IB4/91 (AF093793), UK/93 (UK7/93) (Z83979), Holte (L18988), Gray (L14069), Arkansas (Ark99) (AF094814) and DE0 (DE072) (AF274435). 4.38
- 4.11b The phylogenetic tree showing the nucleotide sequence relationships of isolate V9/03 to other strains and including Malaysian nephropathogenic strain MH5365/95, JP8443 (AY296745), Shanghai (DQ069317), T-Strain (AY775779), Taiwan (AY606321), Korean (AY257060), Connecticut (L18990), H120 (M21970), M41 (A24863), BEU (Beaudette) (AJ311362), China (DQ07083), IB4/91 (AF093793), UK/93 (UK7/93) (Z83979), Holte (L18988), Gray (L14069), Arkansas (Ark99) (AF094814) and DE0 (DE072) (AF274435). 4.39



LIST OF ABBREVIATION

%	percentage
µg	Microgram
µl	microlitre
AF	allantoic fluid
AGPT	agar gel precipitin test
AMV	Avian myeloblastosis virus
ATP	adenosine triphosphate
BEL	Beaudette embryo lethal
bp	base pair
cDNA	copy deoxyribonucleoase acid
CF	complement fixation
cp	connecting peptide
CTP	cytidine triphosphate
ddH ₂ O	distilled deionized water
DNA	deoxyribonuclease acid
dNTP	deoxyribonucleotide triphosphate
DTT	dithiothreitol
<i>E.coli</i>	<i>Escherichia coli</i>
EB	elution buffer
EDTA	ethylenediaminetetraacetic acid
ELISA	enzyme-linked immunosorbent assay
EMBL	England Molecular Biology Laboratory
GTP	guanosine triphosphate
HA	haemagglutination
HCl	hydrochloride acid
HI	haemagglutination-inhibition
IB	infectious bronchitis
IBV	infectious bronchitis virus
kb	kilobase pair
KCl	kalium chloride
kDa	kilodalton
l	litre
LB	Luria Bertani
M	membrane
MAb	monoclonal antibody
MVK	Makmal Veterinar Kawasan Petaling Jaya
MgCl ₂	Magnesium chloride
MHV	murine hepatits virus
mM	miliMolar
Mo-MLV	Moloney murine leukemia virus
MOPS	3- <i>N</i> -morpholino propanesulfonic acid
mRNA	messenger RNA
MVP	Malaysian Vaccine Pharmaceutical
N	Nucleocapsid



NaCl	sodium chloride
NDV	New Castle disease virus
NS	Nonstructural
ORF	open reading frame
PBS	phosphate buffer saline
PCR	polymerase chain reaction
pmol	picomole
RE	restriction enzyme
RNA	ribonucleic acid
RT-PCR	reverse transcriptase-polymerase chain reaction
RT-PCR-RFLP	reverse transcriptase-polymerase chain reaction-restriction fragment length polymorphism
S	spike
SDS	Sodium Dodecyl Sulfate
SgRNAs	subgenomic mRNA
sp	signal peptide
TBE	Tris-borate EDTA buffer
TTP	thymidine triphosphate
U	unit
UK	United Kingdom
UPM	Universiti Putra Malaysia
USA	United States of America
UTR	untranslated region
UV	ultraviolet
V/60	volt per 60 minutes
VN	virus neutralization
VNT	virus neutralization test
VRI	Veterinary Research Institute
w/v	weight per volume



CHAPTER 1

GENERAL INTRODUCTION

Infectious bronchitis virus (IB) is an acute, highly contagious respiratory and urogenital disease of chickens. The disease, also called avian infectious bronchitis virus is of significant to the economic importance to the poultry industry as it causes poor weight gain and feed efficiency. The disease was first reported in 1931 in the United State of America (Beach and Schalm, 1936). It is one of the organisms isolated and produces airsacculitis resulting in condemnations of broilers at processing (King and Cavanagh, 1991).

IBV together with Turkey and pheasant coronaviruses, belongs to Group 3 of the genus *Coronavirus*, family *Coronaviridae* (Cavanagh, 2001). All the members of this genus have a linear, non-segmented, positive sense, single stranded RNA genome of approximately 27 kb in length. The first 20 kb encode the viral RNA-dependent RNA polymerase and proteases. The remainder of the genome encodes five structural proteins, the spike (S) consisting S1 and S2, envelope (E), membrane (M) and nucleocapsid (N) proteins, four small non-structural proteins 3a, 3b, 5a and 5b and a 3' untranslated region (UTR).

Variations in the S1 and N genes, in particular, are believed to be critical for emergence of variants because of their role in virus replication and immunity. Moreover, S1 and N



have been used most frequently to determine the relatedness of emerging IBV. The S1 glycoprotein is located on the surface of the virion and carries epitopes and determinants for virus neutralizing antibodies, protective immunity and cell tropism (Casais *et al.*, 2003 ; Cavanagh *et al.*, 1986 ; Cavanagh *et al.*, 1988). The N protein, located in the capsid of the virion is involved in RNA replication and carries group-specific antigenic determinants (Ignjatovic and Galli, 1995). The 3'UTR region is involved in initiation of negative-strand RNA synthesis and has also been used to assess variation in emerging IB strains and other members of coronavirus Group 3 (William *et al.*, 1993).

IBV evolution is currently considered to be driven by three factors such as the inherent propensity of its RNA genome to mutate, the continuous use of live, often multiple vaccines and the immunological pressure exerted on circulating viruses by the enduring presence of immune bird populations. These three factors probably play a role in performance and involve various mechanisms, such as point mutations, deletion, insertions and recombination to generate new variants (Cavanagh *et al.*, 1992a ; Kuster *et al.*, 1990 ; Wang *et al.*, 1994).

Serotyping of IBV isolate is essential in the design of prevention and control strategies as well as in epidemiological studies. Several serotyping methods have been used including virus neutralization (VN) (Zwaagstra *et al.*, 1992), reverse transcription-polymerase chain reaction-restriction fragment length polymorphism (RT-RFLP) (Kwon *et al.*, 1993), serotype-specific reverse transcriptase-polymerase chain reaction (RT-PCR) (Lai and Cavanagh, 1997) and multiplex PCR (Wang and Khan, 1999). For the control of disease

