



UNIVERSITI PUTRA MALAYSIA

**WOOD QUALITY OF TEN-YEAR-OLD SENTANG (*Azadirachta
excelsa*) GROWN FROM SEEDLINGS AND ROOTED CUTTINGS**

NORDAHLIA BINTI ABDULLAH SIAM

T FH 2009 1



**WOOD QUALITY OF TEN-YEAR-OLD SENTANG (*Azadirachta excelsa*)
GROWN FROM SEEDLINGS AND ROOTED CUTTINGS**

By

NORDAHLIA BINTI ABDULLAH SIAM

**Thesis Submitted to the School of Graduates Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirement for the Degree of Master of Science**

January 2009



Specially dedicated to:

**My beloved father Abdullah Siam b. Alias and mother
Hamidah bt. A.R. Hassan**

Your love always in my heart

My beloved husband Khairul Iman b. Khairuddin

Your love has kept me going on

**My beloved father and mother in law Khairuddin b.
Kamaruddin and Mariam bt. Din**

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Master of Science

**WOOD QUALITY OF TEN-YEAR-OLD SENTANG (*Azadirachta excelsa*)
GROWN FROM SEEDLINGS AND ROOTED CUTTINGS**

By

NORDAHLIA BINTI ABDULLAH SIAM

2009

Chairman: Associate Professor Zaidon Ashaari, PhD

Faculty: Forestry

Seeds of sentang were exhibit recalcitrant and cannot be stored for a long period of time. So efforts have been made to propagate this species vegetatively via rooted cuttings technique. It is envisaged that for the technique to be deemed successful, the quality of wood that it produces should be comparable with or better than the wood extracted from the natural forest or established plantation via seedlings. The main objective of this study was to determine and compare the wood qualities of the trees grown from seedlings and rooted cuttings and to study the property variations within the tree. The specific objectives of these studies were to evaluate the macroscopic features, anatomical, physical, mechanical properties and resistance to fungal decay of sentang which grown from these two propagation techniques. Correlations between anatomical, physical and mechanical properties of the wood were also analysed. The studies of both types of wood were done at three different height levels, i.e bottom, middle and top and also in the sapwood and heartwood



portion. Data were subjected to Analysis of Variance (ANOVA) and further analysed using Least Significant Difference (LSD) at $p \leq 0.05$.

The barks of both types of wood were smooth and pinkish grey. There was significant difference in colour of sapwood and heartwood between these two propagation techniques where the sapwood from seedling trees was lighter than those from rooted-cutting trees. However, the heartwood of the latter was darker. The mean percentage of sapwood-heartwood ratio was not significantly different between these two types of wood and the amount of sapwood-heartwood ratio increased towards the top. The anatomical structure in both types of wood is similar with typical sentang. Vessels are diffuse, solitary or multiple, round to oval shaped and filled with dark-coloured dried extractives deposits in the heartwood. Apotracheal and paratracheal parenchyma are present in both types of wood. Tangential section shows multiseriate rays which mostly 2-3 cells wide and heterocellular rays were observed in the radial section in both types of wood. The rooted-cutting wood showed higher value in fibre dimensions, vessel diameter and proportion of vessels, whereas vessel density and proportion of fibre were higher in the seedling wood and proportion of rays was not significantly different between both types of wood. Both the seedling and rooted-cutting wood showed a decreasing trend towards the top in fibre dimensions, vessel diameter and proportion of rays. Increasing trend was observed in vessel density and proportions of vessels and fibres. The sapwood exhibited higher value in fibre dimensions, proportion of rays and larger vessel diameter but lower value in vessel density and proportions of vessels and fibres than the heartwood in the seedling wood.

However, in the rooted-cutting wood there was no significant difference between sapwood and heartwood in proportion of rays, vessels and fibres.

Both types of wood were not significantly different in basic density. Moisture content at green condition and shrinkages in green to air-dry and green to oven-dry conditions were higher in the rooted-cutting wood, but tangential and radial shrinkages from green to oven-dry conditions showed no significant difference between both types of wood. The physical properties of both types of wood showed a decreasing trend from the bottom to the middle of the tree and then a slight increasing trend towards the top. Basic density was higher in the sapwood but moisture content and shrinkages in both conditions were higher in the heartwood.

Compression and shear parallel to the grain were higher in the seedling wood, however MOE was higher in the rooted-cutting wood and MOR showed no significant difference between the seedling and rooted-cutting wood. Mechanical properties showed increasing trend towards the top and higher values in the heartwood in both types of wood. All the mechanical properties in the seedling and rooted-cutting wood were significantly related with vessel diameter, proportion of fibres and volumetric shrinkage with small correlation. Basic density was weakly correlated with mechanical properties in this present study. Both types of wood were classified as resistant with rooted-cutting wood being more resistance to white rot fungus (*Lentinus sajor-caju*) than the seedling wood. The resistance of the wood to fungal decay decreased towards the top and higher resistance was shown in the heartwood than the sapwood portion.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**KUALITI KAYU SENTANG (*Azadirachta excelsa*) BERUMUR SEPULUH
TAHUN YANG DITANAM SECARA BIJI BENIH DAN KERATAN
PENGAKAR**

Oleh

NORDAHLIA BINTI ABDULLAH SIAM

2009

Pengerusi: Profesor Madya Zaidon Ashaari, PhD

Fakulti: Perhutanan

Biji benih sentang adalah rekalsitran dan tidak boleh disimpan dalam jangka masa yang lama. Oleh itu, usaha telah diambil untuk menanam spesies ini secara vegetatif menggunakan teknik keratan pengakar. Teknik ini dijangka akan berjaya di mana kualiti kayu yang dihasilkan seharusnya sama atau lebih baik daripada kayu yang diambil dari hutan asli atau dari perladangan yang ditanam secara biji benih. Objektif utama kajian ini adalah untuk menentukan dan membandingkan kualiti kayu pada pokok yang ditanam secara biji benih dan keratan pengakar dan untuk melihat variasi sifat kayu pada pokok tersebut. Objektif spesifik penyelidikan adalah untuk mengenalpasti sifat makroskopik, anatomi, fizikal, mekanikal dan ketahanan kayu terhadap kulat pereput bagi pokok sentang dari dua teknik penanaman ini. Kajian terhadap korelasi di antara sifat-sifat anatomi, fizikal dan mekanikal juga dijalankan. Kajian bagi kedua-dua jenis kayu dijalankan pada tiga paras ketinggian yang berbeza iaitu bahagian bawah, tengah dan atas serta pada gubal dan teras kayu. Data dianalisis



menggunakan analisis varian (ANOVA) dan seterusnya dianalisis menggunakan perbezaan minimum yang ketara (LSD) pada $p \leq 0.05$.

Kulit pokok bagi kedua-dua jenis kayu adalah licin dan berwarna kelabu kemerah muda. Terdapat perbezaan bererti bagi warna pada kayu gubal dan kayu teras bagi kedua-dua teknik penanaman ini. Kayu gubal pada kayu anak benih adalah lebih cerah berbanding kayu keratan pengakar. Manakala, kayu teras lebih gelap pada kayu keratan pengakar. Purata peratus kayu gubal tiada perbezaan bererti antara kedua-dua teknik penanaman ini dan meningkat ke bahagian atas pokok. Struktur anatomi pada kedua-dua jenis kayu adalah menyamai dengan sentang tipikal. Sel salur adalah tunggal, beganda, berbentuk bulat atau oval, dipenuhi dengan ekstraktif pada bahagian kayu teras. Terdapat parenkima apotrakeal dan paratrakeal pada kedua-dua jenis kayu. Bahagian tangen menunjukkan ruji multiseriat yang lazimnya mengandungi 2-3 sel dan ruji heterosel didapati di bahagian radial pada kedua-dua jenis kayu. Kayu keratan pengakar menunjukkan nilai yang tinggi dalam sifat gentian, diameter sel salur dan peratusan sel salur, manakala ketumpatan sel salur dan peratusan gentian adalah tinggi bagi kayu biji benih dan peratusan ruji menunjukkan nilai yang tidak bererti diantara kedua-dua teknik ini. Keputusan dari sifat anatomi didapati kayu biji benih dan kayu keratan pengakar menunjukkan penurunan ke bahagian atas bagi sifat gentian, diameter sel salur dan peratusan ruji, manakala ketumpatan sel salur, peratusan sel salur dan gentian meningkat ke bahagian atas. Sifat gentian, diameter sel salur, peratusan ruji adalah tinggi di bahagian kayu gubal, manakala ketumpatan sel salur, peratusan sel salur dan gentian adalah tinggi di bahagian kayu teras pada kayu biji benih. Manakala, pada kayu keratan pengakar,

tiada perbezaan bererti antara bahagian kayu gubal dan kayu teras bagi peratusan ruji, sel salur dan gentian.

Ketumpatan asas menunjukkan nilai tidak bererti di antara kedua-dua jenis kayu manakala kandungan lembapan pada keadaan basah dan pengecutan bagi keadaan basah ke kering udara dan basah ke kering oven adalah tinggi pada kayu keratan pengakar, tetapi pengecutan tangen dan pengecutan radial dari keadaan basah ke kering oven menunjukkan nilai tidak bererti antara kedua-dua teknik penanaman. Kedua-dua jenis kayu menunjukkan penurunan dari bahagian bawah ke bahagian tengah kemudian meningkat sedikit ke bahagian atas bagi sifat fizikal. Ketumpatan asas tinggi di bahagian kayu gubal, manakala kandungan lembapan dan pengecutan bagi kedua-dua keadaan tinggi di bahagian kayu teras.

Mampatan dan kekuatan ricihan selari ira tinggi pada kayu biji benih, tetapi MOE adalah tinggi pada kayu keratan pengakar dan MOR menunjukkan nilai tidak bererti antara kayu biji benih dan kayu keratan pengakar. Sifat mekanikal menunjukkan peningkatan ke bahagian atas dan nilai yang tinggi di bahagian kayu teras bagi kedua-dua jenis teknik penanaman. Semua sifat mekanikal pada kayu biji benih dan kayu keratan pengakar adalah berkolerasi secara bererti dengan diameter sel salur, peratusan gentian dan pengecutan volumetrik. Ketumpatan adalah berkolerasi sangat rendah dengan sifat mekanikal pada kajian ini. Kedua-dua kayu dikelaskan sebagai rintang dengan kayu dari keratan pengakar lebih rintang terhadap kulat pereput putih. Ketahanan kedua-dua kayu menurun ke bahagian atas dan kerintangannya adalah tinggi di bahagian kayu teras berbanding kayu gubal.

ACKNOWLEDGEMENTS

In the Name of Allah, The Most Merciful and The Most Gracious. Praise goes to ALLAH Almighty for I have been blessed with the strength and ardour to finally accomplish this thesis.

First of all, I would like to express my sincere gratitude and appreciation to Associate Professor Dr. Zaidon Ashaari, Chairman of the Supervisory Committee, for his constant guidance, assistance and suggestions throughout the preparation of this thesis. Heartiest thanks are also due to Dr. Ani Sulaiman for her constant support, encouragement and invaluable advice. I would also like to express my profound gratitude to Prof. Dr Mohd Hamami Sahri for his invaluable suggestions and constructive comments.

I am indebted to the Director General of the Forest Research Institute Malaysia (FRIM) Dato' Dr. Abd. Latif Mohmood for the scholarship grants (RA Scheme). Sincere appreciation is also extended to Dr. Mohamad Omar Mohamad Khaidzir, Dr. Salmiah Ujang, Dr. Ahmad Zuhaidi Yahya, Dr. Mohd Khairun Anwar Uyup and Dr Hamdan Husain for their cooperation and constant support in this project. Cooperation from other FRIM staff Mr. Mohd Zahari Mohd Noor, Mrs Norjihan Abu Bakar, Mrs Rohana Idris, Mr Lim S.C, Mr. Sabri Ahmad, Mr Mohd Jamil Abd. Wahab, Mrs Chong Y.T, Mr. Roszalli Mohd, Mr. Abd Karim Nasir, Mr. Abd Ghani Mohamad, Mr. Rosdi Koter, Mr. Kamarul Hizam Hamsan, Mr. Mahat Judin, Mr. Zakaria Yusoff and others is also greatly appreciated.



To my father Mr. Abdullah Siam Alias, my Mother Mrs. Hamidah A. R. Hassan, my grandmother (Mrs Hjh. Sujirah Tasim) my brothers (Mohd Azrai'i and Muhammad A'mmar) and my sister (Nor Dalila Sa'adah), I render my gratitude and thanks for their advices, patience, concern, support and sacrifices. My appreciations also extended to my father and mother in law (Mr. Khairuddin Kamaruddin and Mrs. Mariam Din) and brothers (Khairil Nizam and Muhd Nur Izham) for their encouragement and support. Special thank to my beloved husband Mr. Khairul Iman Khairuddin, who are always concern with my study and have given me support to improve myself, your love has kept me going on.



I certify that an Examination Committee has met on **9th January 2009** to conduct the final examination of Nordahlia binti Abdullah Siam on her Master of Science thesis entitled “Wood Quality of 10-year-old Sentang (*Azadirachta excelsa*) Grown from Seedlings and Rooted Cuttings” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The committee recommends that the student be awarded the Master of Science.

Members of the Examination Committee were as follows:

H’ng Paik San, PhD

Faculty of Forestry
Universiti Putra Malaysia
(Chairman)

Edi Suhaimi Bakar, PhD

Associate Professor
Faculty of Forestry
Universiti Putra Malaysia
(Internal Examiner)

Faizah Abood, PhD

Associate Professor
Faculty of Forestry
Universiti Putra Malaysia
(Internal Examiner)

Razak Wahab, PhD

Associate Professor
School of International Tropical Forestry
Universiti Malaysia Sabah
(External Examiner)

HASANAH MOHD. GHAZALI, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Zaidon Ashaari, PhD

Associate Professor
Faculty of Forestry
Universiti Putra Malaysia
(Chairman)

Mohd Hamami Sahri, PhD

Professor
Faculty of Forestry
Universiti Putra Malaysia
(Member)

Ani Sulaiman, PhD

Senior Research Officer of Wood Anatomy
Forest Research Institute of Malaysia
(Member)

HASANAH MOHD. GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 9 April 2009



DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledge. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

NORDAHLIA BINTI ABDULLAH SIAM

Date: 16 March 2009



TABLE OF CONTENTS

DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL SHEETS	xi
DECLARATION FORM	xiii
TABLE OF CONTENTS	xiv
LIST OF TABLES	xvii
LIST OF FIGURES	xx
LIST OF PLATES	xxii
LIST OF ABBREVIATIONS	xxiv
CHAPTER	
1 INTRODUCTION	1
1.1 Background	1
1.2 Problem Statement	4
2 LITERATURE REVIEW	6
2.1 Forest Plantation	6
2.1.1 The Importance of Forest Plantation	6
2.1.2 Selected Plantation Species	7
2.2 Background of Sentang	8
2.2.1 Botanical Name	9
2.2.2 Vernacular Name	10
2.2.3 Description of the Tree	10
2.2.4 Anatomical Properties of Sentang	13
2.2.5 Physical Properties of Sentang	15
2.2.6 Mechanical Properties of Sentang	16
2.2.7 Natural Durability of Sentang	18
2.3 Distribution of Sentang	19
2.4 Environmental Consideration of Sentang	20
2.5 Uses of Sentang	21
2.6 Pests and Diseases of Sentang	23
2.7 Propagation of Sentang	24
2.7.1 Seedlings	24
2.7.2 Vegetative Propagation	25
2.8 Properties of Wood from Different Propagation Technique in Different Species	28
2.9 The Concept of Wood Quality	30
2.9.1 Density	30
2.9.2 Uniformity and Ring Width	31
2.9.3 Fibre Length	32
2.9.4 Spiral Grain	32
2.9.5 Proportion of Heartwood	33

2.9.6 Cellular Composition	33
2.9.7 Knots	34
2.9.8 Chemical Composition	34
2.9.9 Brightness of Wood	35
2.9.10 Occurrence of Juvenile Wood and Reaction Wood	35
3 COLLECTION OF SAMPLES	37
3.1 Field Sampling	37
3.2 Preparation of Samples	39
4 MACROSCOPIC FEATURES AND ANATOMICAL PROPERTIES OF 10-YEAR-OLD SENTANG GROWN FROM SEEDLINGS AND ROOTED CUTTINGS	41
4.1 Introduction	41
4.2 Materials and Methods	43
4.2.1 Macroscopic Evaluation	43
4.2.2 Evaluation of Anatomical Properties	44
4.2.3 Maceration	48
4.2.4 Light Microscopy	49
4.2.5 Statistical Analysis	50
4.3 Results and Discussion	51
4.3.1 Macroscopic Features of 10-year-old Sentang Grown from Seedlings and Rooted Cuttings	51
4.3.2 Wood Anatomical Structure of 10-year-old Sentang Grown from Seedlings and Rooted Cuttings	55
4.3.3 Anatomical Properties of 10-year-old Sentang Grown from Seedlings and Rooted Cuttings	58
4.3.4 Variations of Anatomical Properties of 10-year-old Sentang Grown from Seedlings and Rooted Cuttings	61
4.3.5 Correlation Coefficients among Anatomical Properties	73
4.4 Conclusion	75
5 PHYSICAL PROPERTIES OF 10-YEAR-OLD SENTANG GROWN FROM SEEDLINGS AND ROOTED CUTTINGS	77
5.1 Introduction	77
5.2 Materials and Methods	79
5.2.1 Sample Preparation	79
5.2.2 Moisture Content Determination	79
5.2.3 Basic Density Determination	80
5.2.4 Shrinkage Determination	80
5.2.5 Statistical Analysis	81
5.3 Results and Discussion	82
5.3.1 Physical Properties of 10-year-old Sentang Grown from Seedlings and Rooted Cuttings	82
5.3.2 Variations of Physical Properties of 10-year-old Sentang Grown from Seedlings and Rooted Cuttings	89

5.3.3 Correlation Coefficients between Anatomical and Physical Properties	100
5.4 Conclusion	107
6 MECHANICAL PROPERTIES OF 10-YEAR-OLD SENTANG GROWN FROM SEEDLINGS AND ROOTED CUTTINGS	109
6.1 Introduction	109
6.2 Materials and Methods	111
6.2.1 Sample Preparation	111
6.2.2 Static Bending	113
6.2.3 Compression Parallel to the Grain	114
6.2.4 Shear Parallel to the Grain	115
6.2.5 Statistical Analysis	116
6.3 Results and Discussion	117
6.3.1 Mechanical Properties of 10-year-old Sentang Grown from Seedlings and Rooted Cuttings	117
6.3.2 Variations of Mechanical Properties of 10-year-old Sentang Grown from Seedlings and Rooted Cuttings	119
6.3.3 Correlation Coefficients between Anatomical Properties, Basic Density and Volumetric Shrinkage and Mechanical Properties	127
6.4 Conclusion	130
7 RESISTANCE OF 10-YEAR-OLD SENTANG GROWN FROM SEEDLINGS AND ROOTED CUTTINGS TO FUNGAL DECAY	132
7.1 Introduction	132
7.2 Materials and Methods	133
7.2.1 Sample Preparation	133
7.2.2 Decay Study Using White Rot Fungus	134
7.2.3 Statistical Analysis	136
7.3 Result and Discussion	136
7.3.1 Resistance of 10-year-old Sentang Grown from Seedlings and Rooted Cuttings to Fungal decay	136
7.3.2 Variations of Resistance to Fungal Decay of 10-year-old Sentang Grown from Seedlings and Rooted Cuttings	138
7.4 Conclusion	142
8 GENERAL CONCLUSIONS AND RECOMMENDATIONS	143
8.1 Conclusions	143
8.2 Recommendations	146
REFERENCES	148
BIODATA OF STUDENT	165

Table	LIST OF TABLES	Page
2.1	Ranges in girth of sentang in Southern Thailand	9
2.2	Mechanical properties of sentang	18
3.1	Measurements of the 10-year-old sentang trees grown from seedlings and rooted cuttings	38
4.1	Whitish values of the sapwood and heartwood of the 10-year-old sentang grown from seedlings and rooted cuttings	53
4.2	Percentages of sapwood within the stem height and averages of the sapwood percentage of 10-year-old sentang grown from seedlings and rooted cuttings	54
4.3	Anatomical properties of the 10-year-old sentang grown from seedlings and rooted cuttings	58
4.4	Anatomical properties of the 10-year-old sentang grown from seedlings and rooted cuttings compared with those of study on sentang and other fast-growing species from other studies	60
4.5	Analysis of variance of anatomical properties of the 10-year-old sentang grown from seedlings and rooted cuttings	62
4.6	Variations along the tree height of fibre dimensions of the 10-year-old sentang grown from seedlings and rooted cuttings	64
4.7	Variations along the tree height of proportions of rays, vessels and fibres of the 10-year-old sentang grown from seedlings and rooted cuttings	67
4.8	Variations of fibre dimensions in the sapwood and heartwood of the 10-year-old sentang grown from seedlings and rooted cuttings	69
4.9	Variations of proportions of rays, vessels and fibres in the sapwood and heartwood of the 10-year-old sentang grown from seedlings and rooted cuttings	72
4.10	Correlations among the anatomical properties of the 10-year-old sentang grown from seedlings and rooted cuttings	74
5.1	Physical properties of the 10-year-old sentang grown from seedlings and rooted cuttings	84



5.2	Comparison of the physical properties between wood from the 10-year-old sentang grown from seedlings and rooted cuttings and that from mature sentang and other fast-growing species	88
5.3	Summary of analysis of variance of the physical properties of the 10-year-old sentang grown from seedlings and rooted cuttings	90
5.4	Shrinkage variations along the tree height of the 10-year-old sentang grown from seedlings and rooted cuttings	95
5.5	Variations of shrinkage properties in the sapwood and heartwood of the 10- year-old sentang grown from seedlings and rooted cuttings	100
5.6	Correlations between anatomical and physical properties of the 10-year-old sentang grown from seedlings and rooted cuttings	106
6.1	Mechanical properties of the 10-year-old sentang grown from seedlings and rooted cuttings	117
6.2	Mechanical properties of the 10-year-old sentang grown from seedlings and rooted cuttings as compared with those from previous studies on sentang and other commercial species	119
6.3	Summary of analysis of variance of mechanical properties of the 10-year-old sentang grown from seedlings and rooted cuttings	120
6.4	Variations of mechanical properties along the tree height of the 10-year-old sentang grown from seedlings and rooted cuttings	122
6.5	Variations of mechanical properties in the sapwood and heartwood of the 10-year-old sentang grown from seedlings and rooted cuttings	124
6.6	Correlations between mechanical properties and anatomical properties, basic density and volumetric shrinkage of the 10-year-old sentang grown from seedlings and rooted cuttings	129
7.1	Decay resistance of the 10-year old sentang grown from seedlings and rooted cuttings	137



7.2	Analysis of variance of the 10-year-old sentang grown from seedlings and rooted cuttings	139
7.3	Mean percentages of weight loss in the sapwood and heartwood of the 10-year-old sentang grown from seedlings and rooted cuttings	144



Figure	LIST OF FIGURES	Page
3.1	Maps of sampling field at Bukit Hari, FRIM	37
3.2	Selection of sentang trees using completely randomised design at Bukit Hari, Forest Research Institute Malaysia, for wood quality assessment	39
3.3	Locations of sampling points within a selected tree	40
4.1	Axial lines drawn across the disc surface for measurements of disc and heartwood diameters	44
4.2	Sampling method for the preparation of samples for anatomical studies	45
4.3	Flow chart of the process of staining and dehydration of thin wood sections	47
4.4	Arrangement of wood sections	48
4.5	The sampling fields for anatomical studies	50
4.6	Vessel diameters of the 10-year-old sentang grown from seedlings and rooted cuttings	65
4.7	Vessel densities of the 10-year-old sentang grown from seedlings and rooted cuttings	66
4.8	Vessel diameter variation in the sapwood and heartwood of the 10-year-old sentang grown from seedlings and rooted cuttings	70
4.9	Vessel density variations in sapwood and heartwood of the 10-year-old sentang grown from seedlings and rooted cuttings	71
5.1	Variations of green moisture content along the tree height of the 10-year-old sentang grown from seedlings and rooted cuttings	91
5.2	Variations of basic density along the tree height of the 10-year-old sentang grown from seedlings	92
5.3	Moisture contents in the sapwood and heartwood of the 10-year-old sentang grown from seedlings and rooted cuttings	96
5.4	Basic densities of the 10-year-old sentang grown from	98

	seedlings and rooted cuttings in the sapwood and heartwood	
6.1	Sampling method for mechanical properties study	112
7.1	Variations of wood resistance along the tree height of the 10-year-old sentang grown from seedlings and rooted cuttings	140
7.2	Resistance to fungal decay in the sapwood and heartwood of the 10-year-old sentang grown from seedlings and rooted cuttings	142



Plate	LIST OF PLATES	Page
2.1	Pseudo-tap roots in 6-year-old jelutong propagated by tissue culture	28
3.1	10-year-old tap root system of seedlings sentang, Plot 48, Bukit Hari, (FRIM)	38
3.2	10-year-old pseudo-tap root system of rooted cuttings sentang, Plot 44, Bukit Hari, (FRIM)	38
4.1	Barks of the 10-year-old sentang grown from (a) seedlings and (b) rooted cuttings	51
4.2	Discs of the 10-year-old sentang grown from (a) seedlings and (b) rooted cuttings	52
4.3	Cross section from the heartwood of the 10-year-old wood cut from the seedling sentang which was similar in anatomical structure to the wood cut from the rooted-cutting sentang and typical wood of sentang (magnification 50X)	56
4.4	Cross section from the sapwood of the 10-year-old wood cut from the seedling sentang which was similar to the anatomical structure with wood cut from the rooted-cutting and typical wood of sentang (magnification 50X)	56
4.5	Tangential section from the heartwood of the 10-year-old wood cut from the rooted cutting sentang which was similar in anatomical structure to the wood cut from the seedling sentang and typical wood of sentang (magnification 50X)	57
4.6	Radial section from the sapwood of the 10-year-old wood cut from the rooted-cutting sentang which was similar in anatomical structure to the wood cut from the seedling sentang and typical wood of sentang (magnification 50X)	57
6.1	Bending test (MOR and MOE)	114
6.2	Compression parallel to the grain test	115
6.3	Shear parallel to the grain test	116
6.4	Compression at the top and splintering tension at the bottom were observed in both types of wood of sentang	125
6.5	Crushing at the top was observed in both types of wood of sentang	126



6.6	Crushing at the bottom was observed in both types of wood of sentang	126
7.1	Samples of the 10-year-old sentang wood grown from seedlings and rooted cuttings after 12 weeks exposure to the test fungus	135

LIST OF ABBREVIATIONS

ANOVA	Analysis of Variance
ASTM	American Standard Testing Method
BS	British Standard
CFFPR	Compensatory Forest Plantation Project
CWT	Fibre Wall Thickness
DBH	Diameter Breast Height
F (%)	Fibre proportion
FD	Fibre diameter
FL	Fibre Length
FRIM	Forest Research Institute Malaysia
HW	Heartwood
L	Longitudinal Shrinkage
LD	Fibre Lumen Diameter
LSD	Least Significant Difference
LVL	Laminated Veneer Lumber
MC	Moisture Content
MOE	Modulus of Elasticity
MOR	Modulus of rupture
MTC	Malaysian Timber Council
MTIB	Malaysia Timber Industry Board
R	Radial Shrinkage
R (%)	Ray proportion
SAS	Statistical Analysis Software