



**UNIVERSITI PUTRA MALAYSIA**

**INFLUENCE OF SOIL EXCHANGEABLE CATIONS ON GROWTH,  
NUTRIENT UPTAKE AND PHYSIOLOGY OF OIL PALM SEEDLINGS**

**AHMAD AFANDI BIN MURDI**

**FP 2007 12**



**INFLUENCE OF SOIL EXCHANGEABLE CATIONS ON GROWTH,  
NUTRIENT UPTAKE AND PHYSIOLOGY OF OIL PALM SEEDLINGS**

**By**

**AHMAD AFANDI BIN MURDI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the degree of Master of Agricultural Science**

**June 2007**



Dedicated to Suhaili, Amirul Haziq  
Nur Hanisa and Nur Hasya



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Agricultural Science

**INFLUENCE OF SOIL EXCHANGEABLE CATIONS ON GROWTH,  
NUTRIENT UPTAKE AND PHYSIOLOGY OF OIL PALM SEEDLINGS**

By

**AHMAD AFANDI BIN MURDI**

**June 2007**

**Chairman : Associate Professor Ahmad Husni Mohd Hanif, PhD**

**Faculty : Agriculture**

Increasing the efficiency of nutrient uptake can significantly reduce the cost of production, since fertilizer cost is about 30% of total cost of oil palm production. Despite many agronomic trials on the response of oil palm to various fertilizers, there were few published information on the multidisciplinary study of physiological responses with cations e.g. potassium (K), calcium (Ca) and magnesium (Mg). Therefore, this study is necessary to increase efficiency in the oil palm nutrient management system through multidisciplinary aspects of soil and plant physiology. The trial was conducted in the nursery of the Malaysian Palm Oil Board (MPOB) Research Station in Lahad Datu, Sabah. Oil palm seedlings were grown in polybag filled with two types of soil i.e Semporna Family which contain high exchangeable Ca or Lumpongan Family with low exchangeable Ca. Semporna soil was tested at  $3^2$  levels of K and Mg and Lumpongan was tested at  $3^3$  levels of K, Ca and Mg with



three replicates. Among the parameters studied were vegetative measurements, relative chlorophyll content and leaf relative water content. Destructive samplings of whole plants were done and seedlings were separated into rachis, pinnae, stem and roots. The tissues were then used to determine K, Mg and Ca concentrations and uptake. Soil samples were taken at the beginning and the end of trials and subsequently analysed for cations content. Leaf gas exchange parameters were measured using a portable infrared gas analyser two weeks before the final destructive sampling. Pre-treatment analysis showed that the exchangeable K, Mg and Ca in Lumpongan soil were 0.35, 6.74 and 6.07 cmol (+) kg<sup>-1</sup> respectively. Meanwhile, in Semporna soil the exchangeable K, Mg and Ca were 0.17, 1.22 and 33.50 cmol (+) kg<sup>-1</sup>. Application of cation fertilizers significantly increased respective exchangeable cations in both soils. There were non significant increases in seedling dry matter components with regards of cations treatments in both soils. The concentration and uptake of K in seedling components were increased significantly by K fertilizer in both soils. Magnesium fertilizer was not recommended on Lumpongan soil but 6.69 g of Mg per seedling are needed to sustain the growth of seedling in Semporna soil. The seedling recovery efficiency for K and Ca in Lumpongan soil was about 3.65 and 1.21%, respectively. Meanwhile, in Semporna soil the recovery efficiency for K and Mg was about 4.56 and 2.54%, respectively. This study showed that total cation in oil palm seedling leaves was largely determined by soil exchangeable calcium rather than K or Mg. The proportion of individual tissue cation to total cations was fairly constant, i.e about 29% if the soil exchangeable calcium is high (> 25 cmol (+) kg<sup>-1</sup>). This was considered sufficiently balance for oil palm nutrient requirement. However, excessive amount of

exchangeable Mg in soil ( $> 4.75 \text{ cmol (+) kg}^{-1}$ ) such as Lumpongan soil resulted in an imbalance proportion of individual tissue cation to total cation. The oil palm seedling vegetative growth, leaf tissues turgidity and chlorophyll content exhibited less sensitivity to cation treatments as indicated by non significant difference in both soils. Excessive amount of Ca in Semporna soil contributed to low shoot to root ratio of the seedlings by 7 to 9% as compared to Lumpongan soil. The photosynthesis rate on Lumpongan and Semporna soils ranged from  $8.52$  to  $9.45 \mu\text{mol m}^{-2}\text{s}^{-1}$  and  $7.07$  to  $8.66 \mu\text{mol m}^{-2}\text{s}^{-1}$ , respectively. This implies that, cations treatments did not significantly reduce the photosynthesis rate. It was also concluded that the stomata conductance and intercellular  $\text{CO}_2$  concentrations of seedlings in both soils were adequate to support leaf gas exchange as indicated by non significant different between treatments. In conclusion, combinations of K1 (35.11 g K) + Mg0 (0 g Mg) + Ca1 (14 g Ca) and K2 (70.22 g K) + Mg1 (6.69 g Mg) were recommended for sustaining optimum growth, nutrient uptake and physiological requirement in oil palm seedlings grown in Lumpongan and Semporna soils, respectively. These combinations take into account the total dry matter production, uptake, recovery efficiency and physiological characteristics of oil palm seedlings at each cation levels.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains Pertanian

**PENGARUH TUKARGANTI KATION TANAH TERHADAP  
TUMBESARAN, PENGAMBILAN NUTRIEN DAN FISIOLOGI  
ANAKBENIH SAWIT**

Oleh

**AHMAD AFANDI BIN MURDI**

**Jun 2007**

**Pengerusi : Profesor Madya Ahmad Husni Mohd Hanif, PhD**

**Fakulti : Pertanian**

Peningkatan kecekapan pengambilan nutrien akan turut menurunkan kos pengeluaran, ini kerana kos pembajaan merangkumi 30% daripada keseluruhan kos pengeluaran sawit. Meskipun, banyak penyelidikan agronomi yang telah dilaksanakan berkaitan gerakbalas sawit ke atas berbagai jenis baja, tetapi sedikit yang diterbitkan berkaitan kajian fisiologi sawit terhadap kation seperti Kalium (K), Kalsium (Ca) dan Magnesium (Mg). Oleh itu, kajian ini penting untuk meningkatkan kecekapan pengurusan nutrien sawit melalui pendekatan multidisiplin berkaitan tanah dan fisiologi pokok. Kajian dijalankan di tapak semaian Stesen Penyelidikan Lembaga Minyak Sawit Malaysia (MPOB) di Lahad Datu, Sabah. Anakbenih disemai di atas dua jenis tanah iaitu yang mengandungi Ca tukarganti yang tinggi (Famili Semporna) dan tanah rendah kandungan tukarganti Ca (Famili Lumpongan). Tanah Semporna diuji dengan kadar  $3^2$  K dan Mg manakala tanah Lumpongan dengan kadar  $3^3$  K, Ca dan Mg sebanyak tiga ulangan. Di antara parameter yang

diukur ialah pengukuran tampang, kandungan klorofil dan kandungan bandingan air laidaun. Keseluruhan pokok diagihkan mengikut bahagian pelepah, laidaun, batang dan akar. Seterusnya penentuan kandungan dan pengambilan K, Ca dan Mg. Sampel tanah diambil sebelum dan selepas kajian, seterusnya dijalankan penganalisaan kandungan kation. Pengukuran pertukaran gas daun dilakukan menggunakan alat “infrared gas analyser” dua minggu sebelum pengambilan sampel terakhir. Analisis awal tanah Lumpongan menunjukkan kandungan tukarganti K, Mg dan Ca adalah 0.35, 6.74 dan 6.07 cmol (+) kg<sup>-1</sup> manakala, di tanah Semporna kandungan tukarganti K, Mg dan Ca adalah 0.17, 1.22 dan 33.50 cmol (+) kg<sup>-1</sup>. Penggunaan baja kation meningkatkan kandungan setiap tukarganti kation secara ketara di kedua-dua tanah. Rawatan kation menunjukkan peningkatan pengeluaran bahan kering setiap komposisi anakbenih yang tidak ketara di kedua-dua tanah. Kepekatan dan pengambilan K dalam komposisi anakbenih meningkat secara ketara hasil pembajaan K untuk setiap jenis tanah. Pembajaan Mg tidak disyorkan di tanah Lumpongan tetapi baja Mg 6.69 g sepokok diperlukan bagi menampung tumbesaran anakbenih di tanah Semporna. Kecekapan pengambilan semula bagi K dan Ca di tanah Lumpongan ialah 3.65 dan 1.21% manakala, di tanah Semporna kecekapan pengambilan semula untuk K dan Mg ialah 4.56 dan 2.54%. Kajian menunjukkan bahawa, jumlah kation pada anakbenih sawit terutamanya di dalam laidaun sangat dipengaruhi oleh kandungan tukarganti Ca dalam tanah berbanding tukarganti K atau Mg. Pembahagian peratus setiap kation dibanding jumlah kation adalah seimbang pada 29% sekiranya kandungan tukarganti Ca adalah tinggi (> 25 cmol (+) kg<sup>-1</sup>), ianya dianggap mencukupi dan seimbang untuk keperluan nutrien sawit. Walau bagaimanapun, jika kandungan tukarganti Mg adalah tinggi (> 4.75 cmol (+) kg<sup>-1</sup>)



sepertimana yang terdapat pada tanah Lumpongan akan menyebabkan ketidakseimbangan pembahagian kation berbanding dengan jumlah kation. Tumbesaran anakbenih sawit di kedua-dua tanah kurang mempamerkan kepekaannya terhadap rawatan kation. Ini dibuktikan melalui tumbesaran, ketegasan tisu laidaun dan kandungan klorofil yang tidak ketara perbezaanya. Kandungan Ca yang berlebihan seperti di tanah Semporna menyumbang kepada nisbah pucuk kepada akar anakbenih yang rendah di antara 7 hingga 9 % berbanding tanah Lumpongan. Kadar fotosintesis untuk anakbenih di tanah Lumpongan dan Semporna masing-masing pada 8.52 hingga 9.45  $\mu\text{mol m}^{-2}\text{s}^{-1}$  dan 7.07 hingga 8.66  $\mu\text{mol m}^{-2}\text{s}^{-1}$ . Rawatan kation disifatkan tidak memberikan kesan yang ketara terhadap penurunan kadar fotosintesis anakbenih. Turut disimpulkan ialah, purata nilai konduktan stomata dan kepekatan  $\text{CO}_2$  antara sel mencukupi bagi menampung pertukaran gas daun, sepertimana yang dibuktikan melalui perbezaan yang tidak ketara nilai keduanya melalui rawatan kation. Hasil kajian merumuskan bahawa, kombinasi K1 (35.11 g K) + Mg0 (0 g Mg) + Ca1 (14 g Ca) di tanah Lumpongan dan K2 (70.22 g K) + Mg1 (6.69 g Mg) bagi tanah Semporna disyorkan untuk menampung tumbesaran, pengambilan nutrien dan keperluan fisiologi anakbenih sawit. Kombinasi tersebut dipilih setelah mengambil kira pengeluaran berat kering, pengambilan semula dan ciri-ciri fisiologi anakbenih sawit bagi setiap tahap kation.

## ACKNOWLEDGEMENTS

**In the name of ALLAH the Beneficial and the Compassionate.** Thank to ALLAH S.W.T, the Almighty God had given me the strength and blessings to further my study and to complete this thesis.

I would like to express my sincere gratitude to Assoc. Prof. Dr. Ahmad Husni Mohd. Hanif the Chairman of the Supervisory Committee for his advice, guidance and encouragement throughout the course of this study. Special thanks and great appreciation also extended to members of the supervisory committee, Assoc. Prof. Dr. Mohd Rafii Hj Yusop and Dr. Mohd Haniff Harun for their valuable assistance and constructive comments. Special thanks to Dr Osumanu Haruna Ahmed and Tn. Hj. Ahmad Tarmizi Mohammad for their statistical input.

Sincere appreciation and gratitude also extended to Director General of MPOB for giving me financial support and granting my part time study leave. I would like to convey my heartiest thank and appreciation to En. Ab. Hakim Ismail, En. Abdul Fatah , En. Muarad Ahmad, En. Mohd Roslan Noor, The Field Manager of FELCRA Sungai Yeh- Yeh and MPOB Lahad Datu Agronomy staff for their support during field and laboratory analysis. My deepest appreciation goes to my late parents, my beloved wife and my children whom had really support and shared the hard time together. Above all, all praises and thank be to **ALLAH**.



I certify that an Examination Committee has met on 19<sup>th</sup> June 2007 to conduct the final examination of Ahmad Afandi Bin Murdi on his Master of Agricultural Science thesis entitle “Influence of Soil Exchangeable Cations on Growth, Nutrient Uptake and Physiology of Oil Palm Seedlings” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulation 1981. The Committee recommends that the student be awarded the degree of Master of Agricultural Science.

Members of the Examination Committee are as follows:

**Aminuddin Hussin, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Mohamed Hanafi Musa, PhD**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Zakaria Wahab, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Zin Zawawi Zakari, PhD**

Malaysian Palm Oil Board  
(External Examiner)

---

**HASANAH MOHD. GHAZALI, PhD**

Professor and Deputy Dean  
School of Graduates Studies  
Universiti Putra Malaysia

Date: 24 October 2007



This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Agricultural Science. The members of the Supervisory Committee are as follows:

**Ahmad Husni Mohd. Hanif, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Mohd Rafii Hj. Yusop, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Mohd Haniff Harun, PhD**

Crop Physiology Group  
Malaysian Palm Oil Board  
(Member)

---

**AINI IDERIS, PhD**

Professor and Dean  
School of Graduates Studies  
Universiti Putra Malaysia

Date: 15 November 2007



## **DECLARATION**

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

---

**AHMAD AFANDI BIN MURDI**

Date : 4 October 2007

## TABLES OF CONTENTS

		<b>Page</b>
	<b>DEDICATION</b>	ii
	<b>ABSTRACT</b>	iii
	<b>ABSTRAK</b>	vi
	<b>ACKNOWLEDGEMENTS</b>	ix
	<b>APPROVAL</b>	x
	<b>DECLARATION</b>	xii
	<b>LIST OF TABLES</b>	xv
	<b>LIST OF FIGURES</b>	xviii
	<b>CHAPTER</b>	
<b>I</b>	<b>INTRODUCTION</b>	1
<b>II</b>	<b>LITERATURE REVIEW</b>	4
	The Performance of Oil Palm ( <i>Elaeis guineensis</i> Jacq.) in Malaysia	4
	Yield Performance and Nutritional Requirements	5
	Oil Palm Seedlings and Nursery Management	8
	Potassium	10
	Uptake and Physiological Role in Plants	10
	Potassium in Soil	12
	Deficiency Symptoms	13
	Magnesium	14
	Uptake and Physiological Role in Plants	14
	Magnesium in Soil	15
	Deficiency Symptoms	16
	Calcium	17
	Soil Calcium	17
	Uptake and Physiological Role in Plants	19
	Prediction of Optimum Bases in Palm Leaves	20
	Grading K, Ca and Mg Deficiency in Oil Palm	21
	Nutrient Balance	21
	Cations Interaction	22
	Photosynthetic, Stomatal Conductance and Intercellular CO <sub>2</sub> Concentration Characteristics	23
	Leaves Relative Water Content (LRWC)	25
	Leaf Relative Chlorophyll Content	26
	The Soils of Sabah in Relation to Exchangeable Cations	27
	Summary	28

<b>III</b>	<b>MATERIALS AND METHODS</b>	31
	Study Site	31
	Oil Palm Nursery Preparation	31
	Experimental Plot and Design	33
	Fertilizer Forms and Rates	33
	Soil Sampling and Analyses	34
	Determination of Exchangeable $K^+$ , $Ca^{2+}$ and $Mg^{2+}$ by Shaking	35
	Method	
	Determination of Total N	35
	Determination of Phosphorus	36
	Determination of soil pH	37
	Destructive Sampling of Seedlings	37
	Tissue Analyses	38
	Estimation of Selected Nutrient Content in Plant Parts	39
	Recovery Efficiency	39
	Vegetative Growth Measurement	40
	Shoot-Root Relations	41
	Leaf Gas Exchange Measurement	42
	Relative Water Content Measurement	43
	Relative Chlorophyll Content Measurement	43
	Statistical Analysis	44
<b>IV</b>	<b>RESULTS</b>	45
	The Initial Status of Soil Properties	45
	Soil Nutrient Status in Relation to Cations Application	47
	Cations Interaction	50
	Dry Matter Partitioning of Oil Palm Seedlings	52
	Cations Tissue Concentration and Uptake	57
	Recovery Efficiency of Potassium, Magnesium and Calcium	73
	Total Cations in Seedling Components	74
	Main Effects of Cations on Vegetative Growth	80
	Shoot-Root Ratio	83
	Leaf Relative Water Content	83
	Relative Chlorophyll Content	86
	Photosynthesis Rate	86
	Stomatal Conductance	88
	Intercellular $CO_2$ Concentration	90
<b>V</b>	<b>DISCUSSION</b>	92
<b>VI</b>	<b>SUMMARY</b>	97
	<b>REFERENCES</b>	99
	<b>BIODATA OF THE AUTHOR</b>	106

## LIST OF TABLES

<b>Table</b>		<b>Page</b>
1	FFB, Oil Production Level and Mature Planting Area	8
2	Classification for K Status in Rachis Analysis from Frond 17	13
3	Percentage of Individual Cation to Total Bases in Coral Soil	18
4	Soil Exchangeable Cation from Different Soil Regions	18
5	Deficiency Rating for Leaf K and Mg Based on Leaf Nutrient Level as a Proportion of TLC	21
6	Soil Exchangeable Cation and Nutrient Ratios on Coral Soil	22
7	Leaves Physiological Changes in Relation to Relative Water Content	26
8	Partial Legend of the Reconnaissance Soil Map of Sabah	32
9	Fertilizer Treatments Applied to Oil Palm Seedlings	34
10	The Initial Status (Mean $\pm$ S.E) of Soil Characteristics in Lumpongan and Semporna Soils	45
11	The Mean $\pm$ S.E of Soil N, P and pH	47
12	Soil Exchangeable K in Lumpongan Soil in Relation to K Treatment	48
13	Soil Exchangeable Mg in Lumpongan Soil in Relation to Mg Treatment	49
14	Soil Exchangeable Ca in Lumpongan Soil in Relation to Ca Treatment	49
15	Soil Exchangeable K in Semporna Soil in Relation to K Treatment	49
16	Soil Exchangeable Mg in Semporna Soil in Relation to Mg Treatment	50



17	The Analysis of Variance for Exchangeable K, Ca and Mg Concentration in Lumpongan Soil	51
18	Effects of K, Ca and Mg on Dry Matter Production of Oil Palm Seedling Components on Lumpongan Soil	55
19	Effects of K and Mg on Dry Matter Production of Oil Palm Seedling Components on Semporna Soil	56
20	Potassium Concentration in Oil Palm Seedling Components on Lumpongan Soil	59
21	Effects of Treatments on Potassium Uptake in Oil Palm Seedling Components on Lumpongan Soil	60
22	Magnesium Concentration in Oil Palm Seedling Components on Lumpongan Soil	63
23	Effects of Treatments on Magnesium Uptake in Oil Palm Seedling Components on Lumpongan Soil	64
24	Calcium Concentration of Various Seedling Components on Lumpongan Soil	65
25	Effects of Treatments on Calcium Uptake in Oil Palm Seedling Components on Lumpongan Soil	66
26	Potassium Concentration in Oil Palm Seedling Components on Semporna Soil	69
27	Effects of Treatments on Potassium Uptake in Oil Palm Seedling Components on Semporna Soil	70
28	Magnesium Concentration in Oil Palm Seedling Components on Semporna Soil	71
29	Effects of Treatments on Magnesium Uptake in Oil Palm Seedling Components on Semporna Soil	72
30	Effects of Treatments on Total Cation of Various Oil Palm Seedling Components on Lumpongan Soil	76
31	Percentage of K, Ca and Mg to Total Cation of Oil Palm Seedling Components Grown on Lumpongan Soil	77
32	Effects of Treatments on Total Cation of Various Oil Palm Seedling Components on Semporna Soil	78

33	Percentage of K and Mg to Total Cation of Oil Palm Seedling Components Grown on Semporna Soil	79
34	The Effect of Cations on Vegetative Characteristics in Lumpongan Soil	81
35	The Effect of Cations on Vegetative Characteristics in Semporna Soil	82
36	Effects of Cation Treatments on Shoot to Root Ratio on Lumpungan Soil at Two Different MAP	84
37	Effects of Cation Treatments on Shoot to Root Ratio on Semporna Soil at Two Different MAP	84
38	Effects of Cation Treatments on Relative Water Content in Oil Palm Seedlings on Lumpungan Soil at Different MAP	85
39	Effects of Cation Treatments on Relative Water Content in Oil Palm Seedlings on Semporna Soil at Different MAP	85
40	Effects of Cation Treatments on Relative Chlorophyll Content of Oil Palm Seedlings on Lumpungan Soil at Different MAP	87
41	Effects of Cation Treatments on Relative Chlorophyll Content of Oil Palm Seedlings on Semporna Soil at Different MAP	87
42	Effects of Cation Treatments on Mean of Gas Exchange Parameters on Lumpongan Soil	89
43	Effects of Cation Treatments on Mean of Gas Exchange Parameters on Semporna Soil	89
44	The Correlation Coefficient Among Photosynthesis, Stomatal conductance and Intercellular CO <sub>2</sub>	91

**LIST OF FIGURES**

<b>Figure</b>		<b>Page</b>
1	Oil Palm Efficient Nutrient System (OPENS)	6
2	Harvested Vegetative Parts	38
3	Measuring Leaf Gas Exchange Using Portable Photosynthesis System	42
4	The Exchangeable Ca in Relation to Mg x Ca Interaction in Lumpongan Soil	51

## CHAPTER I

### INTRODUCTION

The primary objectives of oil palm agronomy research are to optimize yield and to reduce cost of production. Production cost can be reduced by increasing the efficiency of nutrient uptake, since fertilizer cost is about 30% of total costs of production. In view of the very low nutrient content of most soils where oil palm is grown, adequate and balanced nutrient applications are required for optimal growth and high yield (Tarmizi *et al.*, 2003). Therefore, basic studies are needed to understand the relationships between soil-fertilizer-plant in maintaining optimum nutrition and the mechanisms of oil palm response in terms of dry matter and nutrient allocation. Nutrient use efficiency can be achieved by balancing the external nutrient inputs.

Through intensive research and development, much knowledge and information have been obtained on nutrient management in oil palm. There are still areas where improvement can be made, such as by exploiting of cation nutrient such as potassium, magnesium and calcium. The main function of cation are to balance the charge of anions or each cation absorbed, so as to maintain a neutral charge balance, except when  $\text{NH}_4^+$  is abundant (Henry and Boyd, 1996).

Therefore, each cation will assert an effect on the optimal ratio for optimal growth and maximum yield. The antagonistic interaction of K, Ca and Mg was clearly



shown on Semporna soil families, where Ca constitutes more than 70% of the ratio of the three exchangeable bases, resulting in an imbalance uptake of Mg and K and consequently affects the growth and yield of oil palm (Afandi *et al.*, 2002).

Cation, such as potassium plays an important role in cell extension, stomata regulation and other factors related to water stress. Accumulation of K produces a gradient of osmotic pressure that draws water into root. Plant deficient in K has less ability to absorb water and is subjected more to water stress. The effects of K on oil to bunch ratio in oil palms have been reported (Tarmizi, 2000).

Another essential cation is Mg which is associated with chlorophyll and its ability to convert light energy into biochemical energy during photosynthesis. Inadequate Mg in palms will reduce the oil to bunch ratio (Prabowo and Foster, 1998) and its relation to K in oil palms has been well studied (Chan and Rajaratnam, 1977; Breure, 1982).

Knowledge of some physiological aspects, such as photosynthesis, stomatal conductance and leaf relative water content of the oil palm is important for proper agronomic managements. Any limitations on physiological process, such as nutrient supply will reduce efficiency in the oil palm production system and affecting the dry matter production and yield. In general, the rate of dry matter production of crop is determined by physiological processes, such as photosynthesis, respiration, stomata conductance, translocation and transpiration. Therefore, understanding the

physiological role of mineral nutrients is essential in order to increase fertilizer efficiency.

Despite many agronomic trials on the response of oil palm to various fertilizers, there is little published information on the interdisciplinary study of accumulation and partitioning of cation nutrients in relation to physiological responses during its growth. Measurements of photosynthesis in oil palm production system are necessary for comparing and understanding productivity at leaf, palm or community level as well as their response to environmental stress (Haniff, 2005).

The objectives of this study were: (i) To quantify K, Ca and Mg concentration and uptake in various parts of oil palm seedling and their relation to oil palm seedling growth, and (ii) To quantify physiology responses of oil palm seedlings in relation to K, Ca and Mg uptake.

## CHAPTER II

### LITERATURE REVIEW

#### **The Performance of Oil Palm (*Elaeis guineensis* Jacq.) in Malaysia**

Total area planted with oil palm in Malaysia in 2005 was about 4.05 million hectares. Sabah remained the largest oil palm planted state with 1.2 million hectares or 30% of total area. As the major crop for vegetable oil in the world, palm and kernel oils represent almost 28% of the total vegetable oil production in the global market. The performance of the Malaysian oil palm industry in 2004 and half yearly 2005 remained competitive compared with other crops.

The crude palm oil (CPO) production was increased to the highest level in the Malaysian oil palm industry's history, increasing by 4.7% or 0.63 million tonnes to 13.98 million tonnes from 13.35 million tonnes in 2003. Improvement in the oil extraction rate from 19.75% in 2003 to 20.03% in 2004 and expansion of mature area are the main contribution of the significant increased of CPO production. However, the fresh fruit bunches (FFB) yield per hectare decreased by 2.1% to 18.60 tonnes from 18.99 tonnes in 2003 due to dry period encountered in early 2004 (MPOB, 2005).

The oil palm sector had contributed significantly to the Malaysia's economy during the 1997/98 regional financial crisis. However, its future expansion is going to be

limited by several factors such as depletion of prime soils for planting, shortage of workers and rising costs of labour and material (Afandi *et al.*, 2002).


In Sabah alone, there is a trend lately to develop area previously classified as marginal for agriculture including oil palm. As an example, there is a rapid increase in oil palm hectarage in the Keningau – Sook plain which is located at an altitude of about 370 m above sea level, and situated in a rain – shadow area with distinct dry season (Deratil *et al.*, 2001).

### **Yield Performance and Nutritional Requirements**

The theoretical potential oil yield of oil palm has been estimated to be about 17 t ha<sup>-1</sup> (Corley, 1985) however the recorded average oil yields in Malaysia is about 3.6 t ha<sup>-1</sup> (Ng *et al.*, 2003). Due to its large production of FFB yield and oil per hectare per year, therefore oil palm required large demand for nutrients as compared to the other crops such as soybean, seed rape and sunflower. An attempt had been made to reach the theoretical potential oil yield of oil palm through tissue culture (Ng *et al.*, 2003).

The limiting factors that may restrict yields include inadequate rainfall, shallow soil depth, slope and poor drainage, which may reduce nutrient requirement estimates. Tarmizi *et al.*(1999) incorporated the environmental data, soil chemical and texture, foliar nutrient ratio, total leaf cation balance and other factors in the oil palm fertilizer recommendation system called “Oil Palm Efficient Nutrient System” or OPENS (*Figure 1*).





**Figure 1: Oil Palm Efficient Nutrient System (OPENS)**

The nutrient requirements will depend significantly on the maximum site yield potential if the agro-management practices, such as planting of cover crops, pruning intensity and planting density are standardized. This is because fertilizers increased dry matter production mainly by improving canopy efficiency (Squire, 1983).