



UNIVERSITI PUTRA MALAYSIA

**VEGETATION AND SELECTED ENVIRONMENTAL FACTORS OF
BUKIT CHARAS LIMESTONE**

SOH WUU KUANG

FSAS 2002 29

**VEGETATION AND SELECTED ENVIRONMENTAL FACTORS OF
BUKIT CHARAS LIMESTONE**

By

SOH WUU KUANG

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of Requirement for the Degree of Master of Science**

November 2002



TO MOM AND DAD WITH LOVE



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

**VEGETATION AND SELECTED ENVIRONMENTAL FACTORS OF
BUKIT CHARAS LIMESTONE**

By

SOH WUU KUANG

November 2002

Chairperson: Rusea Go, Ph.D.

Faculty: Science and Environmental Studies

The Limestone flora of Peninsular Malaysia is exceptionally rich in diversity and endemism but is under extreme threat. Moreover, the ecology of limestone plants is little understood. Bukit Charas stands out as one of the last sanctuaries for the flora of the Panching Limestone Formation. Therefore a floristic and ecological study were undertaken to document and conserve the hill. The flora in this study stood at 249 species in 175 genera and 82 families. The largest family is the Euphorbiaceae (19) followed by the Rubiaceae (18), Araceae (13), Annonaceae (10) and Lauraceae (7). Six rare species; (*Suregada multiflora* var *lamellata*, *Ardisia suffruticosa*, *Tectaria cherasica*, *Emarhendia bettiana*, *Monophyllaea hendersonii* and *Polyalthia* sp.) are recognised from Bukit Charas. Cluster analysis has identified four types of vegetation groups at the base of the hill (BASE), at the slope of the hill (SLOPE), at the talus slope (TALUS) and at the cave (CAVE). Plants, characteristic to the limestone habitat are found in the TALUS and CAVE groups. Twelve environmental variables have been used for ordination. Unrotated principal component analysis has reduced these variables to two components, i.e. calcium and magnesium; and kalium and carbonate. Two primary environmental gradients were identified with



the Varimax-rotated principal component analysis of environmental data: topographic situation and soil fertility. The TALUS and CAVE groups were significantly associated with high level of soil exchangeable calcium and magnesium. The TALUS group is also associated with high organic matter and the CAVE group with high carbonate content. The results of the species and quadrat ordination of the detrended correspondence analysis were similar to the principal component analysis. Three hypotheses are proposed: Firstly, the plant endemism in limestone habitat is greatly driven by edaphic factor and not by the need to avoid competition. Secondly, the high level of exchangeable calcium and magnesium, and high soil pH in the limestone soil act as important limiting factors for limestone endemics. These conditions are important requirements for limestone habitats. Thirdly, there exist limiting factors within a limestone habitat that create a niche partition among the limestone endemic species. The current threat to Bukit Charas flora is agricultural activities at the surrounding base of the hill. A buffer zone that stretched 50 m from the forest margin is proposed as a strategy to conserve Bukit Charas. Future study should be expanded to other hills for comparison and green house experiment should be undertaken to test the hypotheses.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

VEGETASI DAN FAKTOR ALAM PERSEKITARAN TERPILIH BATU KAPUR BUKIT CHARAS

Oleh

SOH WUU KUANG

November 2002

Pengerusi: Rusea Go, Ph.D.

Fakulti: Sains and Pengajian Alam Sekitar

Flora batu kapur di Semenanjung Malaysia adalah kaya dengan diversiti and tumbuhan endemik, walaubagaimanapun ia sedang diancam kepupusan. Tambahan pula pengetahuan kita tentang ekologi flora batu kapur adalah cetek. Bukit Charas adalah salah satunya kawasan perlindungan terakhir bagi flora batu kapur Panching. Oleh itu, satu kajian flora dan ekologi untuk mendokumentasi dan memulihara Bukit Charas telah dijalankan. Jumlah flora dalam kajian ini adalah sebanyak 249 spesies, 175 genera and 82 famili. Famili terbesar adalah Euphorbiaceae (19), ini diikuti oleh Rubiaceae (18), Araceae (13), Annonaceae (10) dan Lauraceae (7). Sebanyak enam spesies di Bukit Charas adalah nadir (*Suregada multiflora var lamellata*, *Ardisia suffruticosa*, *Tectaria cherasica*, *Emarhendia bettiana*, *Monophyllaea hendersonii* dan *Polyalthia* sp.). Analisis kluster telah mengenal pasti empat jenis kumpulan vegetasi, iaitu vegetasi di dasar bukit (BASE), lereng bukit (SLOPE), lereng batu-batuan (TALUS) and gua (CAVE). Tumbuhan yang mencirikan habitat batu kapur didapati di dalam kumpulan TALUS dan CAVE. Sebanyak dua belas faktor persekitaran telah digunakan untuk ordinasi. Analisis komponen prinsipal 'Unrotated' telah mengurangkan faktor-faktor ini

kepada dua komponen, iaitu kalsium dan magnesium; dan potasium dan karbonat. Dua kecerunan persekitaran telah dikenalpasti melalui analisis komponen prinsipal 'Varimax-rotated': keadaan topografi dan kesuburan tanah. Kumpulan TALUS dan CAVE berkait rapat dengan kandungan kalsium and magnesium tanah yang tinggi. Kumpulan TALUS berkait rapat dengan kandungan organik tanah yang tinggi manakala, kumpulan CAVE berkait rapat dengan kandungan karbonat yang tinggi. Keputusan ordinasi spesies dan kuadrat 'detrended correspondance analysis' adalah sama dengan keputusan analisis komponen prinsipal. Tiga hipotesis telah dicadangkan: Pertama, taburan tumbuhan yang endemik kepada habitat batu kapur adalah dipengaruhi oleh faktor edafik dan bukan oleh faktor untuk mengelak daripada persaingan. Kedua, kandungan kalsium and magnesium, dan pH yang tinggi adalah ciri penting yang menghadkan taburan tumbuhan endemik kepada habitat batu kapur. Ketiga, di dalam habitat batu kapur itu sendiri, terdapat faktor penghad yang mewujudkan pemisahan 'niche' di kalangan tumbuhan endemik batu kapur. Ancaman utama Bukit Charas kini adalah daripada aktiviti pertanian yang dijalankan di dasar bukit. Oleh itu, satu saranan untuk mewujudkan satu zon mampan sejauh 50 m dari gigi hutan dicadangkan sebagai strategi untuk pemuliharaan Bukit Charas. Pada masa yang akan datang, kajian yang melibatkan bukit batu kapur lain dicadangkan supaya dapat dibuat perbandingan and selain daripada itu, eksperimen rumah hijau juga patut dibuat untuk mengesahkan hipotesis yang dibuat.

ACKNOWLEDGEMENTS

This thesis is the offspring of a great many people, for whom a mere thanks seems terribly inadequate. First and foremost, I am grateful to have Dr. Rusea Go as my supervisor and friend. I wish to thank her for the guidance, encouragement and support in completing this thesis and in realising my dream to become a good botanist. She had personally toiled over my field-work and helped me in the time of difficulty. To her a world of thanks and an enormous debt of gratitude.

I deeply acknowledge the kind and valuable assistance of those who helped me in my field-works. A special thanks to Latifah Zainal Abidin, Mohd Azman Abdul Majid, Mariam Jutta Abdullah, Tessa Loh May Fong, Joannes Unggang, Aslitawaty Abu Bakar, Phoon Sook Ngoh, Ibrahim Idham Muhiddin, Badrol Ahmad and Bibian Michael Diway, who selflessly contributed to the field-works; to little Kent for stimulating the working atmosphere. A special thanks to the folks of Kampung Kolek for their hospitality and generosity for allowing us to stay during the course of the field-work in Bukit Charas.

A genuine thanks to the staff at FRIM who have welcomed my presence. In particular, I am grateful to the curator of the Herbarium of FRIM, Dr. Saw Leng Guan for allowing me to use the facility; to Mr. Mat Asri Ngah Sanah and Mr. Kamarudin Salleh who guided me through the tedious and yet exciting task of identifying plants.



A grateful thanks to my co-supervisors, Dr. Saberi Othman and Dr. Japar Sidik Bujang, who guided me in research, and who read and commented on the manuscript throughout; to a warm special friend, Mariam Jutta Abdullah, who kindly offered assistance in editing the text.

A warm and very special thanks to my blood-brother, Tan Teng Ju, who helped me with the soil analysis and endured complaints of fatigue from the long hours of field-works and lab works.

Finally, an extra special word of thanks to my parents, Soh Onn Fatt and Ho Kim Lan; to my loving sister, Soh Sheue Yann; to Victor Lee Boon Chai; to my brother, Soh Wu Shyong. Thank you for encouraging and supporting me throughout my undertaking. To them, much more thanks.



I certify that an Examination Committee met on 11th November 2002 to conduct the final examination of Soh Wuu Kuang on his Master of Science thesis entitled "Vegetation and Selected Environmental Factors of Bukit Charas Limestone" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Univeristi Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Umi Kalsom Yusuf, Ph.D.

Associate Professor
Department of Biology
Faculty of Science and Environmental Studies,
Univeristi Putra Malaysia
(Chairperson)

Rusea Go, Ph.D.


Department of Biology
Faculty of Science and Environmental Studies,
Univeristi Putra Malaysia
(Member)

Japar Sidik Bujang, Ph.D.

Associate Professor,
Department of Biology
Faculty of Science and Environmental Studies,
Univeristi Putra Malaysia
(Member)

Saberi Othman, Ph.D.

Associate Professor,
Department of Biology
Faculty of Science and Environmental Studies,
Univeristi Putra Malaysia
(Member)



SHAMSHER MOHAMAD RAMADILI, Ph.D.
Professor / Deputy Dean,
School of Graduate Studies,
Universiti Putra Malaysia

Date: **25 NOV 2002**

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

Rusea Go, Ph.D.

Department of Biology
Faculty of Science and Environmental Studies,
Univeristi Putra Malaysia
(Chairperson)

Japar Sidik Bujang, Ph.D.

Associate Professor,
Department of Biology
Faculty of Science and Environmental Studies,
Univeristi Putra Malaysia
(Member)

Saberi Othman, Ph.D.

Associate Professor,
Department of Biology
Faculty of Science and Environmental Studies,
Univeristi Putra Malaysia
(Member)



AINI IDERIS, Ph.D.
Professor / Dean,
School of Graduate Studies,
Universiti Putra Malaysia.

Date: : 9 JAN 2003

DECLARATION

I hereby declare that the thesis is based on my original work except for equations and citations which have been duly acknowledge. I also declare that it has not been previously or concurrently submitted for any other degrees at UPM or other institutions.

(SOH WUU KUANG)

Date:

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xv
LIST OF FIGURES	xvii
LIST OF PLATES	xix
LIST OF ABBREVIATIONS	xx
 CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	6
2.1 The Limestone Flora	6
2.2 The Ecology of Vegetation Over Limestone	10
2.2.1 Limestone Hill Habitats	10
2.2.2 Topography and Edaphic Conditions	13
2.2.3 Nutrient Cycling	17
2.2.4 Vegetation and Environmental Factors	18
2.2.5 Calcifuge and Calcicole Habit of Plants	21
2.3 The Threats to Limestone Flora	23
2.4 Conservation of Limestone Flora	26
2.5 Bukit Charas	28
2.5.1 General Geography and Geology	28
2.5.2 Climate	30
2.5.3 The Flora	30
3 METHODOLOGY	32
3.1 Preliminary Checklist	32
3.2 Preliminary Surveys	32
3.3 Field Method	33
3.4 Vegetation Variables	36
3.5 Environmental Variables	37
3.5.1 Samples Preparation	38
3.5.2 Soil Moisture and Organic Matter Determinations	39
3.5.3 pH Determination	40
3.5.4 Total Carbonate	40
3.5.5 Available Phosphate (Olsen Bicarbonate Extractable)	41
3.5.6 Exchangeable Cations (Na, K, Mg and Ca)	42



	3.5.7	Total Nitrogen Determination	43
	3.6	Assessment of the Status of Species	44
	3.7	Assessment of Threats and Management Needs	44
	3.8	Statistical Analysis	45
	3.8.1	Analytical Approach	45
	3.8.2	Data Preparation	47
	3.8.3	Environmental Variables Analysis	48
	3.8.4	Vegetation Analysis	51
	3.8.5	Linking the Vegetation and Environmental Data	52
	3.8.6	Computer Programmes	53
4		RESULTS	54
	4.1	The Flora	54
	4.1.1	The Size of the Flora of Bukit Charas	54
	4.1.2	Species Endemism	58
	4.1.3	Bukit Charas Vegetation in Relation to Other Limestone Areas in Peninsular Malaysia	61
	4.1.4	Rare species of Bukit Charas	62
	4.1.5	The Vegetation of Bukit Charas	70
	4.1.5.1	The Vegetation of the Hill Base	70
	4.1.5.2	The Vegetation of the Talus Slope	70
	4.1.5.3	The Vegetation of the Depression and Hill Slope	71
	4.1.5.4	The Vegetation of the Cave Overhang Ledges and Entrances	72
	4.1.5.5	The Vegetation of the Hill Summit and Cliff	72
	4.2	Environmental and Vegetational Analysis	73
	4.2.1	Cluster Analysis	73
	4.2.2	Environmental Situations of the Vegetation Groupings	78
	4.2.2	Principal Component Analysis (PCA) of Environmental Variables	82
	4.2.2.1	Unrotated PCA	82
	4.2.2.2	Rotated PCA	94
	4.2.3	Detrended Correspondance Analysis (DCA)	102
	4.2.4	Shrub Analysis	113
	4.2.5	Assumptions in Analysis	115
5		DICUSSION	117
117	5.1	Vegetation and Environmental Variations	
	5.2	Conservation Importance of The Bukit Charas Flora	122
	5.3	Threats to Bukit Charas: Human Influence and other Biotic Factors	123
	5.4	Recommendations	125
	5.4.1	Recommendations for the Protection of the Flora of Bukit Charas	125

5.4.2	Recommendations for Future Research Works	126
6	CONCLUSION	127
	REFERENCES	128
	APPENDICES	135
	BIODATA OF THE AUTHOR	160



LIST OF TABLES

Tables	Page
1 A six-point Domin cover scale	37
2 Pearson correlation matrix of environmental variables	50
3 Flora of Bukit Cheras	54
4 Species endemism at Bukit Cheras	56
5 Taxa endemic to Peninsular Malaysia	57
6 Species endemic to Pahang	59
7 Species restricted to Pahang and one other state in Peninsular Malaysia	59
8 Limestone endemic species at Bukit Charas	60
9 Tree species listed under the IUCN Red List Categories	61
10 List of quadrats and species in each community grouping of cluster analysis	76
11 The mean and median of twelve environmental variables for the four communities groupings	77
12 Soil analyses of the limestone tropical rain forest	80
13 Summary of the total variance explained by the components in PCA	83
14 Factor loadings of variables in the unrotated PCA	83
15 Probabilities of the Mann-Whitney U statistic from tests of the difference in median PCA scores (Component 1 and Component 2) for soil samples associated with vegetation groups	89
16 Factor loadings of variables in the Varimax-rotated PCA	95
17 Probabilities of the Mann-Whitney U statistic from tests of the difference in median Varimax-rotated PCA scores (Component 1 and Component 2) for soil samples associated with vegetation groups	98
18 The eigenvalue of the first four axes of the DCA of the vegetation data	103



19	Pearson correlation coefficients of DCA axis scores and environmental factors	103
20	The eigenvalue of the first four axes of the DCA of the vegetation data	111
21	Pearson correlation coefficients of DCA axis scores and environmental factors	112
22	Records of temperature for the year 1968-1999 at Kuantan Station	135
23	Records of rainfall for the year 1951-19999 at Kuantan Station	136
24	Preliminary floristic checklist of Bukit Charas	137
25	Floristic checklist of Bukit Charas	138
26	Environmental data of 46 quadrats	144
27	Herbs abundance data on transect one and two	146
28	Herb abundance data on transect three and four	148
29	Herbs abundance data on transect five and six	150
30	Shrub data on transect two and three	152
31	Shrub abundance data on transect three and four	153
32	Shrub abundance data on transect five and six	154
33	Cluster analysis amalgamation steps	155
34	Quadrat scores of unrotated and Varimax-rotated PCA	156
35	Species scores centroid of the unrotated and Varimax-rotated PCA	157
36	Quadrat scores of DCA	158
37	Species scores of DCA	159



LISTS OF FIGURES

Figure		Page
1	Sketch-map of the Area west of Kuantan, Pahang, Showing the Location the Pancing Limestone and Bukit Charas (Metcalf, 1980)	29
2	Locations of Study Site and Transect Lines	35
3	Dendogram of Cluster Analysis	74
4	Factor Loadings of Unrotated PCA	84
5	The Quadrat Ordination Diagram of Unrotated PCA	84
6	Quadrat Ordination Diagram to Show the Values of Exchangeable Calcium in Each Quadrats (as quartiles)	85
7	The Quadrat Ordination Diagram to Show the Values of Exchangeable Magnesium (as quartiles)	85
8	Quadrat Ordination Diagram to How the Exchangeable Kalium Values in the Quadrats (as quartiles)	86
9	Centroid Ordination of Species Based on Unrotated PCA of the Environmental Factors	90
10	The Distribution of Limestone Endemic Species on Quadrat Ordination Diagram	91
11	The Distribution of Species Indifferent to Limestone Habitat	91
12	The Distribution of Species that are Consider Stranger to Limestone Habitat	92
13	The Factors Loadings of Varimax-rotated PCA	96
14	The Quadrat Ordination of Varimax-rotated PCA	96
15	Quartiles of Exchangeable Calcium Plotted on Quadrat Ordination Diagram of Varimax-rotated PCA	99
16	Quartiles of Exchangeable Magnesium Plotted on the Quadrat Ordination Diagram of Varimax-rotated PCA	99



17	Quartiles of pH Plotted on the Quadrat Ordination Diagram of Varimax-rotated PCA	100
18	Quartiles of Soil Organic Matter Plotted on the Quadrats Ordination Diagram of Varimax-rotated PCA	100
19	Centroid Ordination of Species Based on Varimax-rotated PCA of the Environmental Factors. Data	101
20	DCA Quadrat Ordination of the Vegetation Data	104
21	DCA Species Ordination of the Vegetation Data	105
22	DCA Quadrat Ordination of the Vegetation Data With Quadrats 31 and 32 Omitted	108
23	DCA Species Ordination of Vegetation Data With Quadrat 31 and Quadrat 32 Omitted	109
24	Quartiles of Exchangeable Calcium Content Plotted on DCA Quadrat ordination Diagram of the Vegetation Data	110
25	Quartiles of Exchangeable Magnesium Plotted on the DCA Quadrat Ordination	110
26	The Shoot Percentage of Shrub Indifferent to Limestone Habitat Plotted on the Quadrat Ordination Diagram of Unrotated PCA	114
27	The Shoot Percentage of Shrub Species Stranger to Limestone Habitat Plotted on the Quadrat Ordination Diagram	114

LIST OF PLATES

PLATE		Page
1	<i>Monophyllaea hendersonii</i> growing at the cave ledge	63
2	<i>Emarhendia bettiana</i> , a plant endemic to the Panching Limestone Formation	65
3	<i>Suregada multiflora</i> var. <i>lamellata</i> , a rare tree with red ribbed fruits (dried specimen)	65
4	A community of <i>Ardisia suffruticosa</i>	67
5	<i>Tectaria cherasica</i> , a fern endemic to Bukit Charas	67
6	A dried specimen of <i>Polyalthia</i> sp. (SWK 075)	69
7	The inflorescence of <i>Polyalthia</i> sp. (SWK 075)	69



LIST OF ABBREVIATIONS

- DCA Detrended correspondence analysis
G.P.S. Global Positioning System
KEP The herbarium of the Forest Research Institute Malaysia
KLU The herbarium of Rimba Ilmu
PCA Principal component analysis

Species acronyms:

- Agl *Agloanema simplex*
Alo *Alocasia denudata*
Ami *Amischotolype monosperma*
Amo *Amorphophallus prainii*
Ana *Anadendrum montanum*
Ard *Ardisia suffruticosa*
Ari *Arisaema laminatum*
Beg *Begonia ignorata*
Chl *Chloranthus erectus*
Cyt *Cyrtandra pendula*
Dio *Dioscorea* sp.
Dra *Draceana pendula*
Ela *Elatostema subscabrum*
Ema *Emarhendia bettiana*
Epi *Epithema saxatile*
Ery *Erythrospermum candidum*
Gym *Gymnostachyum decurrens*



Imp *Impatiens albo-flava*
 Mon *Monophyllaea hendersonii*
 Myx *Myxopyrum nervosum*
 Pas *Passiflora* sp.
 Peli *Peliosanthes teta*
 Phy *Phryhium pubinerve*
 Pil *Pilea fruticosa*
 Rap *Rhaphidophora maingayi*
 Rin *Rinorea honeri*
 Sch *Schismatoglottis calyptrata*
 Tyl *Tylophora* sp
 Ven *Ventilago oblongifolia*
 1 Sapling 1
 4 Sapling 4
 5 Sapling 5
 6 Sapling 6
 10 Sapling 10
 Adi *Adiantum latifolium*
 Nid *Asplenium nidus*
 Pel *Asplenium pellucidum*
 Bol *Bolbitis sinuate*
 Het *Heterogonium pinnatum*
 Lgy *Lygodium salicifolium*
 Nep *Nephrolepis falcata*

Phy *Phymatosorus nigrescens*

Del *Selaginella delicatula*

Min *Selaginella minutifolia*

Kec *Tectaria keckii*

CHAPTER 1

INTRODUCTION

Limestone flora of Peninsular Malaysia is a vegetation type that is exceptionally rich in diversity and endemism, while being under extreme threat. In this situation, it is often only practical or possible to conserve part of the target area. Usually detailed information on individual species may be lacking but botanists are often confident that those areas are rich in diversity (Given, 1994).

To date, studies on the limestone flora in Peninsular Malaysia has been in three major categories. They are the floristics, the ecology, and the conservation of limestone flora.

In the early days the emphasis was on the documentation of the general flora. The first comprehensive study on the limestone flora of the Peninsular Malaysia was undertaken by Henderson (1939). The account on the limestone flora is based entirely on vast botanical collections and observations on plants growing on limestone derived soil. The limestone hills in Peninsular Malaysia were classified into two broad groups, 'dry hills' and 'wet hills', and a brief description on the microhabitat of limestone flora was given. The plants collected from limestone hills were also enumerated with a total of 745 species recorded (Henderson, 1939).

The same attempt was made by Anderson (1965) to describe and record the limestone flora of Sarawak based on field observation. However, the study was a preliminary one and still awaits detailed investigation. Over 600 species of plant growing on limestone hill have been recorded in this study. The vegetation of Sarawak's limestone hills has been classified into eight broad subdivisions and the microhabitats of limestone flora were generally described. It is interesting to note that for the first time, the calcifuge and calcicole habitats of plants in relation to the soil properties was mentioned and discussed briefly, however, many of the findings were mere speculations.

The floristic study of limestone flora in Peninsular Malaysia has been amplified by Chin's work (Chin, 1977, 1979, 1983a, 1983b), which provided a revision of Henderson's account. As a result, a regional floristic key was constructed and 1216 species of vascular plants was enumerated and grouped into four degree of affinity towards limestone habitat. Chin (1977) subdivided the Peninsular Malaysia limestone habitat into nine groups and described the flora in each distinct habitat. Pteridophytes were included in the study for the first time in limestone flora research. The plants on the limestone hills have been grouped into four degree of fidelity (Braun-Blanquet, 1932).

In the late 70s investigations on the forest ecology over limestone was undertaken. Studies on the limestone vegetation in Peninsular Malaysia have been studied by Crowther (1982; 1986) and in Sarawak, particularly in Mulu National Park have been studied by Vallack (Proctor *et al.*, 1983). Both studies looked into