



**UNIVERSITI PUTRA MALAYSIA**

**DISTRIBUTION AND PATHOGENIC POTENTIAL  
OF SOIL FUSARIA FROM SELECTED OIL  
PALM HABITATS IN WEST MALAYSIA**

**HO YIN WAN**

**FP 1984 3**

Distribution and pathogenic potential  
of soil fusaria from selected oil  
palm habitats in West Malaysia

by

HO YIN WAN

A thesis  
submitted in partial fulfilment  
of the requirement for the degree of  
Doctor of Philosophy in the Dept. of Plant Protection,  
Faculty of Agriculture, Universiti Pertanian Malaysia  
(University of Agriculture Malaysia)

March 1984



## ACKNOWLEDGEMENTS

I wish to express my appreciation and sincere gratitude to my supervisor, Professor G. Varghese of the Plant Protection Department, Faculty of Agriculture, University of Agriculture Malaysia, for initiating this research project and for his invaluable guidance and advice throughout the course of this study and in the preparation of this manuscript.

I am also much indebted to Dr. G.S. Taylor, Botany Department, University of Manchester, England for supervising the studies conducted in England and who has offered many helpful comments and suggestions regarding those studies and in the preparation of this manuscript and to Professor J. Colhoun, Emeritus Professor, University of Manchester for being a consultant to this project, to Dr. D. Moore, Genetics Department, University of Manchester and Dr. M. Emes, Physiology Department, University of Manchester for their help and advice on the electrophoretic studies.

I also wish to acknowledge with thanks Mr. A. Johnston, Director of the Commonwealth Mycological Institute, Kew, England for permission to use the Institute's facilities and to Dr. C. Booth, of the Commonwealth Mycological Institute who confirmed all my identifications of the Fusarium species and who very kindly let me have access to his excellent collection of publications on Fusarium

by workers all over the world.

The statistical advice and assistance of Dr. Yap Thoo Chai of the Agronomy Department, University of Agriculture Malaysia, Dr. D.K. Friesen of Soil Science Department, University of Agriculture Malaysia, and Dr. Quah Soon Cheang of Biology Department, University of Agriculture Malaysia have been invaluable and are therefore gratefully acknowledged.

Appreciation is extended to the technical staff of both University of Manchester and University of Agriculture Malaysia for their co-operation and assistance in the laboratory and field, to Socfin Company Limited, Malaysia and Binga Plantations, Unilever Limited, Zaire for supplying oil palm seeds and seedlings and to Kumpulan Guthrie Sendirian Berhad, United Plantations Berhad, Highland and Lowlands Berhad and Kumpulan Ladang-Ladang Trengganu for their assistance and permission to sample soil in their oil palm estates.

I am grateful to the Inter-University Council for Higher Education Overseas for the award of a fellowship which enables me to carry out part of the research in England.

Finally, I wish to thank my husband, Dr. Ong Seng Huat for his constant encouragement and support throughout this project especially during the period which I have to spent away from home in England.

## CONTENTS

APPROVAL SHEET	
TITLE PAGE	i
ACKNOWLEDGEMENTS	ii
TABLE OF CONTENTS	iv
LIST OF TABLES	ix
LIST OF FIGURES AND ILLUSTRATIONS	xii
ABSTRACT	xxi
 CHAPTER I INTRODUCTION	 1
1.1 The genus <u>Fusarium</u>	1
1.2 Taxonomy of <u>Fusarium</u>	2
1.3 Biochemical approaches to taxonomy of <u>Fusarium</u>	9
Electrophoretic studies	10
Serological studies	11
1.4 Distribution of soil fusaria	15
1.5 Pathogenic significance of soil fusaria	18
Vascular wilt of cotton	18
Vascular wilt of banana	20
Vascular wilt of date palm	21
Vascular wilt of oil palm	22
1.6 Oil palm ( <u>Elaeis guineensis</u> Jacq.) as a plantation crop in Malaysia	26
1.7 Objectives of the present study	31



CHAPTER II	GENERAL METHODOLOGY AND TECHNIQUES	34
2.1	The fungus-culture techniques and identification	34
	Single spore isolation	34
	Decontaminating cultures	34
	Growing conditions for the fungus	35
	Stimulation of sporulation	36
	Preservation of cultures	37
2.2	The host - oil palm seedlings	39
	Germination of oil palm seeds	39
	Growing conditions for the oil palm seedlings	40
2.3	Preparation of inoculum	41
2.4	Inoculation of the oil palm seedlings	42
2.5	Prevention of cross contamination	43
2.6	Photography and drawings	44
2.7	Experimental design and statistical analysis	44
CHAPTER III	DISTRIBUTION OF SOIL FUSARIA IN OIL PALM HABITATS	46
	Materials and methods	46
	(i) Habitats	46
	(ii) Sampling method	48
	(iii) Isolation and enumeration of <u>Fusarium</u> species	52
	(iv) Isolation of <u>Fusarium</u> species from oil palm roots	53



	Results	54
	Discussion	71
CHAPTER IV	MORPHOLOGY AND IDENTIFICATION OF <u>FUSARIUM</u> ISOLATES	80
	Materials and methods	80
	Results	82
	Discussion	128
	Key to <u>Fusarium</u> species from oil palm soils in Malaysia	134
CHAPTER V	PATHOGENIC POTENTIAL OF SOIL FUSARIA FROM MALAYSIAN OIL PALM HABITATS	136
	Materials and methods	137
	Results	140
	Discussion	148
CHAPTER VI	PATHOGENICITY STUDIES OF MALAYSIAN <u>F. OXYSPORUM</u> AND AFRICAN <u>F.</u> <u>OXYSPORUM</u> f. sp. <u>ELAEIDIS</u>	151
6.1	Cross inoculation studies	153
	Materials and methods	153
	Results	163
	Discussion	174
6.2	Histopathology of Malaysian oil palm seedlings infected with pathogenic <u>F. oxysporum</u> isolates from Africa	177
	Materials and methods	178

Results	179
(i) Symptomology	180
(ii) Histology of uninoculated palms	188
(iii) Histology of inoculated palms	194
Discussion	211
CHAPTER VII COMPARATIVE STUDIES BETWEEN MALAYSIAN <u>F. OXYSPORUM</u> ISOLATES AND PATHOGENIC ISOLATES OF <u>F.</u> <u>OXYSPORUM</u> FROM AFRICA	220
7.1 Morphological comparisons	221
Materials and methods	221
(i) Light microscope studies	221
(ii) Scanning electron microscope (SEM) studies	221
Results	222
Discussion	233
7.2 Physiological comparisons	235
Materials and methods	235
(i) Effect of temperature	235
(ii) Effect of media	235
(iii) Effect of pH	236
Results and Discussion	237
7.3 Electrophoretic studies	246
Materials and methods	247



(i)	Culture methods and preparation of samples	247
(ii)	Preparation of extracts	247
(iii)	Determination of protein content in the extract	248
(iv)	Vertical disc-electrophoresis	248
(v)	Polyacrylamide iso-electric focusing	252
	Results	257
(i)	Vertical disc-electrophoresis	257
(ii)	Polyacrylamide iso-electric focusing	260
	Discussion	270
	CHAPTER VIII GENERAL DISCUSSION	276
	BIBLIOGRAPHY	285
	APPENDIX	
1	Composition of solid culture media used in this thesis	307
2	Composition of liquid media	309
3	Preparation of sections of palm materials for anatomical studies	310
4	Determination of protein content in extract - Lowry's Folin Test	314

## LIST OF TABLES

TABLE 1.	TOTAL PLANTED HECTARAGE UNDER OIL PALM IN WEST AND EAST MALAYSIA AS AT 31st DECEMBER, 1980 (DEPT. OF STATISTICS, MALAYSIA, 1982).	28
TABLE 2.	MAJOR CRITERIA USED IN ASSESSING SOIL SUITABILITY FOR OIL PALM (NG, 1967).	30
TABLE 3.	PHYSICAL PROPERTIES OF SOILS TAKEN AT THE SAMPLING LOCATIONS FOR SOIL FUSARIA.	56
TABLE 4.	MOISTURE, pH AND ORGANIC CONTENT OF SOIL TAKEN AT SAMPLING LOCATIONS FOR SOIL FUSARIA.	57
TABLE 5.	QUANTITATIVE ESTIMATION OF THE TOTAL POPULATION OF <u>FUSARIUM</u> SPECIES IN OIL PALM HABITATS.	59
TABLE 6.	A COMPARISON OF MEANS OF THE TOTAL POPULATION OF <u>FUSARIUM</u> SPECIES IN THE OIL PALM SOILS ACCORDING TO PALM AGE AND SAMPLE ORIGIN.	60
TABLE 7.	QUANTITATIVE ESTIMATION OF THE MOST COMMON <u>FUSARIUM</u> SPECIES IN OIL PALM SOILS.	62
TABLE 8.	A COMPARISON OF MEANS OF THE POPULATION OF THE MOST COMMON <u>FUSARIUM</u> SPECIES IN OIL PALM SOILS ACCORDING TO PALM AGE AND SAMPLE ORIGIN.	63
TABLE 9.	QUANTITATIVE ESTIMATION OF OCCASIONAL <u>FUSARIUM</u> SPECIES IN OIL PALM SOILS.	64
TABLE 10.	QUANTITATIVE ESTIMATION OF RARE <u>FUSARIUM</u> SPECIES IN OIL PALM SOILS.	65
TABLE 11.	ISOLATION OF <u>FUSARIUM</u> SPECIES FROM ROOTS OF YOUNG, MATURE AND OLD PALMS IN THE FOUR LOCATIONS.	70
TABLE 12.	NUMBER OF ISOLATES OF EACH <u>FUSARIUM</u> SPECIES CAUSING AN APPARENT REDUCTION IN GROWTH OF OIL PALM SEEDLINGS.	141

TABLE 13.	EFFECT OF INOCULATING MALAYSIAN OIL PALM SEEDLINGS WITH ISOLATES OF <u>F. SOLANI</u> , <u>F. MONILIFORME</u> AND <u>F. MONILIFORME</u> var. <u>SUBGLUTINANS</u> ON AERIAL PARAMETERS OF GROWTH.	144
TABLE 14.	EFFECT OF INOCULATING MALAYSIAN OIL PALM SEEDLINGS WITH ISOLATES OF <u>F. SEMITECTUM</u> , <u>F. EQUISETI</u> , <u>F. OXYSPORUM</u> var. <u>REDOLENS</u> , <u>F. HETEROSPORUM</u> , <u>F. ACUMINATUM</u> AND <u>F. LATERITIUM</u> ON AERIAL PARAMETERS OF GROWTH.	145
TABLE 15	EFFECT OF INOCULATING MALAYSIAN OIL PALM SEEDLINGS WITH ISOLATES OF <u>F. OXYSPORUM</u> ON AERIAL PARAMETERS OF GROWTH.	146
TABLE 16	ISOLATES OF <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> FROM AFRICA.	152
TABLE 17	EFFECT OF INOCULATING MALAYSIAN OIL PALM SEEDLINGS WITH AFRICAN ISOLATES ON DISEASE DEVELOPMENT AND ROOT INFECTION.	164
TABLE 18	EFFECT OF INOCULATING MALAYSIAN OIL PALM SEEDLINGS WITH AFRICAN ISOLATES ON AERIAL PARAMETERS OF GROWTH.	165
TABLE 19	EFFECT OF INOCULATING MALAYSIAN OIL PALM SEEDLINGS WITH AFRICAN ISOLATES, WITH OR WITHOUT AN INITIAL PERIOD OF WATER STRESS, ON AERIAL PARAMETERS OF GROWTH.	167
TABLE 20	EFFECT OF INOCULATING MALAYSIAN OIL PALM SEEDLINGS WITH AFRICAN ISOLATES, WITH OR WITHOUT AN INITIAL PERIOD OF WATER STRESS, ON DISEASE DEVELOPMENT AND ROOT INFECTION.	168
TABLE 21	EFFECT OF INOCULATING AFRICAN OIL PALM SEEDLINGS WITH MALAYSIAN ISOLATES ON AERIAL PARAMETERS OF GROWTH AND ROOT INFECTION.	169
TABLE 22	EFFECT OF INOCULATING AFRICAN OIL PALM SEEDLINGS WITH ISOLATES WITH AND WITHOUT AN INITIAL PERIOD OF WATER STRESS, ON AERIAL PARAMETERS OF GROWTH AND ROOT INFECTION.	170

TABLE 23	EFFECT OF INOCULATING MALAYSIAN OIL PALM SEEDLINGS WITH MALAYSIAN ISOLATES, WITH OR WITHOUT AN INITIAL PERIOD OF WATER STRESS, ON AERIAL PARAMETERS OF GROWTH AND ROOT INFECTION.	172
TABLE 24	CORRELATION COEFFICIENTS BETWEEN DRY WEIGHT AND VARIOUS AERIAL PARAMETERS OF LEAFINESS OF MALAYSIAN OIL PALM SEEDLINGS INOCULATED WITH THE AFRICAN ISOLATES.	173
TABLE 25	CULTURAL CHARACTERISTICS OF 13 ISOLATES OF <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> FROM AFRICA.	223
TABLE 26	CULTURAL CHARACTERISTICS OF <u>F. OXYSPORUM</u> FROM MALAYSIA.	
TABLE 27	EFFECT OF TEMPERATURE ON THE GROWTH OF THE MALAYSIAN <u>F. OXYSPORUM</u> ISOLATES AND THE <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> ISOLATES FROM AFRICA.	238
TABLE 28	EFFECT OF SOLID AND LIQUID MEDIA ON THE GROWTH OF MALAYSIAN <u>F. OXYSPORUM</u> AND <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> ISOLATES FROM AFRICA.	239
TABLE 29	EFFECT OF pH ON GROWTH ON PSA OF THE MALAYSIAN <u>F. OXYSPORUM</u> ISOLATES AND THE <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> ISOLATES FROM AFRICA.	245
TABLE 30	ELECTRODE SOLUTIONS AND ELECTRIC CONDITIONS FOR ISOELECTRIC FOCUSING.	254
TABLE 31	COMPOSITION OF SOLUTIONS FOR FIXING, STAINING, DESTAINING AND PRESERVING GELS WITH PROTEINS SEPARATED BY ELECTROFOCUSING.	255
TABLE 32	INDEX OF SIMILARITY IN SOLUBLE PROTEIN BANDS WITHIN AND BETWEEN THE PATHOGENIC AFRICAN ISOLATES AND THE MALAYSIAN ISOLATES.	259

## LIST OF FIGURES AND ILLUSTRATIONS

FIGURE 1.	SPOROCHIA PRODUCED ON CARNATION LEAF AGAR.	38
FIGURE 2.	OIL PALM SEEDLINGS OF TWO-LEAF STAGE GROWING IN A SAND BED.	41
FIGURE 3.	MAP OF PENINSULAR MALAYSIA SHOWING THE FOUR SAMPLING LOCATIONS.	47
FIGURE 4.	YOUNG OIL PALMS (1-2 YEARS OLD) WITH A MIXED LEGUMINOUS COVER.	49
FIGURE 5.	MATURE OIL PALMS (7 YEARS OLD) WITH NATURAL COVER OF GRASSES AND FERNS.	49
FIGURE 6.	OLD OIL PALMS (OVER 25 YEARS OLD) WITH NATURAL COVER OF GRASSES AND FERNS.	50
FIGURE 7.	DIAGRAMMATIC REPRESENTATION OF SAMPLING AREA IN OIL PALM HABITAT.	51
FIGURE 8.	<u>FUSARIUM</u> COLONIES ON PEPTONE-PCNB AGAR MEDIUM.	55
FIGURE 9.	HISTOGRAMS SHOWING MEAN NUMBERS OF <u>FUSARIUM</u> PROPAGULES ISOLATED FROM SOILS OF DIFFERENT LOCATIONS, SAMPLE ORIGIN (RHIZOSPHERE, AVENUE) AND PALM AGE AREAS.	66
FIGURE 10.	FREQUENCY OF ISOLATION OF VARIOUS <u>FUSARIUM</u> SPECIES FROM THE FOUR LOCATIONS.	67
FIGURE 11.	MICROCONIDIA OF <u>F. SOLANI</u> .	83
FIGURE 12.	MACROCONIDIA OF <u>F. SOLANI</u> .	83
FIGURE 13.	CULTURES OF <u>F. SOLANI</u> .	85
FIGURE 14.	CULTURES OF <u>F. SOLANI</u> .	88
FIGURE 15.	LINE DRAWINGS OF THE CONIDIA OF THE VARIOUS MORPHOLOGICAL TYPES OF <u>F. SOLANI</u> .	92
FIGURE 16.	MICROCONIDIOPHORES WITH MICROCONIDIA OF <u>F. OXYSPORUM</u> .	93





FIGURE 17.	ELABORATELY BRANCHED MACROCONIDIO- PHORES PRODUCING MASSES OF MACRO CONIDIA OF <u>F. OXYSPORUM</u> .	93
FIGURE 18.	MACROCONIDIA AND MICROCONIDIA OF <u>F. OXYSPORUM</u> .	94
FIGURE 19.	SCANNING ELECTRONMICROGRAPHS SHOWING ANASTOMOSES OF MACROCONIDIA IN <u>F.</u> <u>OXYSPORUM</u> .	95
FIGURE 20.	CHLAMYDOSPORES OF <u>F. OXYSPORUM</u> .	97
FIGURE 21.	CULTURE OF <u>F. OXYSPORUM</u> SHOWING SCLEROTIAL BODIES.	98
FIGURE 22.	MORPHOLOGICAL TYPE 1 (M1) OF <u>F.</u> <u>OXYSPORUM</u> .	98
FIGURE 23.	MORPHOLOGICAL TYPES 2 (M2) AND 3 (M3) OF <u>F. OXYSPORUM</u> .	101
FIGURE 24.	MORPHOLOGICAL TYPE 4 OF <u>F. OXYSPORUM</u> .	103
FIGURE 25.	LINE DRAWINGS OF THE CONIDIA OF THE VARIOUS MORPHOLOGICAL TYPES IN <u>F.</u> <u>OXYSPORUM</u> .	104
FIGURE 26.	TOP VIEW OF <u>F. OXYSPORUM</u> var. <u>REDOLENS</u> CULTURE.	106
FIGURE 27.	MACROCONIDIA OF <u>F. OXYSPORUM</u> var. <u>REDOLENS</u> .	106
FIGURE 28.	BRANCHED CONIDIOPHORES WITH POLY- BLASTIC CONIDIOGENOUS CELLS OF <u>F. SEMITECTUM</u> .	108
FIGURE 29.	MACROCONIDIA OF <u>F. SEMITECTUM</u> .	108
FIGURE 30.	BRANCHED SIMPLE PHIALIDES OF <u>F.</u> <u>SEMITECTUM</u> PRODUCING SECONDARY MACROCONIDIA	109
FIGURE 31.	MACROCONIDIA OF <u>F. SEMITECTUM</u> PRODUCING SMALL MICROCONIDIUM- LIKE STRUCTURE.	109
FIGURE 32.	CULTURES OF <u>F. SEMITECTUM</u> .	111
FIGURE 33.	MICROCONIDIOPHORE WITH A MICRO- CONIDIUM OF <u>F. MONILIFORME</u> .	112
FIGURE 34.	MICROCONIDIA OF <u>F. MONILIFORME</u> IN CHAINS.	112

FIGURE 35.	MICROCONIDIA OF <u>F. MONILIFORME</u> IN CHAINS.	114
FIGURE 36.	MACROCONIDIA OF <u>F. MONILIFORME</u> ,	114
FIGURE 37.	CULTURES OF <u>F. MONILIFORME</u> .	115
FIGURE 38.	LINE DRAWINGS OF <u>F. MONILIFORME</u> .	116
FIGURE 39.	POLYPHIALIDES OF <u>F. MONILIFORME</u> var. <u>SUBGLUTINANS</u> .	120
FIGURE 40.	CULTURES OF <u>F. ACUMINATUM</u> .	121
FIGURE 41.	MACROCONIDIA OF <u>F. ACUMINATUM</u> .	122
FIGURE 42.	REVERSE SIDE OF <u>F. EQUISETI</u> (A) AND <u>F. HETEROSPORUM</u> (B) CULTURES.	124
FIGURE 43.	CONIDIA OF <u>F. EQUISETI</u> .	124
FIGURE 44.	CULTURE OF <u>F. LATERITIUM</u> .	126
FIGURE 45.	CONIDIA OF <u>F. LATERITIUM</u> .	126
FIGURE 46.	<u>FUSARIUM HETEROSPORUM</u> .	127
FIGURE 47.	CULTURAL VARIANTS (PATCHES AND SECTORS) IN <u>F. SOLANI</u> AND <u>F.</u> <u>ACUMINATUM</u> ).	132
FIGURE 48.	REDUCTION OF GROWTH IN THE ABSENCE OF DISEASE SYMPTOMS FOLLOWING INOCULATION WITH <u>F. SOLANI</u> ISOLATE.	147
FIGURE 49.	REDUCTION OF ROOT DEVELOPMENT IN THE ABSENCE OF DISEASE SYMPTOMS FOLLOWING INOCULATION WITH <u>F.</u> <u>SOLANI</u> ISOLATE.	147
FIGURE 50.	CALIBRATION CURVE OF SOIL MOISTURE CONTENT AND ITS EQUIVALENT WATER POTENTIAL.	156
FIGURE 51.	AN OIL PALM SEEDLING IN THE WATER STRESS EXPERIMENT.	157
FIGURE 52.	GRAPH OF CORRECTION FACTOR FOR GREEN LEAF AREA BY METER READING.	161
FIGURE 53.	YOUNG UNDERGROUND (a) AND OLD AERIAL (b) PNEUMATHODES (PURVIS, 1956).	181

FIGURE 54.	VARIOUS STAGES OF SYMPTOM DEVELOPMENT OF VASCULAR WILT ON MALAYSIAN OIL PALM SEEDLINGS.	182
FIGURE 55.	INFECTED OIL PALM BULB CUT TRANSVERSELY AND LONGITUDINALLY TO SHOW VASCULAR DISCOLOURATION.	182
FIGURE 56.	DISEASED OIL PALM BULBS CUT TRANSVERSELY TO SHOW DISCOLOURATION OF VASCULAR SYSTEM.	183
FIGURE 57.	MALAYSIAN OIL PALM SEEDLING INFECTED WITH VASCULAR WILT DISEASE SHOWING WILTED LEAVES AND POORLY DEVELOPED NECROTIC PRIMARY ROOTS (DISEASE INDEX = 4).	185
FIGURE 58.	LEAVES FROM AN OIL PALM SEEDLING INITIALLY SHOWING SYMPTOMS OF VASCULAR WILT BUT LATER RECOVERED.	186
FIGURE 59.	ROOT SYSTEMS OF OIL PALM SEEDLINGS WITH INITIAL SYMPTOMS OF VASCULAR WILT BUT LATER RECOVERED.	187
FIGURE 60.	TRANSVERSE SECTIONS OF A HEALTHY ROOT FROM A CONTROL PALM.	189
FIGURE 61.	TRANSVERSE SECTION OF A HEALTHY BULB FROM A CONTROL PALM SHOWING THE VASCULAR BUNDLES (v) SCATTERED IN THE GROUND PARENCHYMA.	192
FIGURE 62.	TRANSVERSE SECTIONS OF A HEALTHY LEAF FROM A CONTROL PALM.	193
FIGURE 63.	TRANSVERSE SECTION OF A SLIGHTLY INFECTED OIL PALM ROOT SHOWING STRONGLY LIGNIFIED CELLS IN THE HYPODERMIS (h), LIGHTLY STAINED PARENCHYMA CELLS IN THE CORTEX (c) AND INFECTED VASCULAR CYLINDER (v).	196
FIGURE 64.	TRANSVERSE SECTION OF AN INFECTED PRIMARY OIL PALM ROOT SHOWING CONTINUOUS VASCULAR CONNECTION BETWEEN PRIMARY ROOT (pr) AND INFECTED LATERAL ROOT (lr), MYCELIUM NOT DETECTED IN CORTICAL CELLS (c).	196



FIGURE 65.	TRANSVERSE SECTIONS OF AN INFECTED OIL PALM ROOT SHOWING VESSEL WITH CONIDIA (c), HYPHAE (h), CHLAMYDOSPORE (ch) AND TRACHEIDS (t) PLUGGED WITH CONIDIA AND HYPHAE.	197
FIGURE 66.	TRANSVERSE SECTION OF AN INFECTED OIL PALM ROOT SHOWING INFECTED XYLEM ELEMENTS (x) BUT UNAFFECTED PHLOEM CELLS (p).	198
FIGURE 67.	LONGITUDINAL SECTION OF AN INFECTED OIL PALM ROOT SHOWING XYLEM VESSEL (v) WITH HYPHAE, ADJACENT TRACHEIDS OCCLUDED WITH GUMS AND DEPOSITS (g) AND DISINTEGRATING CORTICAL CELL (c).	199
FIGURE 68.	LONGITUDINAL SECTION OF AN INFECTED OIL PALM ROOT SHOWING XYLEM ELEMENTS COMPLETELY OCCLUDED WITH GUMS AND OTHER MATERIALS (ARROWS).	199
FIGURE 69.	LONGITUDINAL SECTIONS OF AN INFECTED OIL PALM ROOT SHOWING TYLOSES IN XYLEM VESSELS.	200
FIGURE 70.	TRANSVERSE SECTION OF AN INFECTED OIL PALM ROOT SHOWING DEPOSITION OF ADDITIONAL WALL LAYER IN PHLOEM CELLS (p) AND CAVITIES IN PHLOEM BEING FILLED WITH DARK STAINING MATERIALS (s).	200
FIGURE 71.	TRANSVERSE SECTIONS OF AN INFECTED OIL PALM ROOT SHOWING DISINTEGRATION OF PHLOEM CELLS (p), VESSEL FILLED WITH CONIDIA (v) AND TRACHEIDS PLUGGED WITH GUM (g).	201
FIGURE 72.	LONGITUDINAL SECTION OF AN INFECTED OIL PALM ROOT SHOWING DISINTEGRATING XYLEM ELEMENTS (ARROWS) AND COMPLETELY DISINTEGRATED CORTICAL CELLS RESULTING IN CAVITIES (c) ON EITHER SIDE OF THE CENTRAL CYLINDER	203
FIGURE 73.	TRANSVERSE SECTION OF AN INFECTED OIL PALM ROOT SHOWING DEVELOPMENT OF CAVITIES AND GAPS (g) IN THE HYPODERMIS (h) AND CORTEX (c).	203

FIGURE 74.	LONGITUDINAL SECTION OF AN INFECTED OIL PALM ROOT SHOWING LUMEN OF HYPODERMAL CELL (h) PLUGGED WITH GUMS, AND DISINTEGRATING EPIDERMAL CELLS (e) AND CORTICAL CELLS (c).	204
FIGURE 75.	TRANSVERSE SECTION OF AN INFECTED OIL PALM ROOT SHOWING CAVITY (c) IN THE PITH FORMED BY DISINTEGRATING CELLS.	204
FIGURE 76.	TRANSVERSE SECTION OF AN INFECTED OIL PALM ROOT SHOWING THE DEVELOPMENT OF LARGE CAVITIES (ARROWS) FORMED FROM THE DISINTEGRATION OF INNER CORTICAL CELLS BETWEEN THE LACUNAE.	205
FIGURE 77.	TRANSVERSE SECTION OF AN INFECTED OIL PALM ROOT DURING LATE PATHOGENESIS SHOWING THE CENTRAL VASCULAR CYLINDER (cv) COMPLETELY DETACHED FROM THE OUTER CORTEX (oc).	205
FIGURE 78.	TRANSVERSE SECTION OF AN INFECTED OIL PALM ROOT SHOWING INTACT ENDODERMIS (e), DISINTEGRATING CORTICAL CELLS (c) AND EXTENSIVE HYPHAL DEVELOPMENT IN THE PHLOEM, XYLEM AND PITH CELLS (ARROWS).	206
FIGURE 79.	LONGITUDINAL SECTION OF AN INFECTED OIL PALM BULB SHOWING XYLEM VESSELS WITH HYPHAE (h) AND CHLAMYDOSPORES (ch) OR OCCLUDED WITH GUMS (g).	206
FIGURE 80.	TRANSVERSE SECTION OF AN INFECTED OIL PALM BULB SHOWING HYPHAE PASSING THROUGH THE WALLS OF CONTIGUOUS VESSEL ELEMENTS (ARROWS)	208
FIGURE 81.	TRANSVERSE SECTION OF AN INFECTED VASCULAR BUNDLE OF AN OIL PALM BULB SHOWING DISINTEGRATING PHLOEM CELLS (p) AND GROUND PARENCHYMA CELLS (gp) WITH CRUSHED OR DISTORTED CELL WALLS AND A BAND OF CELLS OCCLUDED WITH GUMS (g).	208

FIGURE 82.	TRANSVERSE SECTION OF AN INFECTED OIL PALM BULB SHOWING SMALL CAVITIES OR GAPS (g) FORMED FROM THE DISINTEGRATION OF OCCLUDED XYLEM ELEMENTS AND LARGE CAVITIES (gp) FORMED FROM DISINTEGRATING GROUND PARENCHYMA CELLS.	209
FIGURE 83.	TRANSVERSE SECTION OF AN INFECTED OIL PALM BULB SHOWING OCCLUDED VASCULAR BUNDLES (v) AND LARGE CAVITIES (c) BETWEEN THE VASCULAR BUNDLES.	209
FIGURE 84.	LONGITUDINAL SECTION OF AN INFECTED OIL PALM LEAF PETIOLE SHOWING AN OCCLUDED VASCULAR BUNDLE.	212
FIGURE 85.	TRANSVERSE SECTION OF A WILTED LEAF FROM AN INFECTED OIL PALM SEEDLING SHOWING CRUSHED EPIDERMAL (e), HYPODERMAL (h) AND MESOPHYLL (m) CELLS.	212
FIGURE 86.	CULTURES OF <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> .	224
FIGURE 87.	CULTURES OF <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> WITH ABUNDANT MYCELIUM.	224
FIGURE 88.	MACROCONIDIA OF <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> (FROM LOBE 2 ISOLATE),	225
FIGURE 89.	CULTURES OF <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> WITH A VIOLET TO PURPLISH PIGMENTATION.	226
FIGURE 90.	CULTURE OF YALIGIMBA ISOLATE.	226
FIGURE 91.	SCANNING ELECTRONMICROGRAPHS OF MACROCONIDIA OF <u>F. OXYSPORUM</u> FROM MALAYSIA (A) AND <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> FROM AFRICA (B).	229
FIGURE 92.	SCANNING ELECTRONMICROGRAPHS OF APICAL AND FOOT CELLS OF MACROCONIDIA OF <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> FROM AFRICA (A) AND <u>F. OXYSPORUM</u> FROM MALAYSIA (B-C).	230
FIGURE 93	SCANNING ELECTRONMICROGRAPHS OF BRANCHED MACROCONIDIOPHORES OF <u>F. OXYSPORUM</u> FROM MALAYSIA (A) AND <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> FROM AFRICA (B).	231



FIGURE 94.	SCANNING ELECTRONMICROGRAPHS OF MICROCONIDIA OF <u>F. OXYSPORUM</u> FROM MALAYSIA (A) AND <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> FROM AFRICA (B).	232
FIGURE 95.	ISOLATES OF <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> FROM AFRICA GROWN ON POTATO SUCROSE AGAR (PSA) AND KOMADA'S MEDIUM (KM).	241
FIGURE 96.	ISOLATES OF MALAYSIAN <u>F. OXYSPORUM</u> ON KOMADA'S MEDIUM.	242
FIGURE 97.	ISOLATES OF MALAYSIAN <u>F. OXYSPORUM</u> ON KOMADA'S MEDIUM.	243
FIGURE 98.	DIAGRAMMATIC REPRESENTATION OF SOLUBLE PROTEIN OF AFRICAN <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> AND MALAYSIAN <u>F. OXYSPORUM</u> ISOLATES.	258
FIGURE 99.	DIAGRAMMATIC REPRESENTATION OF ESTERASE PATTERNS OF <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> ISOLATES FROM AFRICA AND <u>F. OXYSPORUM</u> ISOLATES FROM MALAYSIA.	261
FIGURE 100.	DENSITOMETRIC TRACING OF ESTERASE PATTERNS (SEPARATED BY VERTICAL DISC ELECTROPHORESIS) OF SOME MALAYSIAN <u>F. OXYSPORUM</u> ISOLATES and <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> ISOLATES FROM AFRICA.	262
FIGURE 101.	SOLUBLE PROTEINS OF SOME AFRICAN AND MALAYSIAN ISOLATES SEPARATED BY ISOELECTRIC FOCUSING ON POLY-ACRYLAMIDE GELS OF pH 3.5-9.5.	263
FIGURE 102.	DIAGRAMMATIC REPRESENTATION OF SOLUBLE PROTEIN PATTERNS OF AFRICAN <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> ISOLATES AND MALAYSIAN <u>F. OXYSPORUM</u> ISOLATES.	264
FIGURE 103.	SOLUBLE PROTEINS OF SOME AFRICAN AND MALAYSIAN ISOLATES SEPARATED BY ISOELECTRIC FOCUSING ON POLY-ACRYLAMIDE GEL OF pH 5.5-8.5.	266
FIGURE 104.	DIAGRAMMATIC REPRESENTATION OF SOLUBLE PROTEIN PATTERNS OF AFRICAN <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> AND MALAYSIAN <u>F. OXYSPORUM</u> ISOLATES.	267

FIGURE 105.	ESTERASES OF SOME AFRICAN AND MALAYSIAN ISOLATES SEPARATED BY ISOELECTRIC FOCUSING ON POLY- ACRYLAMIDE GELS OF pH 3.5-9.5.	268
FIGURE 106.	DIAGRAMMATIC REPRESENTATION OF ESTERASE ZYMOGRAMS OF AFRICAN <u>F. OXYSPORUM</u> f. sp. <u>ELAEIDIS</u> ISOLATES AND MALAYSIAN <u>F.</u> <u>OXYSPORUM</u> ISOLATES ON POLYACRY- LAMIDE GELS WITH A pH RANGE OF 3.5-9.5.	269

## ABSTRACT

An Abstract of the thesis presented to the Senate of Universiti Pertanian Malaysia in partial fulfilment of the requirements for the Degree of Doctor of Philosophy

### DISTRIBUTION AND PATHOGENIC POTENTIAL OF SOIL FUSARIA FROM SELECTED OIL PALM HABITATS IN WEST MALAYSIA

By

Ho Yin Wan

March, 1984

Supervisor : George Varghese, Ph.D.

Faculty : Agriculture

A total of eight species and two varieties of Fusarium was isolated from the sampling sites in the oil palm habitat. Fusarium solani and Fusarium oxysporum were the most prevalent species followed by Fusarium semitectum. The other species and varieties isolated showed a more





sporadic occurrence. Generally, soils from oil palm rhizospheres and young palm areas contained a larger number and greater variety of Fusarium species than soils from the avenues and older palm areas.

Pathogenicity tests of Fusarium species isolated showed that none were capable of producing vascular wilt or other diseases on oil palm seedlings. Some of the isolates, however, caused a reduction of growth in the test seedlings.

Comparative studies of F. oxysporum isolates from oil palm habitat in Malaysia with F. oxysporum f. sp. elaeidis isolates from Africa showed that the two groups of isolates were indistinguishable in their cultural, morphological and isozyme characteristics. Subsequent pathogenicity tests proved that the F. oxysporum isolates from Africa were pathogenic, causing vascular wilt on the Malaysian oil palm seedlings whilst the F. oxysporum isolates from Malaysia were non-pathogenic to the wilt-susceptible African oil palm seedlings and Malaysian oil palm seedlings. Inoculation of Malaysian F. oxysporum isolates on Malaysian oil palm seedlings and wilt-susceptible African oil palm seedlings, subjected to an initial period of water stress, also did not result in showing any disease symptoms.

Histopathological studies of Malaysian oil palm seedlings inoculated with pathogenic F. oxysporum f. sp. elaeidis indicated that resistance of the symptomless palms to the vascular wilt is probably biochemical in nature.