



UNIVERSITI PUTRA MALAYSIA

**THE COCOA BLACK ANT-MEALYBUG RELATIONSHIP: ARTIFICIAL
ESTABLISHMENT OF CATAENOCOCCUS HISPIDUS (HOMOPTERA:
PSEUDOCOCCIDAE) ON COCOA**

ANG BAN NA

FP 1988 5

THE COCOA BLACK ANT-MEALYBUG RELATIONSHIP:
ARTIFICIAL ESTABLISHMENT OF CATAENOCOCCUS HISPIDUS
(HOMOPTERA: PSEUDOCOCCIDAE) ON COCOA

by
Ang Ban Na

A thesis submitted in partial fulfilment of the requirements for
the degree of Master of Agricultural Science
in the Faculty of Agriculture, Universiti Pertanian Malaysia.

June, 1988



ACKNOWLEDGEMENTS

I wish to thank my supervisors Associate Professor Dr. Khoo Khay Chong and Puan Rita Muhamad for their keen interest and support throughout the course of this project. I am indebted to Dr. Khoo for arranging for financial assistance without which this study would have been difficult.

I am very grateful to Mr. B.J. Wood, Research Director of Ebor Research, Sime Darby Plantations, for his interest and help: the research grant and material support provided by Sime Darby are gratefully acknowledged.

I also wish to record my thanks to Mr. Chung Gait Fee, Agronomist, Ebor Research, for generously making available his time and facilities. Appreciation also goes to Mr. Chong Choon Fong, Prang Besar Research Station, Harrisons Malaysia Plantations Berhad, for granting me access to the HMPB cocoa fields in Teluk Intan.

Thanks are also due to Dr. Khoo Soo Ghee, Zoology Department, University of Malaya, for helping with identification of the mealybugs, Prof. M.J. Way, Imperial College, Silwood Park, for his interest and useful suggestions and Dr. Yusuf Ibrahim, Plant Protection Department, Universiti Pertanian Malaysia for helping in the translation of the abstract to Bahasa Malaysia.

My sincere thanks also goes to the staff members of the Plant Protection Department, Universiti Pertanian Malaysia, Ebor Research and Prang Besar Research Station in Teluk Intan. In



particular I would like to mention the following: R. Selvarajan, Hishammudin Zainuddin, Abd. Rahman Mohamed, Kasbi Sharif, K. Chandrasekaran, N. Konasagran, Catherine Arul and P. Selvaraju.

Finally, I wish to thank my family for all their encouragement and understanding throughout the course of this study.

TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGEMENTS	ii
TABLE OF CONTENTS	iv
LIST OF TABLES	vii
LIST OF FIGURES	ix
LIST OF PLATES	x
ABSTRACT	xi
ABSTRAK	xiv
CHAPTER 1 INTRODUCTION	1
CHAPTER 2 LITERATURE REVIEW	5
CHAPTER 3 METHOD FOR ESTIMATION OF NUMBER OF INDIVIDUALS IN A COLONY OF MEALYBUGS BY ITS AREA	14
Literature review	14
Colony composition	15
Materials and method	16
Results and Discussion	18
Estimation of number of individual in first instar colony through its area	20
Materials and method	20
Results and Discussion	20
CHAPTER 4 TIMING OF MEALYBUG INTRODUCTION WITH RESPECT TO THAT OF THE ANTS	24
Literature review	24
Preliminary experiment to determine influence of mealybug and timing of its introduction on the establishment of the ants	25
Materials and methods	25



Results and discussion	28
Optimum length of time between mealybug and ant introduction for mealybug establishment	31
Materials and Method	31
Results and discussion	32
CHAPTER 5 THE EFFECT OF RAINFALL AND NATURAL ENEMIES ON THE ESTABLISHMENT OF MEALYBUGS IN THE ABSENCE OF ANTS	35
Literature Review	35
Materials and method	36
Results and discussion	38
Conclusion	50
CHAPTER 6 CRAWLER BEHAVIOUR	52
Literature review	52
Influence of starvation on mobility of first instar crawlers	53
Materials and method	53
Results and discussion	54
Response of first instar crawlers to light and gravity	54
Materials and method	56
Results and discussion	59
Effect of artificial scarification of pods, on crawler establishment	62
Materials and method	64
Results and discussion	65
Influence of humidity and temperature on the survival of starved first instar crawlers	65
Materials and method	68
Results and discussion	69

Conclusion	72
CHAPTER 7 INTRODUCTION OF ANTS AND MEALYBUGS AT VARIOUS TREE DENSITIES	73
Literature review	73
Materials and method	75
Results and discussion	81
CHAPTER 8 GENERAL DISCUSSION	88
LITERATURE CITED	93
APPENDICES	101



LIST OF TABLES

<u>Table no.</u>	<u>Description</u>	<u>Page</u>
1	The number of first and second instar nymphs in small, medium and large colonies of <u>Cataenococcus hispidus</u>	19
2	Area (mm^2) of colony and actual number of nymphs in two day and four day old colonies	22
3	Observed and estimated number of nymphs in first instar colonies of <u>C. hispidus</u> . Estimate was based on area of colony (mm^2)	23
4	Mean number of adult female <u>C. hispidus</u> per plot (3 trees) when mealybugs were introduced at various intervals (weeks) before the ant	33
5	Natural enemies observed on the pods in treatments where mealybugs were exposed to natural enemies and to both natural enemies and rainfall	49
6	Mean distance travelled (mm) in five minutes by starved crawlers	55
7	Response of crawlers to light	60
8	Mean number of crawlers found in each site of arena when fresh peduncle was placed between them and source of light	61
9	Geotactic response of crawlers	63
10	Number of nymphs establishing on artificially scarified and unscarified pod surface. Two hundred adult females were used for introduction	66
11	Mean number of live crawlers at various intervals of time when kept at 55% and 90% relative humidity without food. Ambient temperature = $26 \pm 2^\circ \text{C}$. Initial numbers of crawlers per replicate = 10	70
12	Mean number of live crawlers when kept at 15, 20, 30 and 35°C without food.	



	Relative humidity = $95 \pm 5\%$. Initial number of crawlers per replicate = 10	71
13	Composition of <u>D. thoracicus</u> in one-month old artificial nests	82
14	Mean number of trees that were initially not used for introduction colonised by ants	87

LIST OF FIGURES

<u>Figure no.</u>	<u>Description</u>	<u>Page</u>
1	Coverslip used for measuring area of mealybug colony	17
2	Leaf nest used for collecting <u>D. thoracicus</u>	27
3	The establishment of <u>D. thoracicus</u> when mealybugs were introduced before, at the same time and after ants	29
4	Cages used to exclude rain (A) and rain and natural enemies (B)	37
5	Mean number of nymphs on new pod when protected from rain and natural enemies, rain only, natural enemies only and exposed to both	39
6	Mean number of nymphs on old pod when protected from rain and natural enemies, rain only, natural enemies only and exposed to both	41
7	Mean number of adults on old pod when protected from rain and natural enemies, rain only, natural enemies only and exposed to both	47
8	Amount of rainfall during the duration of the experiment on the effect of rainfall and natural enemies on establishment of mealybugs	43
9	Arena used to test response of first instar crawlers to light	57
10	Layout of field when ants and mealybugs were introduced at a density of 50% of the trees	76
11	Layout of field when ants and mealybugs were introduced at a density of 13% of the trees	77
12	Plywood tray used to hold mealybug infested pods	79
13	Establishment of the ant <u>D. thoracicus</u> and mealybugs <u>C. hispidus</u> at various percentage of trees initially used for introduction	83

LIST OF PLATES

<u>Plate no.</u>	<u>Description</u>	<u>Page</u>
I	Mealybugs protected from rain and natural enemies. There is a good colonization of nymphs on the new pod and adult females were still present on the old pod. Photograph taken 12 DAI	44
II	Mealybugs protected from rain but exposed to natural enemies. Virtually no nymphs had established on the new pod but adult females were still present on the old pod. Photograph taken 12 DAI	45
III	Mealybugs exposed to rain and natural enemies. There were virtually no nymphs on the new pod and no adult females on the old pod. Photograph taken 12 DAI	46
IV	Pod artificially scarified before mealybug introduction. Note the good colonization of nymphs on the artificially scarified surface (left half) and virtually no colonization on the unscarified surface (right half).	67
V	Artificial ant's nests in individual newspaper bags loaded onto a truck for transportation	80

An abstract of the thesis presented to the Senate of Universiti
Pertanian Malaysia in partial fulfilment of the requirement for
the degree of Master of Agricultural Science

THE COCOA BLACK ANT-MEALYBUG RELATIONSHIP:
ARTIFICIAL ESTABLISHMENT OF CATAENOCOCCUS HISPIDUS
(HOMOPTERA: PSEUDOCOCCIDAE) ON COCOA

by

Ang Ban Na

June, 1988

Supervisor : Associate Professor Dr. Khoo Khay Chong

Co-supervisor: Puan Rita Muhamad

Faculty : Agriculture

The cocoa black ant Dolichoderus thoracicus (Smith), plays an important role in the control of the cocoa mirid Helopeltis theobromae (Miller) in Malaysia. The ant has a mutualistic relationship with a variety of homopterans, the commonest one being the mealybug Cataenococcus hispidus (Morrison). Attempts to manipulate D. thoracicus are dependent on the ability to manipulate C. hispidus. The study carried out was aimed at obtaining information on C. hispidus to enable the mealybug to be established more successfully.



A method of estimating the number of individuals in a first instar colony was devised. The relationship between the number of individuals (Y) and the area of the colony (X) in mm² is described by the linear equation: $Y=8.55X - 2.98$. The area of colony was measured by using a transparent glass grid.

When using adult female C. hispidus for introduction, it was shown that the optimum time to do so was three days before the ants. It was also found that when the ants were established without the mealybugs, establishment of the former was significantly lower.

Rainfall and natural enemies, mainly coccinellids, neuropterans and slugs, are important mortality factors that seriously hamper establishment of mealybugs. Exposed to rain and natural enemies, the number of nymphs that eventually established on cocoa pods was only four percent of those which received protection from both factors.

The first instar crawlers are the important dispersing stage, in Homoptera. Therefore, studies were conducted on crawler behaviour of C. hispidus. The crawlers show a positive phototactic response and negative geotactic response. There was a strong preference by the crawlers to establish on the artificially scarified surface of pod. Mortality of unfed crawlers was higher at 55±10% relative humidity than at 95±5% relative humidity. When unfed crawlers were kept at 15, 20, 30 and 35°C, survival was highest at 20°C.

Ants and mealybugs were introduced onto 100, 50 and 13% of the trees in a cocoa area. It was found that higher introduction densities resulted in more trees being colonised. But, in terms of number of colonised trees per initially infested tree, the lower densities were more efficient.

Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia sebagai memenuhi sebahagian daripada syarat-syarat untuk Ijazah Master Sains Pertanian.

PERHUBUNGAN SEMUT HITAM KOKO-KOYA: PERTAPAKAN BUATAN
CATAENOCOCCUS HISPIDUS (HOMOPTERA: PSEUDOCOCCIDAE)
DALAM TANAMAN KOKO

oleh

Ang Ban Na

June, 1988

Penyelia : Profesor Madya Dr. Khoo Khay Chong
Penyelia Bersama : Puan Rita Muhamad
Fakulti : Pertanian

Semut hitam koko. Dolichoderus thoracicus (Smith), memainkan peranan penting di dalam mengawal kepinding nyamuk, Helopeltis theobromae (Miller) koko di Malaysia. Semut ini mempunyai perhubungan mutualisma dengan beberapa jenis homopteran, di antara mana yang paling kerap ialah koya Cataenococcus hispidus (Morrison). Usaha-usaha manipulasi ke atas semut adalah bergantung kepada kebolehan usaha-usaha manipulasi terhadap koya. Kajian ini dijalankan dengan tujuan untuk memperolehi maklumat-maklumat berkenaan C. hispidus agar kegiatan penempatan koya ini boleh diusahakan dengan berjaya.

Satu kaedah untuk menganggarkan bilangan individu di dalam koloni instar pertama telah dirangkakan. Perhubungan diantara bilangan individu (Y) dan luas kawasan koloni (X) dalam mm² adalah dijelaskan melalui persamaan linear: $Y=8.55X - 2.98$. Luas kawasan koloni telah diukur dengan gelas lutsinar bergrid.

Apabila menggunakan dewasa betina C. hispidus untuk memulakan infestasi, telah didapati bahawa masa optimum untuk perlakuan sedemikian supaya pertapakan koya yang maksima diperolehi, ialah tiga hari sebelum semut dimasukkan. Juga telah didapati bahawa apabila semut-semut itu bertapak dengan ketiadaan koya maka adalah didapati pertapakan semut-semut ini berkurangan dengan bererti.

Hujan dan musuh-musuh semulajadi, terutamanya coccinellid, neuroptera and lintah bulan adalah faktor-faktor mortaliti penting yang mana sungguh-sungguh menghalang pertapakan koya. Apabila terdedah kepada hujan dan musuh-musuh semulajadi, bilangan nimfa yang akhirnya bertapak pada lenggai-lenggai buah koko hanya lah empat peratus daripada yang mendapat perlindungan daripada kedua-dua faktor tersebut.

Peringkat instar pertama merupakan peringkat yang utama untuk persebaran bagi Homoptera. Oleh itu, kajian dijalankan terhadap tabiat nimfa instar pertama koya. Mobiliti nimfa ini telah didapati berkurang dengan bererti setelah didedahkan kepada kebuluran. Nimfa instar pertama menunjukkan gerak balas fototaksis positif dan gerak balas geotaksis negatif. Telah

didapati nimfa-nimfa amat tertarik untuk bertapak di permukaan parut buatan atas lenggai. Bila disimpan pada kelembapan udara $55 \pm 10\%$ dan $95 \pm 5\%$ mortaliti nimfa yang kebuluran lebih tinggi pada kelembapan udara 55%. Bila nimfa yang kebuluran tersimpan pada 15, 20, 30 dan 35°C , kemandirian adalah paling tinggi pada 20°C .

Apabila semut dan koya dimasukkan ke 100, 50 dan 13% daripada pokok dalam satu kawasan koko, didapati bahawa kepadatan penempatan yang tinggi menghasilkan lebih banyak pertapakan pada pokok-pokok. Tetapi dari segi bilangan pokok yang mempunyai pertapakan semut dan koya per bilangan pokok asal yang digunakan untuk penempatan, penempatan pada kepadatan lebih rendah adalah lebih efisien.

CHAPTER 1

INTRODUCTION

Clausen (1940) noted that ants were the first group of insects to be utilised in an attempt to control insect pest by the biological method. He cited two uses of ants in tropical tree crops by date palm cultivators in Yemen and by citrus growers in southern China. Ants are a group of predators whose predatory values have not been fully investigated (Room, 1975). It has been suggested that where the fauna associated with ants contain important pest species, a reduction in pest damage by manipulation of the dominant ants become possible (Jutsum et al., 1981).

The role of the cocoa black ant, Dolichoderus bituberculatus (Mayr.) in protecting cocoa trees from Helopeltis antonii (Sign.) and H. theivora (Waterh.) in Indonesia have been observed since 1904 (Levert, 1940). A similar relationship in Malaysia was reported for D. bituberculatus against H. theivora theobromae (Miller) (Collingwood, 1977; Azhar, 1984). Some estates in Indonesia practice biological control of Helopeltis soon after beneficial effects of the cocoa black ants were realised. One estate in particular, the Siloewak Sawangan Estate effectively controlled Helopeltis for over 40 years, in a 1500 ha plantation with the cocoa black ant (Geisberger, 1983). Khoo (1988) pointed

out that D. bituberculatus reported in Indonesia and Malaysia is probably D. thoracicus (Smith), because collections of D. bituberculatus in the British Museum did not originate from these two countries. But in this thesis the name D. bituberculatus used by authors before Khoo (1988) would be retained.

In Indonesia D. bituberculatus is associated with the mealybug Planococcus lilacinus (Ckll.) which provides the ant its main food source, the honeydew (Roepke, 1916; Goot, 1917). D. thoracicus in Teluk Intan, Malaysia is associated with the mealybug Cataenococcus hispidus (Morrison), besides other homopterans like scales and aphids. Room (1973) stressed that in order to maintain the numerical superiority of an introduced ant species, its rate of worker production must be kept as high as possible. Therefore it must be ensured that the supply of food is adequate. In most cases it would probably be necessary to introduce the Homoptera tended by the beneficial ants.

It was found that dispersal of P. lilacinus in the field was not effectuated by the cocoa black ant as believed by cocoa planters (Roepke, 1916; Goot, 1917). Dispersal was mainly a function of the mealybugs own movements and wind. Therefore natural dispersal was found to be too slow to achieve a good distribution quickly. Geisberger (1983) concluded that the mealybugs would have to be introduced or a cocoa black ant introduction will fail.

Success in establishment of an introduced Homoptera is dependent upon many factors. It has been found that presence of

attendant ants of the Homoptera enhances establishment (Goot, 1917; Cornwell, 1957). Success of an introduction is also dependent upon timing of introduction with regards to growth stage of the host, e.g. flushing or fruiting (Room, 1973). In homopterans the mobile nymphs which are the principle dispersive stage faces many hazards before they settle to feed. For example, in the scale Aulacaspis tegalensis (Zhnt.) it was found that extreme temperature, low humidity, rain and lack of suitable feeding sites were the major mortality factor for dispersing crawlers (Greathead, 1972). Propensity of crawlers to commence feeding may also be influenced by light (Moran and Cobby, 1979; Gilreath and Smith, 1987). In the cochineal insect, Dactylopius austrinus (de Lotto) propensity of crawlers to settle was enhanced through subjecting them to an artificial "flight" (Moran et al., 1982).

Therefore before the cocoa black ant-mealybug relationship can be used for biological control of Helopeltis a manipulation technique for both the ant, D. thoracicus and mealybug, C. hispidus needs to be developed. Considering the importance of mealybugs to the establishment of the ants, and the many factors that are known to influence establishment of homopterans: this thesis focuses on the manipulation of the mealybugs with the aim of increasing its establishment when introduced.

Studies were undertaken with the following objectives:

1. To develop a method for estimation of number of individuals in mealybug colony.
2. To determine the optimum time to introduce the mealybugs

with respect to that of the ants.

3. Determine the effect of rainfall and natural enemies on the establishment of the mealybugs in the absence of the ants.
4. To study some aspects of crawler behaviour
5. To determine the influence of density of trees in a plot used for introduction of the ants and mealybugs, on the success of the ants and mealybugs in colonising the plot.

CHAPTER 2

LITERATURE REVIEW

This literature review gives a general introduction to the biology of mealybugs, ant-homoptera mutualism and methods that have been employed to artificially establish homopterans. More literature relevant to each experiment are given in the respective sections.

Biology of mealybugs

Mealybugs belong to the family Pseudococcidae of the order Homoptera. The adult females are wingless, more or less degenerate, and are covered with a waxy or powdery coating. The males when present are very much smaller than the females and they do not possess any functional mouth part. Being rather tiny and short-lived, the males are not often observed and the forms that are commonly encountered are the immature stages and adult females. There are four instars in the females and five in the males (Khoo, 1974; Williams, 1985). Mealybugs may be oviparous, viviparous or ovoviviparous (Williams, 1985). In some species the males are totally absent and various types of parthenogenesis occur along with functional hemaphroditism (McKenzie, 1967). The number of eggs or nymphs that can be produced by one female varies according to mealybug species (McKenzie, 1967). As many as

908 nymphs have been produced per female Dysmicoccus brevipes (Ck11.) (Ito, 1938) and 160-299 eggs per female P. lilacinus (Khoo, 1974). The active newly-hatched or born nymphs are the dispersive stage, being able to crawl from plant to plant or be carried by wind or ants to other plants (Khoo, 1974). C. hispidus is viviparous and its biology has been studied by Ang (1988).

ANT ASSOCIATION

Ants as a group are well known mutualists with honeydew-producing Homoptera (Nixon, 1951; Way, 1963; Carroll and Janzen, 1973). In Pseudococcidae the degree of dependence on the ants may vary from strong and almost necessary associations to weak casual seasonal relationships (McKenzie, 1967).

Benefits to ants

Honeydew as source of food. The obvious benefit the ants derive from the homopterans is the acquisition of honeydew as a rich and easily accessible food (Nixon, 1951). In nutritional value, honeydew is more complete than might be expected (McKenzie, 1967). Way (1963) compiled evidence that honeydew may contain sugars, free amino acids and amides, proteins, minerals and B-vitamins. Not all ants forage for honeydew but those that do can at times be the most numerous (Samways, et al., 1982). An example of this was observed by Grant and Moran (1986) in undisturbed woodland savanna, where the dominant ants were almost exclusively honeydew-dependent. It has been suggested that