

## **UNIVERSITI PUTRA MALAYSIA**

# TAXONOMIC AND ELECTROPHORETIC STUDIES OF SELECTED SPECIES OF GANODERMA (KARST.)

**FARIDAH ABDULLAH** 

FP 1996 15



## TAXONOMIC AND ELECTROPHORETIC STUDIES OF SELECTED SPECIES OF GANODERMA (KARST.)

## By

## **FARIDAH ABDULLAH**

Dissertation Submitted in Fulfilment of the Requirements for the Degree of Doctor of Philosophy in the Faculty of Agriculture, Universiti Pertanian Malaysia.

October, 1996



#### **ACKNOWLEDGEMENTS**

I would like to thank Assoc. Prof. Dr. Lim Tong Kwee in his capacity as chairman of the supervisory committee, from January 1986 to June 1990; and for having introduced me to a subject that caught my interest and which shall be my area of research for a long time to come. Thanks are also extended to Prof. Dr. Rosli Mohamad who had taken over the chairmanship when Dr. Lim left the University in 1990 and also to Assoc. Prof. Dr. Kamaruzaman Sijam in his capacity as a member of the Supervisory Committee. Their comments and contributions had been very helpful towards the completion of the thesis. I would also like to extend my appreciation to the Plant Pathology laboratory staff at the Faculty of Agriculture who had rendered help both in the laboratory and in the field. Similarly, I would like to record my gratitude to the Head, Department of Biology for having allowed the use of the Department's laboratories during my studentship tenure and to colleagues in the Department who had rendered help in whichever way they can.

My heartfelt appreciations are particularly extended to Dr. Gurmit Singh from United Plantations in Teluk Intan, Dr. Ismail HamzahandEn. Khairuddin Hashim, bothof Golden Hope Plantations in Selangor, Mr. Bauer from Boh Plantations, and staff from Lian



Hup Estate, Johore, for providing the samples for my studies. Due thanks are also extended to the Building Research Establishment, England and to Prof. Wright from Argentina, for having made available samples from their respective regions. I would also like to thank Dr. Pegler for giving me the opportunity to browse through *Ganoderma* specimens at The Herbarium, during my visit to Kew, England.

A word of thanks is especially recorded to Dr. Peter Omundsen from the University of California (UCLA), Riverside; who has kindly helped with the numerical analysis of electrophoretic results. I am also indebted to a few friends and occasional visitors to UPM, who have helped me with translations of texts in Dutch, French, Japanese, Chinese and Latin into English. Without their help, much information on the subject may have still remained inaccessible.

I would like to thank Universiti Pertanian Malaysia for having granted me study leave and to Jabatan Perkhidmatan Awam (JPA) for having awarded a scholarship throughout my studentship. Last but not least I would like to extend my heartfelt gratitude to friends and members of my family, particularly my children, who had inspired the staying power when the going gets tough. May God bless all of you.

## TABLE OF CONTENTS

|                               |                     | page  |
|-------------------------------|---------------------|-------|
| ACKNOWLEDGEMENTS              |                     | ii    |
| LIST OF TABLES                |                     | xii   |
| LIST OF FIGURES               |                     | xiv   |
| LIST OF PLATES                |                     | xvi   |
| LIST OF ABBREVIATIONS         |                     | xxiii |
| ABSTRACT                      |                     | xxiv  |
| ABSTRAK                       |                     | xxvii |
| CHAPTER                       |                     |       |
| I INTRODUCTION                |                     | 1     |
| II LITERATURE REVIE           | EW                  | 8     |
|                               | ustry: A Historical | 8     |
| Pathological Asp              | pects               | 10    |
| Basal Stem Rot of in Malaysia | of Oil Palm         | 12    |
| Pathogenicity Te              | ests                | 18    |
| Disease Control               |                     | 20    |
| Ganoderma On (                | Other Palms         | 22    |



|     | Taxonomy of the Genus Ganoderma           | 23 |
|-----|---|----|
|     | Growth Studies                            | 30 |
|     | Effect of Nutritional Factors on          |    |
|     | Growth                                    | 30 |
|     | Vitamins in Fungal Nutrition              | 32 |
|     | Effect of Temperature on Growth           | 32 |
|     | Other Factors Affecting Growth            | 33 |
|     | Cross-Plating for Compatability Tests     | 35 |
|     | Electrophoretic Studies                   | 35 |
| III | MATERIALS AND METHODS                     | 40 |
|     | A Survey of <i>Ganoderma</i> Incidence In |    |
|     | Oil Palm and Coconut Plantations          | 40 |
|     | Sporophore Collections                    | 42 |
|     | Diversity of Ganoderma From Oil           |    |
|     | Palms and Coconut Stumps                  | 45 |
|     | Diversity Based on Sporophore             |    |
|     | Morphology                                | 45 |
|     | Diversity Based on Mycelial Types         | 47 |
|     | Sporophore Induction of Ganoderma         |    |
|     | on Solid Subtrates                        | 48 |



| Substrate Preparation                   | 49 |
|---|----|
| Inoculation of Isolates on Solid        |    |
| Substrates                              | 49 |
| Growth and Development on Solid         |    |
| Substrates                              | 51 |
| Colour Reproduction of Induced          |    |
| Sporophores                             | 53 |
| Effect of Interrupted Humidity During   |    |
| Ganoderma Sporulation                   | 53 |
| Groups I and II: Comparative Sporophore |    |
| Morphology                              | 55 |
| Growth Studies of Groups I and II       | 59 |
| Response to A Range of                  |    |
| Temperatures                            | 59 |
| Response to Agar Media                  | 60 |
| Ganoderma Incidence In Rubber, Tea and  |    |
| Cocoa Plantations                       | 61 |
| Sampling of Ganoderma from              |    |
| Woody Hosts                             | 61 |
| Ganoderma from Temperate Regions        | 62 |
| Growth Studies of Groups III and IV     | 65 |
| Response to Temperature                 | 65 |
| Somatic Compatibility Tests of Group    |    |
| III Isolates                            | 65 |



|    | Groups III and IV: Sporophore Induction        |    |
|----|--|----|
|    | and the Effect of Supplements in the Substrate | 66 |
|    | I  |    |
|    | Intraspecific and Interspecific Diversity of   |    |
|    | Ganoderma Based on Isozyme                     | 60 |
|    | Electrophoresis                                | 68 |
|    | Numerical Analysis                             | 68 |
|    | Horizontal Polyacrylamide Gel                  |    |
|    | Electrophoresis (PAGE)                         | 69 |
|    | Fungal Cultures and Enzyme                     |    |
|    | Extraction                                     | 70 |
|    | Scoring Isozymic Bands                         | 71 |
|    | Specific Diversity from Dendrogram             | 74 |
| IV | RESULTS  | 75 |
|    | Ganoderma Incidence in Oil Palm                |    |
|    | and Coconut Plantations                        | 75 |
|    | Foliar signs                                   | 75 |
|    | Presence of Sporophores                        | 76 |
|    | Symptoms on Trunk                              | 80 |
|    | Symptoms on Roots                              | 80 |
|    | Colour Types (CT's) and Diversity of           |    |
|    | Ganoderma Sporophores From Oil                 |    |
|    | Palms and Coconut                              | 81 |
|    |  |    |



| C1 1: Chestnut Brown  | 81  |
|---|-----|
| CT 2 : Orange-Reddish Brown   | 84  |
| CT 3 : Yellow-Buff  | 85  |
| CT 4 : Purplish-Black   | 86  |
| CT 5 : Deep Brown   | 88  |
| CT 6: Ochrous Brown   | 89  |
| CT 7: Blackish-Brown  | 89  |
| Mycelial Types (MT's) of <i>Ganoderma</i> From Oil Palms and Coconut Stumps | 91  |
| Characteristics of MT 1   | 90  |
| Characteristics of MT 2   | 93  |
| Characteristics of MT 3   | 95  |
| Characteristics of MT 4   | 96  |
| Relationship of Mycelial Type to Sporophore Colour                          | 97  |
| Sporophore Induction of Isolates from Oil Palms                             | 99  |
| Spore Observations  | 104 |
| Colour Reproduction of Induced Sporophores                                  | 107 |
| Induced Sporophores from EGr Series   | 110 |



| Effect of Reduced Humidity on Ganoderma                  |     |
|--|-----|
| Sporulation  | 110 |
| Groups I and II: Comparative Sporophore                  |     |
| Morphology   | 111 |
| Sporophore Attachment                                    | 113 |
| Degree of 'Lacca' Deposition                             | 114 |
| Topical Morphology of the Pileus                         | 115 |
| Margin Characteristics                                   | 116 |
| Pore Tubes and Context Layer                             | 118 |
| Literature Reference                                     | 120 |
| Colour Morphology  | 121 |
| Sessile/Stipitate Sporophore                             | 123 |
| Pileus Morphology  | 123 |
| Sporophore Anatomy                                       | 123 |
| Peripheral Margin  | 124 |
| Plant-Host Specificity                                   | 124 |
| Groups I and II: Response to a Range of                  |     |
| Temperatures   | 126 |
| Groups I and II: Response to a Range of                  |     |
| Agar Media   | 130 |
| Ganoderma Incidence In Rubber, Tea and Cocoa Plantations | 133 |
| Diversity of <i>Ganoderma</i> from Woody                 | 127 |



|   | Sporophores of <i>Ganoderma</i> from Woody Hosts                       | 1: |
|---|--|----|
|   | Mycelial Types of Ganoderma from                                       | 4  |
|   | Woody Hosts  Mycelial Types of <i>Ganoderma</i> From Temperate Regions | 1: |
|   | Groups III and IV: Response to Temperature                             | ]  |
|   | Somatic Compatibility of Group III Isolates                            | 1  |
|   | Groups III and IV: Sporophore Induction and the                        | 1  |
|   | Effect of Supplemented Substrates                                      | 1  |
|   | Horizontal Polyacrylamide Gel Electrophoresis                          |    |
|   | (PAGE)   | 1  |
|   | Isozyme Staining   |    |
|   | Diversity of Isolates Based on Isozymic Bands                          |    |
|   | Diversity Based on Numerical Analysis                                  | 1  |
|   | Specific Diversity from Dendrogram                                     |    |
| V | DISCUSSION   | 1  |
|   | Ganoderma Incidence in Plantations                                     | 1  |
|   | Ganoderma on Stumps  |    |



| Sporophore Induction  | 173 |
|---|-----|
| Characterisation by Sporophore  Morphology                        | 174 |
| Characterisation from Colony Cultures                             | 176 |
| Characterisation by Isozyme Electrophoresis                       | 177 |
| Relationship of <i>Ganoderma</i> from Coconut Stumps to Oil Palms | 181 |
| Ganoderma on Woody Hosts  | 181 |
| VI CONCLUSION   | 183 |
| REFERENCES  | 187 |
| APPENDICES  | 197 |
| BIOGRAPHICAL SKETCH   | 209 |



## LIST OF TABLES

| <b>Fable</b> |  | page |
|--------------|--|------|
| 1            | Group I Isolates from Oil Palm : Collection Sites, Codes and Numbers                         | 43   |
| 2            | Group II Isolates from Coconut Stumps: Collection Sites, Codes and Numbers                   | 44   |
| 3            | Isolates from Group I Used in Sporophore Induction   | 50   |
| 4            | Specimens Used in Comparative  Morphological Characterisations                               | 56   |
| 5            | Isolates From Group II Used in Growth Studies  | 56   |
| 6            | Group III Isolates from Perennial Woody Crops: Collection Sites, Codes and Numbers           | 63   |
| 7            | Group IV Isolates From Woody Hosts of Temperate Regions: Collection Sites, Codes and Numbers | 64   |
| 8            | Isozymes Used in Electrophoresis   | 72   |



| 9  | Diversity of Ganoderma Based on MT's  | 93  |
|----|---|-----|
| 10 | MT 1 Isolates Enlisted According to Their CT Subgroups  | 94  |
| 11 | Development of <i>Ganoderma</i> Sporophore Cultures on Palm-Press Fibres vs. Rubber Sawdust     | 100 |
| 12 | Effect of Temperature on Mycelial Growth of <i>Ganoderma</i> From Oil Palms (Group I)           | 128 |
| 13 | Effect of Temperature on Mycelial Growth of <i>Ganoderma</i> From Coconut Stumps (Group II)     | 129 |
| 14 | Effect of Different Media on Mycelial Growth of <i>Ganoderma</i> From Oil Palms (Group I)       | 138 |
| 15 | Effect of Different Media on Mycelial Growth of <i>Ganoderma</i> From Coconut Stumps (Group II) | 139 |
| 16 | Effect of Temperature on Mycelial Growth of <i>Ganoderma</i> Isolates From Woody, Hosts         | 147 |



## LIST OF FIGURES

| Figure |   | page |
|--------|---|------|
| 1      | Ganoderma Taxonomy as Proposed by Karsten, 1881 (Pegler and Young, 1973)                    | 24   |
| 2      | Ganoderma Taxonomy as Proposed by Patouillard, 1889 (Pegler and Young, 1973)                | 25   |
| 3      | Ganoderma Taxonomy as Proposed by Imazeki,1939 (Pegler and Young, 1973) and Steyaert (1980) | 27   |
| 4      | Sites in West Malaysia Where <i>Ganoderma</i> Sporophores Were Collected                    | 41   |
| 5      | Diagram of Parts of A Sporophore and Their Mycological Terms                                | 46   |
| 6      | A Schematic Diagram of the Sporophore Pileus Showing (i) Undulations and (ii) Corrugations  | 58   |
| 7      | Esterase A Banding Patterns in  Ganoderma  xiv  | 156  |



| 8  | Esterase D Banding Patterns in Ganoderma   | 159 |
|----|--|-----|
| 9  | Cholinesterase Banding Patterns in Ganoderma   | 161 |
| 10 | Acid Phosphatase Banding Patterns in Ganoderma   | 162 |
| 11 | Malate Dehydrogenase Banding Patterns in <i>Ganoderma</i>  | 164 |
| 12 | A UPGMA Cluster Analysis of <i>Ganoderma</i> Based on Modified Roger's Distance Measure Calculated From The Isozyme Data | 170 |
| 13 | A Simplified Version of the Dendogram Depicting the Main Clusters  | 171 |



## LIST OF PLATES

| Plate |   | page |
|-------|---|------|
|       |   |      |
| 1     | Ganoderma Sporophores Arising From      |      |
|       | Exposed Roots of the Oil Palm,          |      |
|       | E. guineensis                           | 76   |
| 2     | Ganoderma Sporophores On A Coconut      |      |
|       | Trunk                                   | 77   |
| 3     | Ganoderma Sporophores Collected         |      |
|       | from Oil Palm Trunks                    | 78   |
| 4     | An Imbricate Ganoderma Sporophore       | 79   |
| 5     | A Sessile Sporophore from Coconut Stump | 79   |
| 6     | EGUB 01 Sporophore Representing         |      |
|       | CT 1 With Its Gradation of Chestnut     |      |
|       | Brown Colour                            | 83   |
| 7     | EGB 009 Sporophore Representing         |      |
|       | CT 2 With Its Orange-Reddish Pileus     |      |
|       | and Conspicuous Growth Rings            | 85   |



| 8  | EGPN 307 Representing CT 3 With Its Sub  |    |
|----|--|----|
|    | laccate Condition and Yellow-Buff Colour | 86 |
| 9  | EGB 005 Sporophore Representing CT 3,    |    |
|    | Showing the Same Shade of Purplish-Black |    |
|    | Throughout                               | 87 |
| 10 | EG 114 Sporophore Representing CT 5      | 88 |
| 11 | Induced Sporophore of CNSB 02            |    |
|    | Representing Sporophores In CT 6         | 90 |
| 12 | Induced Sporophore of CNUB 01            |    |
|    | Representing CT 7                        | 90 |
| 13 | EGB 001 on MA at 28°C, Representing MT 1 |    |
|    | With Its Dense and Uniform Mycelia       | 92 |
| 14 | Culture Colony of EGP 111 on MA at 28°C, |    |
|    | Representing MT 2 Showing Patchy Mycelia | 93 |
| 15 | EGSP 05 on MA 28°C, Representing MT 3    |    |
|    | Showing (a) Young White Mycelia          |    |
|    | and(b)Aged Yellow Mycelia                | 96 |
| 16 | CNUB 01 On MA at 28°C,                   |    |
|    | Representing MT 4                        | 97 |



| 17 | Mycelial Establishment on (a) Palm-Press  |     |
|----|---|-----|
|    | Fibres and (b) Rubberwood Sawdust         |     |
|    | Substrates                                | 100 |
| 18 | Transformation of the Prin ordium Into    |     |
|    | a Slender Stipe Within 7 Days' Growth.    |     |
|    | Inset: Close-up View                      | 101 |
| 19 | Prolific Liberation of Spores from        |     |
|    | Induced Sporophores                       | 102 |
| 20 | Ganoderma Spores Showing Vacoules in the  |     |
|    | Cytoplasm and the Two-Layered Cellwall,   |     |
|    | with the Inner Layer Having Ornamented    |     |
|    | Outgrowth (Magnification:1000x)           | 104 |
| 21 | EG 114 Spores with Suprahilar             |     |
|    | Plage (arrowed)                           | 105 |
| 22 | EG 114 Spores in Lactophenol              | 106 |
| 23 | Freshly-Liberated Spores of Ganoderma     |     |
|    | Showing Germination Tubes                 | 106 |
| 24 | Field Collected EGB 005 and Its Induced   |     |
|    | Form (inset)                              | 108 |
| 25 | The Yellow-Orange Line in CT 2 Sporophore |     |
|    | During Its Early Growth Stages            | 109 |





| 26 | (a) Field-Collected EGB 009, and (b) Its Induced Form                                     | 109 |
|----|---|-----|
|    | (0) 200 200 400 1 0000000000000000000000000   | 100 |
| 27 | Freshly-Collected <i>Ganoderma</i> Sporophores From Coconut Stumps                        | 112 |
|    | <b>T</b>  |     |
| 28 | The Same Group of Sporophores From Plate 27, After They Were Cleaned                      | 113 |
|    |   |     |
| 29 | Ganoderma Sporophore from Coconut Stump With Its Distinct, Raised Basal Portion (arrowed) |     |
|    | In the Absence of a Stipe   | 114 |
| 30 | Comparison of Laccate Surface;  |     |
|    | (a) G. applanatum (Argentina), (b) Two Local Specimens from Coconut Stumps,               |     |
|    | (c) A Sporophore from Oil Palm and (d) G. applanatum (Chile)                              | 115 |
|    |   |     |
| 31 | Anatomical Section of <i>Ganoderma</i> from Oil Palm Showing A                            |     |
|    | Flabelliform Margin   | 117 |
| 32 | Anatomical Section of Ganoderma   |     |
|    | from Coconut Stump Showing A Thin, Incurved Margin  | 117 |
| 33 | Ganoderma from (A) Coconut Stump  |     |
|    | and (B) Oil Palm Before Dissection  | 119 |



| 34 | Sporophores from (A) Oil Palm Compared to (B) Coconut Stump After Dissection. Note (a) margin shape, (b) pore tubes, and (c) contextuatissue containing melanoid vs colourless cellular streaks (arrowed) | ıl<br>119 |
|----|---|-----------|
| 35 | Specimen from Oil Palm With Its Sulcate Appearance, Clear Concentric Rings and Short Stipe  | 120       |
| 36 | Specimen from Plate 35 After Dissection Showing the Thin, Curved Margin, Atypical of Specimens from Oil Palm  | 121       |
| 37 | Growth Response of EGP III at 15° C to 40°C On the Eighth Day of Growth   | 126       |
| 38 | Growth Response of CNG 05 at 15°C to 40°C On the Eighth Day of Growth   | 126       |
| 39 | G. pseudoferreum syn. G. philippi on Hevea Stump  | 132       |
| 40 | Tea bushes Showing Foliar Browning, Which Were Easily Spotted Out as Patches  | 133       |
| 41 | A Dead Tea Bush From Whose Roots  Ganoderma was Isolated  | 133       |



| 42 | Infected <i>T. cacao</i> with Advanced Stage |      |
|----|--|------|
|    | Foliar Symptoms                              | 135  |
| 43 | Infected Cocoa Root Showing (a) the White    |      |
|    | Mycelia and (b) Reddish 'Crusts' (arrowed)   |      |
|    | formed on the Lesions                        | 135  |
| 44 | Sporophore of G. pseudoferreum syn. G. phili | ippi |
|    | Showing the Dull, Non-Laccate, Brownish-Blac | k    |
|    | Upper Surface and the White Margin           | 138  |
| 45 | Undersurface of G. pseudoferreum,            |      |
|    | Which is Off-White and Covered               |      |
|    | With Minute Pores                            | 138  |
| 46 | Culture Colony of <i>G. pseudoferreum</i>    |      |
|    | on (a) MA Plate and in (b) Liquid            |      |
|    | Flask Culture at 28°C                        | 140  |
| 47 | Colony Culture of The Medicinal Isolate      |      |
|    | GLT 01 (G. lucidum) on MA at 28°C            | 142  |
| 48 | Miniature G. lucidum (GLT 01) Sporophore     |      |
|    | On MA at 28°C. Inset: Earlier Stage of       |      |
|    | Development                                  | 143  |
| 49 | Mergence and Intermingling of EGB 003 vs     |      |
|    | EGB 003:(A) Upper Surface and                |      |
|    | (B) Under Surface                            | 146  |



| 50 | Formation of Linear Boundary Between          |     |
|----|---|-----|
|    | CSBC 01 vs. Gl 409: (A) Upper Surface         |     |
|    | and (B) Under Surface                         | 147 |
| 51 | A Marked Boundary Indicating Somatic          |     |
|    | Incompatability Within the EGB Series: 003 vs |     |
|    | 008: (a) Upper Surface and                    |     |
|    | (b) Under Surface                             | 148 |
| 52 | Rapid Crust Formation by Oil Palm             |     |
|    | Isolates on Nitrogen-Supplemented             |     |
|    | Substrates                                    | 149 |
| 53 | Prolific Growth of G. lucidum (GLT 01)        |     |
|    | Sporophores on Nitrogen-Supplemented          |     |
|    | Substrates                                    | 150 |
| 54 | Sporophore of (a) G. lucidum (GLT 01) and     |     |
|    | (b) G. resinaceum (Gr 144A) From the Control  |     |
|    | Treatment                                     | 151 |
| 55 | Esterase A Bands of EGP 114 Showing           |     |
|    | the 3 Sites of Activity                       | 153 |
| 56 | Esterase A Banding Pattern of Ganoderma       |     |
|    | Isolate from Coconut Stump (left)             |     |
|    | and Oil Palm (right)                          | 155 |
| 57 | Duplicate Malate Dehydrogenase Bands of       |     |
|    | Three Coconut Stumps Isolates (left) Against  |     |
|    | A Single Oil Palm Isolate (right)             | 161 |
|    | xxii  |     |



## LIST OF ABBREVIATIONS

#### **MEDIA**

LBA - Lima Bean Agar

MA - Malt Agar

CMA - Corn Meal Agar

PDA - Potato Dextrose Agar

CDA - Corn Dextrose Agar

WA - Water Agar

#### **SUBSTRATES**

RSD - rubberwood sawdust

PPF - palm-press fibers

## **ENZYMES**

EST - Esterase

ACP - Acid Phosphatase

ChE - Cholinesterase

MDH - Malate Dehydrogenase

#### **MEASUREMENTS**

g - gram

L - litre

m - millilitre

mm - millimeter

nmg - no measurable growth

## **ANALYSIS**

OTU - Operational Taxonomic Unit

NTSYS-pc - Numerical Taxonomy Systematics Computer

Programme

UPGMA - Unweighted Paired Group Matching Analysis

xxiii





Abstract of dissertation submitted to the Senate of Universiti Pertanian Malaysia in

fulfilment of the requirements for the degree of Doctor of Philosophy.

TAXONOMIC AND ELECTROPHORETIC STUDIES

**OF** 

SELECTED SPECIES OF GANODERMA (KARST.)

By

FARIDAH ABDULLAH

October 1996

Chairman: Prof. Rosli bin Mohamad

Faculty: Agriculture

Ganoderma, the causal pathogen of Basal Stem Rot, is found

in association with all major plantation crops in Malaysia, but the

most severe debilitation caused is when they are found on oil palms.

It is not known how many species/pathogenic types there are,

whether a differential degree of virulence existed, and whether the

fungus is host-specific. Ganoderma taxonomy is still controversial

and could not effectively aid pathological considerations. This

xxiv