



UNIVERSITI PUTRA MALAYSIA

**BIOEFFICACY AND INFECTION MECHANISM OF SEVERAL
ENTOMOPATHOGENIC FUNGI ON CROCIDOLOMIA BINOTALIS
ZELL.**

NORZITA HASHIM

FP 2000 4

**BIOEFFICACY AND INFECTION MECHANISM OF SEVERAL
ENTOMOPATHOGENIC FUNGI ON *CROCIDOLOMIA BINOTALIS* ZELL.**

By

NORZITA HASHIM

**Thesis Submitted in Fulfilment of the Requirements for the
Degree of Master of Agricultural Science in the Faculty of Agriculture
Universiti Putra Malaysia**

April 2000



DEDICATION

To

My parents, Hashim bin Abdullah and Tinam binti Jusoh

My brothers and sisters



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Master of Agricultural Science.

BIOEFFICACY AND INFECTION MECHANISM OF SEVERAL ENTOMOPATHOGENIC FUNGI ON *CROCIDOLOMIA BINOTALIS* ZELLER

By

NORZITA BINTI HASHIM

April 2000

Chairman: Associate Professor Dr. Yusof Ibrahim

Faculty: Agriculture

The role of entomogenous fungi as natural enemies for cruciferous insect pests has recently been explored and several isolates of hypomycetous fungi have been identified. The ability of three entomopathogenic isolates of *Beauveria bassiana* (Bals.) Vuill, *Paecilomyces fumosoroseus* (Wize) Brown and Smith and *Metarhizium anisopliae* var. *majus* (Metsch.) Sorokin to cause infection against second instar larvae of *Crocidolomia binotalis* Zeller was tested. Dosage-response assays were conducted in an ambient environment of $28 \pm 2^{\circ}$ dark periods.

treatment. Results indicated that all isolates of the entomopathogenic fungi were able to cause mortality on the second instar larvae of *C. binotalis*. Exposure to sprays of a conidial suspension of varying dosages from 20 ml^{-1} to 2×10^7 larval mortality from 10.0 to 100.0%

conidia ml^{-1} , larval mortality was recorded in excess of 80%. ED_{50}

was $1926 \text{ conidia ml}^{-1}$ and was significantly much lower than that of *B. bassiana* which



was 5038 conidia ml⁻¹. Meanwhile, the ED₅₀ for *M. anisopliae* var. *majus* was 20000 conidia ml⁻¹, which was much higher than that for *P. fumosoroseus* and *B. bassiana*. *P. fumosoroseus* also demonstrated the best potency of infection when all its series of concentration took lesser times to reach 50% infectivity (LT₅₀) and *M. anisopliae* var. *majus*.

The infection mechanisms of all isolates of the entomopathogenic fungi were studied histologically by scanning electron microscopy. In general, conidia which landed on the cuticle germinated within 4 to 6 hours after inoculation. The germ tube penetrated the cuticle and the hypha subsequently invaded the tissue. Infected larvae died between 24 - 48 hours of post inoculation. The hyphae broke through the outer cuticle after 2-5 days of post-treatment. Conidiophore were produced and they formed conidia, one or two days later. At this stage, complete invasion of the larvae had occurred.

Results from laboratory bioassays revealed that *P. fumosoroseus* was the most efficacious of the three fungal species tested. As such, the efficacy of *P. fumosoroseus* was tested in the field. Studies were conducted in the farm at Universiti Putra Malaysia, Serdang, Selangor, Malaysia. Four formulations of *P. fumosoroseus* were made, namely conidia in aqueous Tween 80 (0.05%), conidia in kaolin, conidia in palm oil and conidia from cultures in rice flour. Larval mortality was recorded daily until 12 days to determine the cumulative percentage mortality. Mean percent mortality was in excess of 70% for all the treatments except for the untreated control. Conidia in palm oil was able to inflict the highest larval mortality (88.5%) and there was no significant difference

compared with conidia in aqueous Tween 80 which recorded 78.0%. Conidia from rice flour cultures effected 76.0% mortality and this was shown to be significantly less virulent than the conidia in palm oil. A lower larval mortality of 75.0% was observed when sprayed with conidia in kaolin. It is thus concluded that conidia in palm oil have the potential for an effective field application compared with conidia in kaolin and conidia from cultures in rice flour.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains Pertanian.

BIOEFIKASI DAN MEKANISMA JANGKITAN OLEH BEBERAPA KULAT ENTOMOPATOGEN KE ATAS *CROCIDOLOMIA BINOTALIS* ZELLER

Oleh

NORZITA BINTI HASHIM

April 2000

Pengerusi: Profesor Madya Dr. Yusof Ibrahim.

Fakulti: Pertanian.

Peranan kulat-kulat entomogenus sebagai musuh semulajadi kepada serangga perosak krusifer telah dijelajahi baru-baru ini dan beberapa pencilan kulat-kulat hypomycete telah dikenalpasti. Oleh yang demikian, keupayaan jangkitan tiga pencilan kulat-kulat entomopatogen *Beauveria bassiana* (Bals.) Vuill, *Paecilomyces fumosoroseus* (Wize) Brown and Smith dan *Metarhizium anisopliae* var. *majus* (Metsch.) Sorokin telah diuji ke atas larva instar kedua *Crocidolomia binotalis* Zeller. Asai respon-dos telah dilakukan di dalam keadaan persekitaran terkawal pada suhu $28 \pm 2^\circ$ bandingan $80 \pm 15\%$ dan dengan keadaan berselang-seli terang dan gelap.

larva dicatat 12 hari selepas pendedahan kepada rawatan semburan konidia. Keputusan menunjukkan kesemua pencilan kulat-kulat entomopatogen berkebolehan menyebabkan kematian ke atas larva instar ke dua *C. binotalis*. Semburan dengan kepekatan konidia 20 ml^{-1} hingga 2×10^7

100.0

80%. ED₅₀ bagi *P. fumosoroseus* adalah 1926 konidia ml^{-1} iaitu lebih rendah berbanding *B. bassiana* dengan 5038 konidia ml^{-1} . Sementara itu, ED₅₀



majus adalah 20000 konidia ml⁻¹,

bassiana. *P. fumosoroseus* juga menunjukkan potensi jangkitan yang terbaik apabila kesemua siri kepekatan mengambil masa yang singkat untuk mencapai 50% jangkitan (LT₅₀) berbanding *B. bassiana* dan *M. anisopliae* var. *majus*.

Mekanisma jangkitan kesemua pencilan kulat entomopatogen telah dikaji secara histologi melalui mikroskop elektron. Secara umum, bercambah dalam masa 4 - 6 jam selepas inokulasi. Tiub germa menembusi kutikel dan hifa kemudiannya menyerang tisu. Larva yang diserang mati di selepas penginokulan. Konidiofor terbentuk dan mengeluarkan konidia, sehari atau dua hari kemudiannya. Pada ketika ini, jangkitan sepenuhnya ke atas larva telah berlaku.

Keputusan daripada bioasai di makmal menunjukkan *P. fumosoroseus* adalah paling berkesan daripada ke tiga-tiga kulat yang diuji.

diuji di ladang. Kajian telah dilakukan di ladang Universiti Putra Malaysia, Serdang, Selangor, Malaysia.

dalam air suling mengandungi Tween 80 (0.05

dalam minyak kelapa sawit dan konidia di dalam tepung beras.

dicatit setiap hari sehingga 12 hari untuk mendapatkan peratus kematian longgokan.

Peratus kematian adalah melebihi 70% bagi kesemua rawatan kecuali kawalan.

di dalam minyak kelapa sawit berkebolehan menyebabkan kematian larva paling tinggi (88.5

mengandung Tween 80 yang dicatit 78% kematian.

menyebabkan 76.0% kematian dan ini adalah kevirulenan yang lebih rendah dengan bererti daripada konidia di dalam minyak kelapa sawit. Kematian larva yang paling rendah iaitu 75% diperhatikan apabila disembur dengan konidia di dalam kaolin. Dengan ini diputuskan bahawa konidia di dalam minyak kelapa sawit berpotensi untuk kegunaan di ladang berbanding konidia di dalam kaolin dan konidia daripada kultur di dalam tepung beras.

ACKNOWLEDGEMENTS

First of all, I thank God for giving me strength and ability to complete this study. I also sincerely grateful and greatly indebted to my supervisor Assoc. Prof. Dr. Yusof bin Ibrahim for his wise guidance, understanding, support, compassion and sacrifices. invaluable advice and comments are highly appreciated.

My sincere appreciation and gratitude to members of my Supervisory Committee, Assoc. Prof. Dr Tan Yee How, and Assoc. Prof. Dr. Ahmad Said Sajap for their critical comments and suggestions during the course of this project.

I also wish to thank Mr. Manan Tikon from Entomology Laboratory, all the laboratory staffs of the Pathology Lab, Nematology Lab and staffs of the Graduate School for their kind assistance and cooperation in throughout my study.

Special thanks to Kak Nor Putri for her friendship, encouragement and support throughout my hard times.

My utmost gratitude to my parents, brothers and sisters for their boundless support and love. Finally, I wish to express my sincere thanks to all those who have one way or another helped me in making this study a success.



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL SHEETS	x
DECLARATION FORM	xii
LIST OF TABLES	xv
LIST OF PLATES	xvi
 CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 <i>Crocidolomia binotalis</i> Zeller	4
2.1.1 Introduction	4
2.1.2 Biology and Ecology	5
2.1.3 Damage and Importance	11
2.1.4 Natural Enemies	11
2.1.5 Control	12
2.2 <i>Beauveria bassiana</i> (Bals.)	14
2.2.1 Introduction	14
2.2.2 History	16
2.2.3 Mode of Infection	16
2.3 <i>Metarhizium anisopliae</i> var. <i>majus</i> (Metsch.) Sorokin	17
2.3.1 Introduction	17
2.3.2 History	19
2.3.3 Mode of Infection	20
2.4 <i>Paecilomyces fumosoroseus</i> (Wize) Brown and Smith	21
3 MATERIALS AND METHODS	23
3.1 Insect Rearing	23
3.2 Fungal Isolates	23
3.3 Dosage-Response Assay	24
3.4 Histopathology	26
3.4.1 Inoculation of Entomopathogenic Fungi	26
3.4.2 Larva Collection	26
3.5 Field Trial	29
3.5.1 Formulation of Fungus	29
3.5.2 Efficacy Test	32



4.0	RESULTS AND DISCUSSION	34
4.1	Dosage-Response Assay	34
4.2	Histopathology	42
4.2.1	General Observasion	42
4.2.2	Conidia Germination and Penetration of the Cuticle	45
4.2.3	Invasion of the Host Tissue	55
4.2.4	Development of Conidiophores	57
4.3	Field Trial	65
5.0	CONCLUSION	71
	REFERENCES	73
	APPENDICES	86
	BIODATA OF AUTHOR	89



LIST OF TABLES

Table	Page
1. Mean Percent Mortality of Second Instar Larvae of <i>C. binotalis</i> Upon Exposure to Varying Dosages of <i>P. fumosoroseus</i> , <i>B. bassiana</i> and <i>M. anisopliae</i> var. <i>majus</i> at 12 Days of Post Treatment, and the Corresponding CFU/s/mm ² on Bacteriological Agar	35
2. Effect of Isolates of <i>P. fumosoroseus</i> (Pf), <i>B. bassiana</i> (Bb) and <i>M. anisopliae</i> var. <i>majus</i> (Mam) on Second Instar Larvae of <i>C. binotalis</i>	39
3. Days to Death for Varying Dosages of <i>P. fumosoroseus</i> , <i>B. bassiana</i> and <i>M. anisopliae</i> var. <i>majus</i> on Second Instar Larvae of <i>C. binotalis</i>	40
4. Cumulative Percent Mortality of Second Instar Larvae of <i>C. binotalis</i> After Treatment with Conidia from Cultures on Rice Flour (R), Conidia in Palm Oil (P), Conidia in Aqueous Tween 80 (S), Conidia in Kaolin (K) and Untreated Control (C)	66
5. Cumulative Percent Mortality of Second Instar Larvae of <i>C. binotalis</i> After Treatment with Various Formulations of <i>P. fumosoroseus</i>	67



LIST OF PLATES

Plate	Page
1. The Eggs of <i>C. binotalis</i> are Laid in Batches on the Lower Surfaces of the Leaf	6
2. First Instar Larvae of <i>C. binotalis</i> are Slender and the Body is Greenish Yellow in Colour (about 2mm)	6
3. The Second Instar Larva of <i>C. binotalis</i> (about 4 mm)	7
4. Late Larval Instar is Characterised by Three Dorsal and Two Lateral Longitudinal Stripes of Pale Colour (about 16mm) ...	7
5. The Full-grown Caterpillar Descends into the Soil to Pupate	9
6. The Pupa of <i>C. binotalis</i> (about 9mm)	9
7. The Male of <i>C. binotalis</i> with Dark Hairs on the Anterior Margin of Each Forewing	10
8. The Female of <i>C. binotalis</i>	10
9. Larvae of <i>C. binotalis</i> in the Heart of Cabbage, to Produce a Mass of Webbing and Frass	12
10. Spores of <i>B. bassiana</i> on Potato Dextrose Agar	15
11. Spores of <i>M. anisopliae</i> var. <i>majus</i> on Potato Dextrose Agar	19
12. Spores of <i>P. fumosoroseus</i> on Media	22
13. Chinese Mustard Grown in the Glasshouse	24
14. Leaf Preparation for Dosage-Response Assay	25
15. Experimental Layout: Cabbage Plants in an Open Area	30
16. Conidia in Kaolin (Left) and Conidia in Palm Oil (Right)	31
17. Conidia Cultures in Rice Flour	32
18. <i>P. fumosoroseus</i> on Infected Larva	35
19. Larva Covered with Green Powdery Spores of <i>M. anisopliae</i> var. <i>majus</i>	36



20. <i>B. bassiana</i> was Observed to Sporulate Outside the <i>C. binotalis</i> Larva	36
21. Massive Amount of Conidia Attached Along the Cervical Membrane	43
22. Conidia of <i>P. fumosoroseus</i> Collected at the Setal Socket	43
23. Conidia of <i>P. fumosoroseus</i> in the Crochet	44
24. Conidia of <i>B. bassiana</i> on the Head of the Larva	44
25. Conidia of <i>M. anisopliae</i> var. <i>majus</i> on the Cuticle	45
26. Conidia of <i>P. fumosoroseus</i> Adhering on the Cuticle Surface with Tween 80 as the Sticky Agent	46
27. Germinating Conidium at About 4-6 Hours After Inoculation (a) <i>B. bassiana</i> (b) <i>M. anisopliae</i> var. <i>majus</i>	47
28. <i>P. fumosoroseus</i> with Thin-walled Germ Tube Penetrating the Cuticle	48
29. Conidium of <i>B. bassiana</i> Formed an Appressorium with (a) Short Germ Tube (b) Long Germ Tube	49
30. Close-up of Appressorium of <i>B. bassiana</i>	50
31(a & b). Germ Tubes of <i>B. bassiana</i> Penetrated Via the Spiracular Opening ...	52
32. Hyphae of <i>M. anisopliae</i> var.	53
33. Germ Tubes Penetrating Via Setal Socket	53
34. Collapsed Conidia (a) <i>P. fumosoroseus</i> (b) <i>M. anisopliae</i> var. <i>majus</i>	54
35 (a&b). Hyphae Ramification Inside the Hemocoel of Dead Larva	56
36. Healthy Tracheal Trunk	58
37 (a&b). Extensive Hyphal Growth in the Tracheal Trunk	59
38 (a&b). Hyphae Bursting Outward Outside of the Cuticle	60
39. Formation of Conidia (<i>B. bassiana</i>) at the Distal Ends of the Branched Conidiophores on the Infected Larva	62
40. Chains of Conidia (<i>B. bassiana</i>) on the Infected Larva	62



41. Conidiophores and Conidia of <i>M. anisopliae</i> var. <i>majus</i> Growing on the Infected Larva. The Layer of Bottle-shaped Conidiophores Can be Recognised Easily	63
42. The Larva was Enveloped with the Fungus	63
43. Secondary Fungal Growth on the Infected Larva (a) <i>Aspergillus</i> sp. (b) <i>Penicillium</i> sp.	64



CHAPTER 1

INTRODUCTION

Cruciferous vegetables are economically important throughout the world. In Malaysia, they are grown by small landholders around urban centres, in highlands and in specialised production areas. Often farmers use input-intensive agronomic practices because the sale of these vegetables provide an important source of ready cash income.

In recent years, however, crucifer production has been seriously affected by a steady increase in insect pests. The cabbage-heart caterpillar, *Crocidolomia binotalis* Zell. (Lepidoptera : Pyralidae) is considered as the second most important insect pest of cabbage and other crucifers. Severe damage to cabbage has been reported, especially in the Cameron Highlands (Ooi *et al.*, 1979). The larvae live gregariously feeding at first on the under side of cabbage leaves which may be eventually eaten completely. The growing point is often damaged leading to regeneration of multiple heads. As the larva bores into the cabbage head, it gets protected from contact with insecticides (Ooi, 1988). The larva is extremely mobile and can easily travel two or more meters to reach a host plant.

To control the insect pest, most farmers rely solely on chemical pesticides which are often used indiscriminately. A survey by Ooi and Sudderuddin (1978) showed that a mixture of two or more insecticides were applied in excess of the recommended rates

and sprayed as frequent as three times a week. These injudicious practices have resulted in many alarming problems such as development of insecticide resistance, pest resurgence, excessive chemical residues and environmental contamination. Concern about the negative effects of chemical insecticides have led to emphasis on alternative strategies for pest control. Meanwhile, the concept of integrated pest management (IPM) has begun to receive widespread acceptance and this will play an important role in establishing the biological control agents for the future (Jusoh *et al.*, 1982).

Microbial control is defined as part of biological control with the manipulation of pathogens or their by-products to cause the reduction of pest population. The unpredictable results with insect pathogens in field applications and the successful use of broad spectrum pesticides to reduce pest population (Tanada and Kaya, 1993) has caused the development of microbial control agents to lag behind the other methods of insect control, especially the chemical control, though the concept of microbial control originated over a hundred years ago.

Entomogenous fungi have been mentioned to play a uniquely important role in the history of microbial control of insects. Roberts (1981) defined an entomopathogenic fungus as a "pathogen causing an early death of the host by penetrating and proliferating inside the host, which is killed by being deprived of soluble nutrients in its hemolymph, by invasion or digestion of its tissues, and/ or by the release of toxins from the fungus". Research during the last decade has proved that fungi have a prominent place in insect control because the entomopathogenic fungi constitute the largest group of insect

pathogens. Furthermore, some entomopathogenic fungi are undergoing large-scale field test and a significant advance in development and manufacturing of these agents in the future is expected with recent biotechnological innovation (Khachatourians, 1986).

In Malaysia, indigenous isolates of *Beauveria bassiana* (Bals.) and *Paecilomyces fumosoroseus* (Wize) Brown and Smith have been reported recently in laboratory studies to be effective against the cosmopolitan diamondback moth (DBM), *Plutella xylostella* (L.) (Ibrahim *et al.*, 1998; Hashim *et al.*, 1999). *Metarhizium anisopliae* (Metch.) has also been reported to be infectious to termites (Sajap *et al.*, 1990) and rhinoceros beetles (Sajap, 1991). Thus it is of great interest to know whether these isolates are also efficacious against the other lepidopteran pest of cabbage such as the cabbage-heart caterpillar, *C. binotalis*. Hence, the objectives of this study are:-

1. To determine the bioefficacy of the isolates *B. bassiana*, *P. fumosoroseus* and *M. anisopliae* against *C. binotalis*.
2. To critically examine the mechanism of infection through histopathological examination of infected larvae of *C. binotalis*.
3. To compare the field efficacy of several raw preparations of formulations using the best fungal isolate from the entomopathogenic fungi tested against *C. binotalis*.

CHAPTER 2

LITERATURE REVIEW

2.1 *Crocidolomia binotalis* Zeller

2.1.1 Introduction

Crocidolomia binotalis Zeller or cabbage-heart caterpillar, is a pyralid moth. Zeller described *C. binotalis* in 1852 from a specimen collected in South Africa. Early synonyms of *C. binotalis* were *C. comalis* (Guence) (1854) from central India and *C. incomalis* (Guenee) (1854) from Java.

The moths are almost exclusively found in hot humid highland tropics and constitute a more serious pest problem during the dry season since heavy rains can drown small larvae. The native range of the moth probably embraced both Africa and Asia (Waterhouse and Norris, 1987). *C. binotalis* occurs in extensive areas of west, south and east Africa, throughout the warmer region of Asia, in north-eastern Australia, Carolina Is, Cook Is, Fiji, Indonesia, Marianas, New Caledonia, Niuc, Norfolk Is, Papua New Guinea, The Philippines, Western Samoa, Solomon Is and Tonga. This pest is not reported in Europe and the Americas (Waterhouse and Norris, 1987). In Malaysia, it has been placed as second in importance after the DBM. Probably the insect is very

much held in check under existing conditions. However, severe damage to cabbage has been reported in the Cameron Highlands (Ooi *et al.*, 1979).

2.1.2 Biology and Ecology

Lifecycle of *C. binotalis* is completed in approximately 28 days, depending on temperature and humidity. The eggs are laid in batches, overlapping each other on the lower surfaces of the leaves (Plate 1). Smooth leaf surface are preferred for oviposition. Most eggs are laid, close to the midrib or the veins on the leaf. The eggs when newly laid are greenish but later turn brown prior to eclosion. They are flat, ovoid and are packed neatly like roof tiles. An egg measures about 0.36mm in width and 0.54mm in length. The number of eggs per batch varies from 8 to 114 per batch (Ooi *et al.*, 1979). Incubation period takes five to fifteen days in the lowland, but depending on temperature (Water and Norris, 1987).

There are five larval instars, and feeding occupies fourteen to forty-two days (Waterhouse and Norris, 1987). First instar larvae are slender. Head and thorax are dark brown and the body is greenish yellow in color (Plate 2). The second instar is green and the head is brown with light pathes (Plate 3). The late larval instars are characterised by whitish longitudinal stripes, three dorsal and one on each lateral side (Plate 4). All the abdominal segments have tubercles. Between the dorsal and lateral stripes are found black round spots. These are the bases of long hairs on the body.




Plate 1 : The eggs of *C. binotalis* are laid in batches on the lower surfaces of the leaf.



Plate 2 : First instar larvae of *C. binotalis* are slender and the body is greenish yellow in colour (about 2 mm).