



UNIVERSITI PUTRA MALAYSIA

**SOMATIC EMBRYOGENESIS AND PROTOPLAST ISOLATION AND
CULTURE IN PAPAYA (CARICA PAPAYA L.)**

AGUS SUTANTO

FP 2001 16

**SOMATIC EMBRYOGENESIS AND PROTOPLAST ISOLATION AND
CULTURE IN PAPAYA (*CARICA PAPAYA* L.)**

By

AGUS SUTANTO

**Thesis Submitted in Fulfilment of the Requirement for the
Degree of Master of Science in the Faculty of Agriculture
Universiti Putra Malaysia**

September 2001



Dedicated to:

My beloved wife Susilowati

My dearest son Ikhsan Fitrianto

Departed soul of my father Munari

BIODATA OF THE AUTHOR

The author was born on 3 August 1967 in Malang, East Java, Indonesia. He graduated from primary school, junior high school and senior high school in 1980, 1983 and 1985 respectively. In 1985, he continued his study at Brawijaya University Malang. He began to study tissue culture in 1990 and completed his study in the university in 1991 with specialization in tissue culture.

After graduation, he worked at Tlekung Horticultural Research Station, Malang, as a researcher on tissue culture of citrus, apple and mangosteen. He participated in a Banana Tissue Culture Course at Taiwan Banana Research Institute (TBRI) and Banana Breeding Workshop at Malaysian Research and Development Institute (MARDI), both in 1994. In 1995, due to reorganization of the Department of Agriculture, he shifted to Solok Research Institute for Fruits, West Sumatera. He worked on banana breeding and tissue culture of citrus, mango and papaya. In 1996, under the collaboration of the Department of Agriculture and International Network for the Improvement of Banana and Plantain (INIBAP), he went to Maluku islands to collect *Musa* germplasms.

In 1997, he was recommended by the Agency for Agriculture Research and Development (AARD), Department of Agriculture, Republic of Indonesia, to continue his study and began his Master of Science program in November 1998 at the Department of Crop Science, Faculty of Agriculture, Universiti Putra Malaysia in the field of Agricultural Biotechnology.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Master of Science

**SOMATIC EMBRYOGENESIS AND PROTOPLAST ISOLATION AND
CULTURE IN PAPAYA (*CARICA PAPAYA* L.)**

By

AGUS SUTANTO

September 2001

Chairman : Dr. Maheran Abdul Aziz

Faculty : Agriculture

This study was carried out with the main objective of establishing somatic embryo production in papaya that may be used for further genetic improvement of the crop. The specific objectives were to induce somatic embryogenesis from immature zygotic embryos, to establish cell suspension culture of somatic embryos and to isolate and culture protoplasts from *in vitro* leaves.

The first experiment studied the effect of growth regulators on the induction and regeneration of somatic embryos from immature zygotic embryos. The combinations of 2,4-dichlorophenoxyacetic acid (2,4-D) and 6-benzylaminopurine (BAP) at various concentrations were assessed in order to determine the best combination for somatic embryogenesis. The experiment was conducted in a Completely Randomized Design.

The results showed that higher 2,4-D with or without BAP in the medium increased the percentage of somatic embryo formation and the number of somatic embryos per explant. MS medium supplemented with 5.0 mg/L 2,4-D promoted the highest



percentage of somatic embryogenesis (74.55%), while the combination of 1.0 mg/L 2,4-D and 0.01 mg/L BAP produced the highest percentage of callus formation (52.58%). The highest number of somatic embryos per explant (66.61) was obtained when 5.0 mg/L 2,4-D and 0.01 mg/L BAP were added into MS medium. For germination of somatic embryos, the result was very inconsistent; nevertheless, the best treatment for plantlet formation was MS medium without growth hormone (N₀B₀).

The second experiment was the establishment of cell suspension culture of somatic embryos. Four weeks after the transfer of somatic embryos into liquid MS medium containing 2.0 mg/L 2,4-D, pro-embryogenic masses (PEMs) were formed in the suspension. Maturation of embryos was achieved on transferring the heart-shaped embryos to liquid MS medium without growth regulator. Germination of somatic embryos occurred following the subculture of cotyledonary embryos from liquid MS medium with 0.2 mg/L BAP and 0.02 mg/L naphthalene acetic acid (NAA) to solid hormone-free MS medium.

The third experiment was the isolation, purification and culture of protoplasts. *In vitro* leaves derived from shoot tips cultured on medium with 0.05 mg/L or 0.10 mg/L BAP were used as the source of protoplasts. Enzyme solution containing cell protoplast washing (CPW) salts, 0.6% (w/v) Macerozyme R-10, 2.0% (w/v) Cellulase R-10, 0.2% (w/v) Driselase and 9.0% (w/v) mannitol was used for protoplast isolation. Protoplasts were purified using two methods of purification; filtration followed by repeated resuspension, and floatation on dense sucrose solution.

The results showed that the highest protoplast yield and percentage of viable protoplasts were obtained when *in vitro* leaves from shoot tip culture placed on MS medium with 0.05 mg/L BAP were used as the source of protoplasts and the protoplasts purified by floatation on dense sucrose solution. Three days following the protoplast culture in modified KM8p medium, first division of protoplast-derived cells was observed, but they failed to grow further.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**EMBRIOGENESIS SOMA DAN PENGASINGAN SERTA KULTUR
PROTOPLAS BAGI TANAMAN BETIK (*CARICA PAPAYA* L.)**

Oleh

AGUS SUTANTO

September 2001

Pengerusi : Dr. Maheran Abdul Aziz

Fakulti : Pertanian

Penyelidikan ini dilaksanakan dengan objektif utama untuk mewujudkan produksi embrio soma betik, yang boleh digunakan untuk perbaikan genetik tanaman tersebut. Objektif khas adalah induksi dan regenerasi embrio soma daripada embrio zigotik belum matang, mewujudkan kultur ampai sel daripada embrio soma dan isolasi serta kultur protoplas dari daun yang dikulturkan secara *in vitro*.

Eksperimen pertama mengkaji kesan pengawalatur tumbesaran terhadap induksi dan regenerasi embrio soma daripada embrio zigotik belum matang. Kombinasi asid diklorofenoksi asetik (2,4-D) dan 6-bensilaminopurina (BAP) pada beberapa kepekatan dicuba untuk mengenalpasti kombinasi terbaik untuk induksi embrio soma. Kajian ini dijalankan dengan menggunakan Rekabentuk Rawak Lengkap (CRD).

Hasil kajian menunjukkan bahawa media yang mengandungi kepekatan 2,4-D yang tinggi dengan atau tanpa BAP merangsang dan meningkatkan peratus pembentukan embryo soma dan bilangan embryo soma per eksplan. Media MS dengan 5.0 mg/L

2,4-D merangsang peratus pembentukan embrio soma tertinggi (74.55%), sementara kombinasi 1.0 mg/L 2,4-D dan 0.01 mg/L BAP menghasilkan peratus pembentukan kalus tertinggi (52.58%). Bilangan embrio soma per eksplan tertinggi diperolehi apabila 5.0 mg/L 2,4-D dan 0.01 mg/L BAP ditambahkan ke dalam media MS. Untuk percambahan embrio soma, hasil kajian tidak konsisten, tetapi rawatan terbaik untuk pembentukan planlet adalah media MS tanpa pengawalatur tumbesaran (N₀B₀).

Eksperimen kedua adalah perwujudan kultur ampai sel daripada embrio soma. Empat minggu setelah embrio soma dipindah ke dalam media cecair MS yang mengandungi 2.0 mg/L 2,4-D, 'pro-embryogenic masses' (PEM) terbentuk di dalam kultur ampai. Kematangan embrio dicapai dengan memindahkan embrio berbentuk hati ke dalam media MS tanpa pengawalatur tumbesaran. Percambahan embrio soma berlaku berikutan pemindahan embrio berkotiledon dari media cecair MS dengan 0.2 mg/L BAP dan 0.02 mg/L NAA ke media pepejal MS tanpa pengawalatur tumbesaran.

Eksperimen ketiga adalah pengasingan, penulenan dan kultur protoplas. Daun *in vitro* daripada tunas pucuk yang dikultur dalam media MS dengan 0.05 mg/L atau 0.10 mg/L BAP digunakan sebagai sumber protoplas. Larutan enzim yang mengandungi garam penyucian sel protoplas (CPW), 0.6% (w/v) Macerozyme R-10, 2.0% (w/v) Cellulase R-10, 0.2% (w/v) Driselase dan 9.0% (w/v) mannitol digunakan untuk isolasi protoplas. Penulenan protoplas menggunakan dua kaedah, iaitu penurasan yang diikuti dengan pengampaian semula, dan sistem pengapungan pada larutan sukrosa ketumpatan tinggi.

Kajian menunjukkan hasil protoplas dan peratus protoplas bernas yang tertinggi diperolehi apabila daun yang berasal daripada tunas pucuk yang dikulturkan pada media MS dengan 0.05 mg/L BAP digunakan sebagai sumber protoplas dan penulenan protoplas menggunakan sistem pengapungan pada larutan sukrosa ketumpatan tinggi. Tiga hari setelah protoplas dikultur dalam media KM8p yang telah diubahsuai kandungannya, sel-sel yang terbentuk dari protoplas mengalami pembahagian pertama, tetapi gagal untuk meneruskan pertumbuhan.

ACKNOWLEDGEMENTS

I should like to express my gratitude to Dr. Maheran Abdul Aziz, Mr. Azmi Abdul Rashid, M.Phil. and Prof. Dr. Marziah Mahmood for their invaluable guidance and encouragement during the course of my study and the preparation of this manuscript.

I am grateful to the Project Manager of ARMP II (Agriculture Research and Management Project II), Agency for Agriculture Research and Development (AARD), Department of Agriculture, Republic of Indonesia for the scholarship and the opportunity given to me in pursuing a postgraduate program at the Universiti Putra Malaysia.

I would also like to express my deepest thanks to my wife Susilowati, my mother Sukarti and my parents in law Soekardi and Yuliana for their encouragement, patience, moral support and inspiration during the period of study. Finally, I show my appreciation to Mr. Md. Humayun Kabir, for his assistance in the laboratory work during the study. Above all, Allah the Most Gracious and Merciful who gave me the strength to complete the work and made all things well.

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF PLATES	xviii
LIST OF ABBREVIATIONS/NOTATIONS	xx
CHAPTER	
1. GENERAL INTRODUCTION	1 1
1 1 Background	1 1
1 2 Objectives	1 2
2. LITERATURE REVIEW	2 1
2 1 Papaya	2 1
2 1 1 Origin of <i>Carica</i> sp	2 1
2 1 2 Sex Type of <i>Carica</i> sp	2 1
2 1 3 Commercial Cultivars of Papaya in ASEAN	2 2
2 2 Tissue Culture Techniques of Papaya	2 3
2 2 1 Shoot Culture	2 4
2 2 2 Organogenesis	2 5
2 2 3 Somatic Embryogenesis	2 6
2 2 4 Cell Suspension Culture	2 10
2 2 4 1 Isolation of Single Cell	2 10
2 2 4 2 Totipotency of Cells	2 11
2 2 5 Embryo Rescue	2 12
2 2 6 Anther Culture	2 12
2 2 7 Protoplast Culture	2 13
2 2 7 1 Isolation and Purification of Protoplasts	2 14
2 2 7 2 Culture of Protoplasts	2 18
2 3 Cultivar Improvement	2 19
2 3 1 Resistance to Papaya Ringspot Virus	2 19
2 3 2 Postharvest Technology	2 21



3. SOMATIC EMBRYOGENESIS OF IMMATURE ZYGOTIC EMBRYOS	3 1
3 1 Preface	3 1
3 2 Materials and Methods	3 3
3 2 1 Explant Preparation and Culture	3 3
3 2 2 Experiments	3 4
3 2 2 1 Induction of Somatic Embryogenesis	3 4
3 2 2 2 Germination of Somatic Embryos	3 4
3 2 3 Experimental Design and Statistical Analysis	3 4
3 3 Results	3 8
3 3 1 Induction of Somatic Embryogenesis	3 8
3 3 2 Germination of Somatic Embryos	3 14
3 4 Discussion	3 17
3 4 1 Induction of Somatic Embryogenesis	3 17
3 4 2 Recurrent Embryogenesis	3 20
3 4 3 Germination of Somatic Embryos	3 21
4. ESTABLISHMENT OF CELL SUSPENSION CULTURE OF SOMATIC EMBRYOS	4 1
4 1 Preface	4 1
4 2 Materials and Methods	4 2
4 3 Results	4 3
4 4 Discussion	4 10
4 4 1 The Development of Embryogenic Cells in The Suspension	4 10
4 4 2 Scale Up Production of Somatic Embryos	4 13
5. ISOLATION AND CULTURE OF PROTOPLAST FROM <i>IN VITRO</i> LEAVES	5 1
5 1 Preface	5 1
5 2 Materials and Methods	5 2
5 2 1 Explant Preparation	5 2
5 2 2 Isolation of Protoplasts	5 2
5 2 3 Purification of Protoplasts	5 3
5 2 4 Culture of Protoplasts	5 3
5 2 5 Treatments	5 4
5 3 Results	5 6
5 4 Discussion	5 13
5 4 1 Isolation of Protoplasts	5 13
5 4 2 Purification of Protoplasts	5 13
5 4 2 1 Filtration and Resuspension Purification	5 14
5 4 2 2 Floation on Dense Sugar Solution	5 14
5 4 3 Culture of Protoplasts	5 16



6. GENERAL DISCUSSION	6 1
7. CONCLUSION	7 1
REFERENCES	R 1
APPENDICES	A 1
BIODATA OF THE AUTHOR	B 1



LIST OF TABLES

Table		Page
3.1	The combination of 2,4-D (D) and BAP (B) concentrations for the induction of somatic embryogenesis	3.6
3.2	The combination of NAA (N) and BAP (B) concentrations for the germination of somatic embryos	3.7
4.1	Estimation of yield of embryos at various developmental stages	4.8
5.1	The combination of sources of protoplasts (B) and purification systems (P) for protoplast isolation and purification	5.5



LIST OF FIGURES

Figure	Page
3 1 The percentage of somatic embryo (A) and callus formation (B) and the number of somatic embryos per explant (C) after two months of culture on induction medium	3 10
3 2 The percentage of somatic embryos producing shoots only (A), roots only (B) and plantlet (C) on plantlet germination medium after one month of culture	3 15
5 1 The yield of protoplasts per millilitre following purification	5 10
5 2 The percentage of viable protoplasts after purification and FDA staining	5 10
6 1 A flowchart on the integration of tissue culture techniques and genetic engineering for papaya improvement	6 5



LIST OF PLATES

Plate	Page
3 1 Zygotic embryo three days after culture on induction medium	3 9
3 2 The apical dome of papaya zygotic embryo enlarges (white arrow) a week after culture on induction medium	3 9
3 3 Somatic embryos (white arrow) induced directly without callus formation from the apical dome of a zygotic embryo and photographed after three weeks of culture	3 12
3 4 Somatic embryos (white arrow) and callus (black arrow) growing together from the apical dome of a zygotic embryo	3 12
3 5 Globular somatic embryos (white arrows) produced indirectly from callus of a zygotic embryo on medium containing combination of high 2,4-D and BAP concentrations	3 13
3 6 Secondary globular embryos (white arrows) emerged from the surface of primary globular embryos after eight weeks of culture	3 13
3 7 Hypocotyl and callus growing together on germination medium containing BAP and NAA White arrows – new leaves, black arrow = callus	3 16
3 8 Somatic embryos germinating on germination medium	3 18
3 9 Plantlet from somatic embryo eight weeks after subculture on germination medium	3 18
4 1 The variation and viability of cell suspension assessed using FDA after four weeks of subculture to liquid medium A) Under normal light B) The same field but illuminated with UV light Viable cells were shown by greenish or yellowish colour	4 4
4 2 A) Somatic embryos transferred to liquid MS medium containing 2 0 mg/L 2 4-D B) The turbidity of cell suspension four weeks after subculture C) Sieved suspension with some globular somatic embryos D) High-frequency of somatic embryo formation	4 5
4 3 The development of somatic embryos in the cell suspension A) The embryogenic cells with distinct nucleolus (black arrow) B) The embryogenic cell has divided into eight-celled structure C) Pro-embryogenic masses D) Globular somatic embryo cluster with epidermal layer (white arrow) E) Heart-shaped embryo F) Torpedo-shaped embryo G) Cotyledonary embryo	4 6



4 4	Maturation of somatic embryos in liquid hormone-free MS medium:	4 9
4 5	Plantlets on solid hormone-free MS medium eight weeks after culture	4 9
5 1	The sources of protoplasts A) Leaves from MS medium with 0 05 mg/L BAP B) Leaves from medium with 0 10 mg/L BAP	5 7
5 2	The protoplast suspension after floatation purification showing floated protoplasts (P) and debris (D)	5 8
5 3	The protoplasts after purification A) Using filtration and resuspension system B) Using floatation purification system d = subcellular debris, bp = broken protoplasts, i = intact protoplast	5 9
5 4	Viable protoplasts after staining with fluorescein diacetate (FDA)	5 12
5 5	A) Cell wall formation two days after culture B) First cell division after 3 days of culture	5 12



LIST OF ABBREVIATIONS/NOTATIONS

2,4-D	2,4-dichlorophenoxyacetic acid
AARD	Agency for Agriculture Research and Development
ABA	abscisic acid
ACC	1-aminocyclopropane-1-carboxylic
ANOVA	analysis of variance
Arcsin	$\sin^{-1}(X)$
ASEAN	Association of South East Asia Nations
BAP	6-benzylaminopurine
<i>carETR1</i>	ethylene response 1 gene isolated from <i>Carica</i>
cDNA	copy of deoxyribonucleic acid
CPW	cell and protoplast washing medium (Frearson <i>et al.</i> 1973)
CRD	completely randomized design
cv.	cultivar
DMRT	duncan multiple range test
e.g.	exempli gratia (for example)
<i>et al.</i>	<i>et alia</i>
etc.	et cetera
F ₁	first filial generation
FDA	fluorescein diacetate
Fe-EDTA	iron ethylene diamine tetraacetic acid
g/L	gram per liter
h.	hours
IAA	indole acetic acid
IBA	indole butyric acid
<i>i.e.</i>	<i>id est</i> (that is)
IEDCs	induced embryogenic determined cells
INIBAP	International Network for the Improvement of Banana and Plantain
KCal	kilo calories



KM8p	Kao and Michayluk (1975) basal medium
M	molar
MARDI	Malaysian Agriculture Research and Development Institute
mg/L	milligram per litre
mM	millimolar
MS	Murashige and Skoog (1962) basal medium
NAA	naphthalene acetic acid
NCSS	Number Cruncher Statistical System
NN	Nitch and Nitch (1969) basal medium
PEDCs	pre-embryogenic determined cells
PEMs	pro-embryogenic masses
pH	$-\log (H^+)$
PRV	papaya ring spot virus
Ri	root-induction
RITA	<i>Réceptient à Immersion Temporarire Automatisé</i>
RNA	ribonucleic acid
rpm	rotary per minute
sp.	species
TBRI	Taiwan Banana Research Institute
Ti	tumor-induction
TIBA	2,3,5 tri-iodobenzoic acid
USA	United State of America
UV	ultra violet light
v/v	volume to volume
VBCB	virus bintik cincin betik
w/v	weight to volume
WPM	woody plant basal medium (Lloyd and McCown, 1981)
x g	x gravity (relative centrifugal force)
%	percent
√	square root



α	level of significance
μg	microgram
μm	micrometer

CHAPTER 1

GENERAL INTRODUCTION

1.1 Background

Papaya is an important fruit in South East Asia countries such as Thailand, Indonesia, the Philippines and Malaysia. The ripe fruit is usually eaten as dessert, processed into jams, juice, and soft drink and as fillers for sauces. Besides supplying food, an important product of papaya is papain, which is in great demand in the international market, particularly United Kingdom and USA. Papain is used in meat tenderizing, manufacture of chewing gum and cosmetics, tannin hides, degumming of natural silk and to give shrink-resistance to wool. In South East Asia, papain is produced commercially only in the Philippines (Rohani, 1994).

Papaya is a very wholesome fruit and provides a cheap source of vitamins and minerals in the daily diet of the people. The fruit contains 1.0-1.5% protein and a plentiful source of carotene (1,160-2,431 μg per 100 g edible portion), the precursor of vitamin A. It is also a good source of vitamin C (69-71 mg/100 g) and mineral such as calcium (11-31 mg/100 g) and potassium (39-337 mg/100 g). The fruit is also popular with dieters since it is low in fat (0.1%), carbohydrate (7-13%) and calories (35-59 KCal/100 g) (Rohani, 1994).

The popularity of papaya fruit and the ease with which it can be propagated has made it ubiquitous in tropical and subtropical region. Local papayas are

