



UNIVERSITI PUTRA MALAYSIA

**DEVELOPMENT OF MICROPROPAGATION SYSTEM AND
REDUCTION OF HYPERHYDRICITY IN REGENERANTS OF
CARNATION (DIANTHUS CARYOPHYLLUS L. CV. MALDIVES)**

BUDI WINARTO

FP 2002 18

**DEVELOPMENT OF MICROPROPAGATION SYSTEM AND REDUCTION
OF HYPERHYDRICITY IN REGENERANTS OF CARNATION (*DIANTHUS
CARYOPHYLLUS* L. cv. MALDIVES)**

BUDI WINARTO

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirement for the Degree of Master of Science**

October 2002



***Dedicated to:
My wife Nuri Rianti M
My son Yoga Aninditya
My father Sugiyono (Alm.)***



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in Fulfillment
of the Requirement for the degree of Master of Science

**DEVELOPMENT OF MICROPROPAGATION SYSTEM AND REDUCTION
OF HYPERHYDRICITY IN REGENERANTS OF CARNATION (*DIANTHUS
CARYOPHYLLUS* L. CV. MALDIVES)**

By

BUDI WINARTO

October 2002

Chairperson : Dr. Maheran Abdul Aziz

Faculty : Agriculture

This study was carried out with the main objectives of developing a micropropagation system for *Dianthus caryophyllus* cv. Maldives and reducing hyperhydricity for healthy shoot production. The development of a micropropagation system included selection of explant and combination-concentration of growth regulators, optimization, multiplication of shoots, rooting and acclimatization. Hyperhydricity study included selection of types of closure and gelling agents, application of ventilated culture vessel, multiplication of recovered shoots and acclimatization of recovered plantlets. The experiment was factorial arranged in a randomized complete block design with four replications. Each treatment consisted of twelve explants per replicate.

In axillary proliferation of shoots using two types of explant and five combination-concentrations of growth regulators, node explant placed on MS medium containing 1.0 mg/L BA and 0.1 mg/L NAA was the most suitable combination in stimulating



high axillary shoot production with low rate of hyperhydricity. Lowering the concentration of NAA from 0.1 mg/L to 0.05 mg/L in combination with 1.0 mg/L BA in the optimization experiment improved axillary shoot production from 4.9 to 5.6 shoots per explant and reduced hyperhydricity to less than 30%.

In adventitious shoot formation from three explants placed on five concentrations of BA and NAA, the first young and fully developed leaves placed on MS medium supplemented with 0.1 mg/L BA and 0.01 mg/L NAA was the most suitable combination in inducing high adventitious shoot formation (43.3%) with lower hyperhydricity (60.0%) compared to other combinations tested.

MS medium containing 1.0 mg/L BA with 0.05 mg/L NAA and 0.5 mg/L BA with 0.1 mg/L NAA were the most appropriate media in inducing high shoot multiplication, whereas MS medium supplemented with 0.1 mg/L BA with 0.02 mg/L NAA and 0.1 mg/L BA with 0.01 mg/L NAA were most suitable in producing good quality shoots for rooting. High production of good quality shoots were produced only after the first subculture and reduced in the subsequent subcultures.

Half-strength MS medium was appropriate in stimulating high root formation from both axillary proliferated shoots and shoots derived adventitiously. Based on economic consideration the use of carton paper as vessel closure and 7 g/L agar were applied in the next experiment. The treatment induced 87.5% of root formation with high number of roots per explant (6.5 roots) and 2.22 cm root length with good quality roots and shoots.



All potting media for acclimatization, except jiffy-7, stimulated high survival rate (80% - 100%) for both plantlets derived from axillary shoots and those derived from adventitious shoots. In the incubation room, paddy charcoal indicated high survival rate (97.9%) with high leaf chlorophyll content in both types of plantlets, but in the screen house, a mixture of kossas peat + paddy charcoal induced the highest survival rate (100%) and leaf chlorophyll content (0.3046 mg/mg). The potted plants flowered within 4.5 to 5 months after acclimatization.

Plastic wrap in combination with agar Type 900 were the most appropriate treatment in obtaining healthy axillary shoots. The combination exhibited lower hyperhydricity (6.0%) with higher chlorophyll content (0.1288 mg/mg) and maintained a low reduction of leaf chlorophyll content at 58.5% in field derived node explants. In node explants derived from hyperhydritised shoots, the combination reduced hyperhydricity of shoots to 22.7% and increased leaf chlorophyll content to 62.0%. Whereas MS medium containing 0.1 mg/L BA and 0.01 mg/L NAA with single layer of carton paper as closure based on economic consideration was appropriate in the recovery of normal shoots from succulent condition with lower hyperhydricity (67.1%). The recovered shoots were able to multiply and produced good quality axillary shoots until the third sub-culture. They were easily rooted and successfully acclimatized in paddy charcoal and kossas peat + soil (1:1, v/v) with high survival rate (80-100%). The plants were potted and indicated a normal growth and flowered 4 to 5 months after acclimatization.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Master Sains

**PERKEMBANGAN SISTEM PEMBIAKAN MIKRO DAN PENGURANGAN
HIPERHIDRISITI REGENERAN BUNGA TELUKI (*DIANTHUS
CARYOPHYLLUS* L. CV. MALDIVES)**

Oleh:

BUDI WINARTO

Oktober 2002

Pengerusi : Dr. Maheran Abdul Aziz

Fakulti : Pertanian

Penyelidikan ini dilaksanakan dengan objektif utama untuk menghasilkan satu sistem pembiakan mikro untuk *Dianthus caryophyllus* L. cv. Maldives dan mengurangkan hiperhidrisiti bagi pengeluaran pucuk yang sihat. Kajian sistem pembiakan mikro mencakupi pemilihan eksplan dan kombinasi kepekatan pengawalatur tumbesaran, pengoptimuman, penggandaan pucuk, pengakaran, dan aklimatisasi. Kajian mengenai hiperhidrisiti mencakupi pemilihan jenis penutup kelalang dan agen pepadat, penggunaan kelalang kultur berventilasi, penggandaan pucuk yang pulih, dan aklimatisasi pokok yang pulih daripada hiperhidrisiti. Kajian berfaktor ini telah dijalankan menggunakan rekabentuk rawak berblok penuh dengan empat replikasi. Setiap rawatan mengandungi duabelas eksplan per replikasi.

Bagi pembentukan tunas aksil dengan menggunakan dua jenis eksplan dan lima kombinasi kepekatan pengawal atur tumbesaran, nod eksplan yang diletakkan pada medium MS yang mengandungi 1.0 mg/L BA dan 0.1 mg/L NAA adalah kombinasi



yang paling sesuai bagi merangsang pengeluaran pucuk aksil yang tinggi dengan kadar hiperhidrisiti yang rendah. Pengurangan kepekatan NAA dari 0.1 mg/L kepada 0.05 mg/L dalam medium dikombinasikan dengan 1.0 mg/L BA bagi eksperimen pengoptimuman meningkatkan bilangan pucuk aksil yang diperolehi daripada 4.9 kepada 5.6 pucuk per eksplan dan menurunkan kadar hiperhidrisiti kepada kurang daripada 30%.

Bagi pembentukan pucuk adventitious pula, dari tiga jenis eksplan yang diletakkan pada lima kepekatan kombinasi BA dan NAA, eksplan daun pertama yang masih muda dan berkembang yang diletakkan pada medium MS yang mengandungi 0.1 mg/L BA dan 0.01 mg/L NAA adalah kombinasi yang paling sesuai bagi merangsang pembentukan pucuk adventitious yang tinggi (43.3%) dan kadar hiperhidrisiti yang lebih rendah (60.0%) berbanding dengan kombinasi lain.

Media MS yang mengandungi 1.0 mg/L BA dengan 0.05 mg/L NAA dan 0.5 mg/L BA dengan 0.1 mg/L NAA adalah media yang paling sesuai bagi merangsang penggandaan pucuk yang tinggi. Manakala media MS yang dibekalkan dengan 0.1 mg/L BA dengan 0.02 mg/L NAA dan 0.1 mg/L BA dengan 0.01 mg/L NAA adalah yang paling sesuai bagi menghasilkan kualiti pucuk yang baik bagi pengakaran. Pengeluaran pucuk yang tinggi dan berkualiti baik telah dihasilkan sehingga subkultur pertama tetapi menurun pada subkultur seterusnya.

Medium MS berkepekatan separuh adalah sesuai bagi merangsang pembentukan akar yang tinggi pada pucuk yang diperolehi daripada tunas aksil ataupun pucuk yang diperolehi secara adventitious. Berasaskan pertimbangan ekonomi penggunaan satu

lapis kertas karton sebagai penutup kelalang dan 7 g/L agar telah digunakan bagi kajian selanjutnya. Rawatan tersebut menghasilkan 87.5% pembentukan akar dengan jumlah akar per eksplan yang tinggi (6.5) dan panjang akar (2.22) dengan kualiti akar dan pucuk yang baik.

Semua media pasuan untuk kegunaan aklimatisasi, kecuali Jiffy 7, merangsang keupayaan hidup yang tinggi (80%-100%) dan baik bagi anak pokok yang diperolehi daripada tunas aksil maupun yang diperolehi daripada tunas adventitious. Dalam bilik inkubasi, arang sekam telah memberikan keupayaan hidup pokok (97.9%) dan kandungan klorofil daun telah didapati tinggi bagi kedua-dua jenis anak pokok, tetapi rawatan dalam rumah skrin, campuran daripada kossas peat + arang sekam merangsang keupayaan hidup tertinggi (100%) dan kadar klorofil (0.3046 mg/mg) yang paling tinggi. Anak pokok yang dipasukan berbunga dalam masa 4.5 sehingga 5 bulan selepas aklimatisasi.

Kombinasi pembungkus plastik dengan agar Type 900 adalah rawatan yang sesuai bagi mendapatkan pucuk aksil yang lebih sihat. Kombinasi ini menunjukkan kadar hiperhidrisiti yang rendah (6.0%) dengan kadar klorofil daun yang lebih tinggi (0.1288 mg/mg) dan mengekalkan penurunan klorofil yang rendah sehingga 58.5% pada nod eksplan yang diperolehi daripada pokok lapangan. Pada nod yang diperolehi daripada pokok yang hiperhidrisiti, kombinasi rawatan tersebut mengurangkan hiperhidrisiti pucuk kepada 22.7% dan mempertingkatkan kadar klorofil daun kepada 62.0%. Media MS yang mengandungi 0.1 mg/L BA dan 0.01 mg/L NAA dengan satu lapis kertas karton sebagai penutup adalah sesuai bagi membaik-pulihkan pucuk daripada keadaan sukulen dengan hiperhidrisiti yang lebih rendah (67.1%). Pucuk

yang pulih didapati masih mampu mengganda dan menghasilkan pucuk aksil yang berkualiti sehingga subkultur yang ketiga. Didapati pucuk mudah diakarkan dan berjaya diaklimatisasi dalam arang sekam dan kossas peat ditambah tanah (1:1, v/v) dengan keupayaan hidup yang tinggi (80-100%). Anak pokok telah dipasukan dan menunjukkan satu pertumbuhan yang normal. Pokok berbunga dalam masa 4 sehingga 5 bulan setelah aklimatisasi.



ACKNOWLEDGMENTS

I should like to express my gratitude to members of the Supervisory Committee, Dr. Maheran Abdul Aziz, Mr. Azmi Abdul Rashid, M.Phil. and Associate Professor Dr. Mohd. Razi Ismail from the Department of Crop Science, Faculty of Agriculture, Universiti Putra Malaysia for their invaluable guidance and encouragement during the course of my study and the preparation of this manuscript.

I am grateful to the Project Manager of PAATP (Participatory Development of Agricultural Technology Project), Agency for Agriculture Research and Development (AARD), Department of Agriculture, Republic of Indonesia for the scholarship and the opportunity given to me in pursuing a postgraduate program at Universiti Putra Malaysia.

I would also like to express my deepest thanks to my wife Nuri Rianti M. and my son Yoga Aninditya for their encouragement, patience, and moral support during the period of study. Finally, I would like to record my appreciation to Mr. Agus Sutanto for his assistance and to all *in vitro* laboratory members for their cooperation in the laboratory work during the study. Above all, Allah the Most Gracious and Merciful who gave me strength to complete the work and made all things well.



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of Supervisory Committee are as follows:

Maheran Abdul Aziz, Ph.D.
Faculty of Agriculture,
Universiti Putra Malaysia.
(Chairperson)

Azmi Abdul Rashid, M.Phil.
Faculty of Agriculture,
Universiti Putra Malaysia.
(Member)

Mohd Razi Ismail, Ph.D.
Associate Professor,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Member)

AINI IDERIS, Ph.D,
Professor/Dean,
School of Graduate Studies,
Universiti Putra Malaysia.

Date:



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	x
APPROVAL	xi
DECLARATION	xiii
LIST OF TABLES	xvii
LIST OF FIGURES	xx
LIST OF PLATES	xxii
LIST OF ABBREVIATIONS/NOTATIONS	xxiv
 CHAPTER	
1 INTRODUCTION	1.1
1.1 Background	1.1
1.2 Objectives	1.4
 2 LITERATURE REVIEW	 2.1
2.1 Carnation	2.1
2.1.1 <i>Dianthus caryophyllus</i>	2.1
2.1.2 Characteristics of Commercial Carnation	2.2
2.1.3 Floriculture Industry in Malaysia	2.4
2.2 Propagation of Carnation	2.5
2.3 Micropropagation	2.6
2.3.1 Stages in Micropropagation	2.7
2.3.2 Micropropagation Methods	2.8
2.3.3 Factor Affecting Micropropagation	2.10
2.3.4 Problems in Micropropagation	2.11
2.3 Carnation Tissue Culture	2.14
2.4.1 Development Carnation Tissue Culture	2.14
2.4.2 Explants in Carnation Tissue Culture	2.15
2.4.3 Explant Sterilization	2.16
2.4.4 Placement of Explants	2.16
2.4.5 Culture Medium	2.17
2.4.6 Growth Regulators	2.18
2.4.7 Culture Conditions	2.18
2.4.8 Optimization in Carnation Tissue Culture	2.19
2.4.9 Problem in Carnation Tissue Culture	2.20
2.5 Axillary Proliferation in Carnation	2.23
2.6 Adventitious Shoot Formation in Carnation	2.24
2.7 Multiplication of Shoots	2.25
2.8 Root Formation in Carnation	2.26



2.9	Acclimatization	2.28
3	DEVELOPMENT OF MICROPROPAGATION SYSTEM OF CARNATION (<i>Dianthus caryophyllus</i> L. cv. Maldives) : INDUCTION, OPTIMIZATION, AND MULTIPLICATION	3.1
3.1	Preface	3.1
3.2	Materials and methods	3.4
3.2.1	Location of Study	3.4
3.2.2	Plant Material	3.4
3.2.3	Surface Sterilization	3.5
3.2.4	Preparation of Explants	3.5
3.2.5	Basic Medium	3.7
3.2.6	Culture Conditions	3.7
3.2.7	Axillary Proliferation Studies	3.7
3.2.8	Adventitious Shoot Formation Studies	3.10
3.2.9	Multiplication of Shoots	3.15
3.3	Results	3.16
3.3.1	Axillary Proliferation	3.16
3.3.2	Adventitious Shoot Formation	3.25
3.3.3	Multiplication of Shoots	3.44
3.4	Discussion	3.48
3.4.1	Axillary Proliferation	3.48
3.4.2	Adventitious Shoot Formation	3.52
3.4.3	Multiplication of Shoots	3.56
4	DEVELOPMENT OF MICROPROPAGATION SYSTEM OF CARNATION (<i>Dianthus caryophyllus</i> L. cv. Maldives): ROOTING AND ACCLIMATIZATION	4.1
4.1	Preface	4.1
4.2	Materials and Methods	4.2
4.2.1	Root Formation Studies	4.2
4.2.2	Acclimatization Studies	4.6
4.3	Results	4.9
4.3.1	Root Formation	4.9
4.3.2	Acclimatization	4.22
4.4	Discussion	4.38
4.4.1	Root Formation	4.38
4.4.2	Acclimatization	4.45
5	REDUCTION OF HYPERHYDRICITY IN REGENERANTS OF CARNATION (<i>Dianthus caryophyllus</i> L. cv. Maldives)	5.1
5.1	Preface	5.1
5.2	Materials and Methods	5.4
5.2.1	Preparation of Hyperhydritised Shoots	5.4
5.2.2	Determination of Percentage of Hyperhydritised Shoots	5.5
5.2.3	Determination of Leaf Chlorophyll Content	5.5
5.2.4	Determination of Water Content and Dry Weight	5.7
5.2.5	Determination of Relative Water Loss (RWL), Increase in Agar Concentration (ACI), and Evaporation Rate (ER)	5.7



5.2.6	Experiment 1: Assessment of Different Types of Closure and Gelling Agents to Minimize Hyperhydricity	5.8
5.2.7	Experiment 2: Assessment of Ventilated Culture Vessels to Reduce Hyperhydricity	5.10
5.2.8	Multiplication of Recovered Hyperhydritised Shoots	5.13
5.2.9	Acclimatization of Recovered Plantlets	5.14
5.3	Results	5.16
5.3.1	Assessment of Different Types of Closure and Gelling Agents to Minimize Hyperhydricity	5.16
5.3.2	Assessment of Ventilated Culture Vessel to Reduce Hyperhydricity	5.29
5.3.3	Multiplication of Recovered Hyperhydritised Shoots	5.37
5.3.4	Acclimatization of Recovered Plantlets	5.40
5.4	Discussion	5.46
5.4.1	Assessment of Different Types of Closure and Gelling Agents to Minimize Hyperhydricity	5.46
5.4.2	Assessment of ventilated culture vessel to Reduce Hyperhydricity	5.52
5.4.3	Multiplication of Recovered Hyperhydritised Shoots	5.56
5.4.4	Acclimatization of Recovered Plantlets	5.57
6	GENERAL DISCUSSION AND CONCLUSION	6.1
6.1	Development of Micropropagation System of Carnation	6.1
6.2	Reduction of Hyperhydricity in Regenerants of Carnation	6.5
6.3	Commercial Aspect of Development of Micropropagation System in Carnation	6.8
6.4	Conclusion	6.12
6.4.1	Development of Micropropagation System	6.12
6.4.2	Reduction of Hyperhydricity in Regenerants of Carnation	6.14
6.4.3	Suggestion	6.15
	REFERENCES	R.1
	APPENDICES	A.1
	BIODATA OF THE AUTHOR	B.1



LIST OF TABLES

	Page
3.1 The combinations of treatments between BA and NAA concentrations	3.9
3.2 The combination of treatments between BA and NAA Concentrations	3.12
3.3 Interaction effect of explants and CC-GRs on percentage of transferable shoots (PTS, %)	3.21
3.4 Interaction effect of BA and NAA concentrations on number of shoots produced per explant (NSE)	3.24
3.5 Interaction effect of BA and NAA concentrations on height of shoots (HS, cm)	3.24
3.6 Interaction effect of explants and C-GRs on percentage of shoot regeneration (PSR, %)	3.33
3.7 Interaction effect of explants and C-GRs on number of shoots produced per explant (NSE)	3.33
3.8 Interaction effect of BA and NAA concentrations on percentage of hyperhydritised shoots (PHS, %)	3.38
3.9 Interaction effect of BA and NAA concentrations on percentage of transferable shoots (PTS, %)	3.38
3.10 Interaction effect of BA and NAA concentrations on height of shoots (HS, cm)	3.39
4.1 The combination of treatments between rooting media and shoot sources	4.4
4.2 The combination of treatments between types of closure and agar concentrations	4.6
4.3 The combination of treatments between potting media and plantlet sources	4.8
4.4 Interaction effect of shoot sources and rooting media on percentage of root formation (PRF, %)	4.14



4.5	Interaction effect of shoot sources and rooting media on number of roots produced per shoot (NRS)	4.14
4.6	Interaction effect of shoot sources and rooting media on root length (RL, cm)	4.15
4.7	Interaction effect of closures and agar concentrations on percentage of root formation (PRF, %)	4.19
4.8	Interaction effect of closures and agar concentrations on number of roots produced per shoot (NRS)	4.19
4.9	Interaction effect of closures and agar concentrations on root length (RL, cm)	4.19
4.10	Interaction effect of plantlet sources and potting media on leaf chlorophyll content (LCC, mg/mg)	4.24
4.11	Interaction effect of plantlet sources and potting media on number of leaves produced per plant (NLP)	4.25
4.12	Alteration in number of stomata/mm ² of leaf epidermis surface during acclimatization	4.31
4.13	Alteration in leaf chlorophyll content (mg/mg) during acclimatization	4.32
5.1	The combination of treatments between types of closure and gelling agents	5.8
5.2	The combination of treatments between types of closure and selected concentrations of BA and NAA (SC)	5.12
5.3	Interaction effect of types of closure and gelling agents on percentage of hyperhydritised shoots (PHS, %)	5.23
5.4	Interaction effect of types of closure and gelling agents on leaf chlorophyll content (LCC, mg/mg)	5.23
5.5	Interaction effect of types of closure and gelling agents on percentage of hyperhydritised shoots (PHS, %)	5.28
5.6	Interaction effect of types of closure and gelling agents on leaf chlorophyll content (LCC, mg/mg)	5.28
5.7	Interaction effect of selected concentrations of BA and NAA (SC) and closures on water content(WC,%)	5.36
5.8	Acclimatization of recovered plantlets in incubation room	5.42



5.9	Acclimatization of recovered plantlets in glasshouse	5.42
5.10	Alteration in leaf chlorophyll content (LCC, mg/mg), height of plants (HP, cm) and number of leaves per plant (NLP) during gradual acclimatization recovered plantlets	5.44



LIST OF FIGURES

	Page
3.1 Flowchart on development of micropropagation system for <i>D. caryophyllus</i> L. cv. Maldives	3.3
3.2 Effect of explants on axillary proliferation	3.18
3.3 Effect of explants on axillary proliferation	3.18
3.4 Effect of BA on axillary proliferation	3.22
3.5 Effect of NAA on axillary proliferation	3.23
3.6 Effect of explants on adventitious shoot formation	3.28
3.7 Effect of C-GRs on percentage of shoot regeneration (PSR, %) and percentage of hyperhydratised shoots (PHS, %)	3.31
3.8 Effect of C-GRs on number of shoots produced per explant (NSE)	3.32
3.9 Effect of BA concentrations on adventitious shoot formation	3.36
3.10 Effect of NAA concentrations on adventitious shoot formation	3.37
3.11 Effect of selected BA and NAA concentrations on total shoot production from node explants of field plants	3.45
3.12 Effect of selected BA and NAA concentrations on total shoot production of node explants derived from adventitious shoots	3.46
4.1 Effect of rooting media on percentage of root formation (PRF, %)	4.12
4.2 Effect of rooting media on number of roots produced per explant (NRE) and length of roots (LR, cm)	4.13
4.3 Effect of closures on percentage of root formation (PRF, %)	4.17
4.4 Effect of closures on number of roots produced per explant (NRE)	4.17
4.5 Effect of potting media on percentage of plantlet survival in incubation room acclimatization	4.23
4.6 Effect of potting media on percentage of plantlet survival in screen house acclimatization	4.28



5.1	Flowchart on reduction of hyperhydricity in regenerants of <i>D. caryophyllus</i> L. cv. Maldives	5.3
5.2	Effect of closures on percentage of hyperhydritised shoots (PHS, %)	5.17
5.3	Effect of closures on leaf chlorophyll content (LCC, mg/mg)	5.18
5.4	Relative water loss using different types of closure	5.19
5.5	Increase in agar concentration using different types of closure	5.19
5.6	Effect of gelling agents on percentage of hyperhydritised shoots (PHS, %)	5.21
5.7	Effect of gelling agents on leaf chlorophyll content (LCC, mg/mg)	5.21
5.8	Effect of closures on percentage of hyperhydritised shoots (PHS, %)	5.25
5.9	Effect of closures on leaf chlorophyll content (LCC, mg/mg)	5.25
5.10	Effect of gelling agents on percentage of hyperhydritised shoots (PHS, %)	5.26
5.11	Effect of gelling agents on leaf chlorophyll content (LCC, mg/mg)	5.27
5.12	Relative water loss using different types of closure	5.31
5.13	Increase in agar concentration using different types of closure	5.32
5.14	Effect of closures on hyperhydricity	5.33
5.15	Effect of selected concentrations of BA and NAA (SC) on hyperhydricity	5.35
5.16	Effect of selected concentrations of BA and NAA (SC) on total shoot production from node explants of recovered shoots	5.38



LIST OF PLATES

	Page
3.1 <i>Dianthus caryophyllus</i> L. cv. Maldives	3.4
3.2 Preparation of explants	3.6
3.3 Axillary proliferation from shoot tip explants	3.19
3.4 Axillary proliferation from node explants	3.20
3.5 Development of adventitious shoots from leaf explants	3.26
3.6 Adventitious shoots produced from leaf and internode explants 6 weeks after culture initiation	3.29
3.7 Effect of subculturing on adventitious shoot growth from leaf explants	3.30
3.8 Effect of subculturing on adventitious shoot growth from young internodes	3.30
3.9 Adventitious shoots produced on selected C-GRs 2 months after culture initiation	3.40
3.10 Histological study on development of adventitious shoots on leaf explant	3.43
3.11 Differences in axillary shoot multiplication on selected BA and NAA concentrations	3.47
4.1 Effect of rooting media on root formation	4.11
4.2 A-rooted shoot on half-strength MS with 7 g/L agar type 900 and filter paper as closure, B-rooted shoot on half-strength MS with g/L agar type 900 and carton paper as closure, C-rooted shoot on half-strength MS with 7 g/L agar type 900 and plastic wrap as closure (control)	4.21
4.3 Effect of potting media on growth and quality of plantlets	4.26
4.4 Growth of plantlets in screen house	4.29
4.5 Performance of plantlets in different potting media	4.30
4.6 Root development at different stages of acclimatization	4.33



4.7	Abnormal plantlets produced during the study	4.33
4.8	Recovery of normal shoots from the abnormal plantlets	4.34
4.9	Varied performance of leaves during acclimatization	4.36
4.10	Flowers of acclimatized plants 4.5 to 5 months after incubation in the glasshouse	4.37
5.1	Hyperhydritised shoots used as explant source	5.4
5.2	Succulent shoots derived adventitiously from leaf explants	5.11
5.3	Effect of types of closure and selected concentrations of BA and NAA (SC) on growth of succulent shoots	5.30
5.4	Effect of closures on water potential in the flasks	5.34
5.5	Effect of closures on hyperhydricity	5.37
5.6	Similar performance of shoot growth on SC-3 and SC-4	5.39
5.7	Plantlets in paddy charcoal one-month after acclimatization in incubation room	5.41
5.8	Plantlets in kossas peat + soil (1:1, v/v) one-month after acclimatization in incubation room	5.41
5.9	Normal growth of plantlets acclimatized in glasshouse	5.43
5.10	Similar growth performance of recovered plantlets (1) and axillary plantlets (2) during acclimatization in the glasshouse	5.43
5.11	Different performance of flowers produced from recovered and normal plants	5.45



LIST OF ABBREVIATIONS/NOTATIONS

A_{647}	- absorbance at 647 nanometers
A_{664}	- absorbance at 664 nanometers
AARD	- Agency for Agriculture Research and Development
ABA	- abscisic acid
ACI	- agar concentration increase
ANOVA	- analysis of variance
BA/BAP	- 6-benzyladenine/benzylaminopurine
Br	- boron
°C	- centigrade
Ca	- calcium
$CaCl_2$	- calcium chloride
CaMV	- Carnation Mottle Virus
Cl	- chloride
cm	- centimeter
CO_2	- carbon dioxide
C/N	- carbon-nitrogen ratio
CPA	- <i>p</i> -chlorophenoxy acetic acid
CRSV	- Carnation Ring Spot Virus
c.v.	- cultivar
CVMV	- Carnation Vein Mottle Virus
2,4-D	- 2,4-dichlorophenoxy acetic acid
DNMRT	- Duncan's New Multiple Range Test
DW	- dry weight
e.g.	- <i>exempli gracia</i> (for example)
ER	- evaporation rate
et al	- <i>et alia</i>
etc	- <i>et cetera</i>
FAA	- formaldehyde-glacial acetic acid-alcohol
f.sp	- <i>forma specialist</i>
FW	- fresh weight

