



UNIVERSITI PUTRA MALAYSIA

**PURIFICATION OF CITRUS TRISTEZA VIRUS AND GENERATION
OF MONOSPECIFIC POLYCLONAL ANTISERUM**

NURHADI

FP 2002 32

**PURIFICATION OF CITRUS TRISTEZA VIRUS
AND GENERATION OF MONOSPECIFIC POLYCLONAL ANTISERUM**

NURHADI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirement for the Degree of Master of Agricultural Science**

July 2002



DEDICATION

*to my wife Rini,
to my daughters and sons: Nuri, Zaldi, Nina, and Farhan,
you are really supporters, best friends, and assistants,
without your understandings, suggestions, patience, and endless praying,
this work could never have been completed.*

*to my parents, Moh. Usman Solichan and Zubaidah, (in memory)
to my parents-in law, A. Cholik Syukur and Minartiningih,
and to my family members,
without your sacrifice, patience and endless praying,
many problems faced during this study
might not have been solved.*

*finally, but not least, to Allyn (in memory),
you were, and still are, a constant source of inspiration, spirit, and encouragement,
nothing is going to change my love for you.*

it is appropriate that this work be dedicated to you all.

Abstract of thesis presented to the Senate of Universiti Putra Malaysia
in fulfillment of the requirement for the degree of Master of Agricultural Science

**PURIFICATION OF CITRUS TRISTEZA VIRUS
AND GENERATION OF MONOSPECIFIC POLYCLONAL ANTISERUM**

By

NURHADI

July 2002

Chairman : Associate Professor Kamaruzaman Sijam, Ph.D.

Faculty : Agriculture

Symptomatology, molecular weight of coat protein (CP) determination and purification procedure for citrus tristeza virus (CTV) antigen were studied to generate monospecific polyclonal antiserum. Results of survey indicated that CTV was found to attack citrus varieties such as *C. grandis*, *C. aurantifolia*, *C. reticulata*, *C. hystrix*, *C. nobilis* and *C. sinensis* with variable symptoms and degrees of severity. Virus complex varied among citrus varieties and cultivars, and there were four important sub isolates found, namely mandarin stem pitting, orange stem pitting, pomelo small fruit, and pomelo mild isolate.

In partial purification step, antigen was concentrated by polyethylene glycol (PEG) precipitation followed by low speed centrifugation. Semi purified

antigen was then finally purified by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE). A major protein band containing CP was excised and eluted using elution buffer containing 0.25 M Tris-HCl pH 6.8 + 0.1 % SDS. With a starting material consisting of 50-gram bark tissue of semi dormant flush, 750 µg of eluted CP as monospecific antigen was obtained. The use of PEG and NaCl for virus precipitation combined with low speed centrifugation in semi purification step, then followed by electrophoresed of semi purified virus preparation in final purification step effectively minimized virus losses during purification processes and minimized possible contamination of plant components in the final immunogen.

Monospecific polyclonal antiserum against citrus tristeza virus was generated using the viral CP. Upon SDS-PAGE of local CTV isolate designated as UPM/T002, two protein bands specific for CTV with molecular weight of 25 kDa and 33 kDa were obtained. The major band with molecular weight of 25 kDa that reacted strongly with commercial polyclonal antibody in double antibody sandwich enzyme-linked immunosorbent assay (DAS-ELISA), was used as the antigen for injection into female White Leghorn chicken.

Chicken immunoglobulin recovered from immature egg reacted strongly with semi purified CTV up to 1:4096 dilutions in microprecipitine test. In DAS-ELISA, using sap of infected plant, the sensitivity of coating

immunoglobulin and enzyme conjugate were 1:1,000 and 1:500 dilutions respectively. Immunoglobulin reacted specifically with CTV isolate and gave no background reaction to healthy plant sap. The simplicity of the procedure makes it economically acceptable and technically adoptable because the antigen can be prepared with limited chemical and equipment. This study is the first reporting antiserum production from CTV-CP using local CTV isolate from Peninsular Malaysia.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Master Sains Pertanian

**PENULENAN CITRUS TRISTEZA VIRUS
DAN PENJANAAN POLIKLONAL ANTISERUM MONOSPESIFIK**

Oleh

NURHADI

Julai 2002

Pengerusi : Profesor Madya Kamaruzaman Sijam Ph.D.

Fakulti : Pertanian

Pengenalpastian simptom, penentuan berat molekul kod protein dan kaedah penulenan untuk antigen citrus tristeza virus (CTV) telah dikaji untuk menghasilkan antiserum poliklonal monospesifik menggunakan isolat CTV setempat. Hasil survei menunjukkan bahwa CTV dijumpai menyerang varieti limau seperti *C. grandis*, *C. aurantifolia*, *C. reticulata*, *C. hystrix*, *C. nobilis* dan *C. sinensis* dengan simptom berbeza dan berbagai tingkat keparahan. Kompleks virus berbeza diantara varieti dan kultivar dan terdapat empat subisolat penting iaitu 'mandarin stem pitting', 'orange stem pitting', 'pomelo small fruit' dan 'pomelo mild isolat'.

Pada peringkat awal penulenan, antigen dipekatkan dengan menggunakan kaedah presipitasi polyethylene glycol (PEG) diikuti dengan pengemparan



berkuasa rendah. Antigen semi tulen selanjutnya dituliskan dengan kaedah dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE). Jalur utama protein yang mengandungi CP dipotong dan ditulen menggunakan 'elution buffer' yang mengandungi 0.25 M Tris-HCl pH 6.8 + 0.1 % SDS. Bahan awal seberat 50 g tisu kulit batang yang diperolehi daripada pucuk semi dorman berjaya menghasilkan 750 ug CP CTV sebagai antigen monospesifik. Penggunaan PEG dan NaCl untuk presipitasi virus yang dipadukan dengan pengemparan berkuasa rendah pada step semi penulenan, yang diikuti dengan elektroforesis daripada virus semi tulen pada step penulenan akhir, sangat berkesan untuk meminimumkan kehilangan virus semasa proses penulenan dan meminimumkan kehadiran bahan-bahan kontaminasi daripada komponen tumbuhan pada hasil akhir penulenan.

Antiserum poliklonal yang spesifik terhadap CTV telah berjaya dihasilkan menggunakan viral CP. Dengan menggunakan isolat CTV setempat UPM/T002, melalui analisis SDS-PAGE, dua saluran yang mempunyai berat molekul 25-kDa dan 33-kDa dan spesifik terhadap CTV telah dihasilkan. Jaluran tebal yang berat molekulnya 25-kDa dan bertindakbalas kuat dengan antiserum komersial ketika diuji dengan double antibody sandwich enzyme-linked immunosorbent assay (DAS-ELISA), digunakan sebagai antigen untuk tujuan imunisasi pada ayam telur jenis White Leghorn.

Immunoglobulin yang dituliskan daripada telur belum sempurna berinteraksi kuat dengan CTV semi tulen sehingga tahap pencairan 1:4096 semasa ujian mikropresipitin. Pada ujian DAS-ELISA, dengan menggunakan sap daripada tanaman berpenyakit, sensitifiti 'coating immunoglobulin' adalah 1:1,000 pencairan, manakala enzim konjugat pula pada pencairan 1:500. Immunoglobulin bertindak balas spesifik terhadap isolat CTV dan tidak menunjukkan reaksi tidak spesifik terhadap sampel sap sihat. Kesederhanaan kaedah penulenan dalam kajian ini menjadikannya dapat diterima dari segi ekonomi dan teknikal kerana antigennya dapat disediakan pada keadaan peralatan dan bahan kimia minimum. Kajian ini merupakan kajian pertama yang melaporkan penghasilan antiserum daripada telur ayam yang belum sempurna, yang dihasilkan daripada viral CP menggunakan isolat CTV setempat dari Semenanjung Malaysia.

ACKNOWLEDGEMENTS

Alhamdulillah, be praised to Allah SWT, the most beneficent, the most merciful, and the most compassionate for giving me the physical and psychological strength and comfort to complete this research project and thesis.

The author wishes to express his deepest personal appreciation and sincere thanks to Associate Professor Dr. Kamaruzaman Sijam, the chairman of the supervisory committee, for his excellent guidance, encouragement and constructive suggestions throughout the undertaking of this study. Without his persistent support, papers presented at the 23rd and 24th Symposiums of the Malaysian Society for Microbiology would not have been proceeded. The invaluable help of Dr. Inon Sulaiman and Dr. Maheran Abdul Aziz, members of the supervisory committee, in the discussion during preparations of the seminars are gratefully acknowledged, and especially for the detailed editing of all chapters and many constructive suggestions for improvement.

Special thanks are due to the Agency for Agricultural Research and Development (AARD) Government of Indonesia especially for Participatory Development of Agricultural Technology Project (ADB Loan No. 1526-INO) and to Dr. Djoko Said Damardjati, the Secretary of AARD, for the opportunity and financial support rendered to the author to pursue a



postgraduate study in the Department of Plant Protection, Universiti Putra Malaysia. Special thanks are due to Dr. Haryono and Ir. Hartiningsih M.Sc., for their generosity and help in overcoming living problems during the study.

Personal appreciation and thanks are due to Agus Sutanto not only for his assistance in computer ware but also for his potential source of inspiration to the author; to Tri Pudji and Sujadi for their valuable support on electrophoresis; to Sri Kusworini, Machful and Panca Djarot for their assistance in collecting and processing of samples. Encik Yusuf, Encik Zawawi, Puan Junaina and Encik Razali were instrumental in supporting the necessary facilities for the Laboratory of Microbiology, and providing conducive atmosphere for the author to carry out research activities. They were, and still are, providing valuable assistance, and it is appropriate that the author acknowledge them.

Finally but not least, to the author's lab mates, especially Khairul Mazmi, Norputeri Julaida Rahmat, and Ee Fong who have brought me up to the homely atmosphere with their generous hospitality. No words could express my thanks for you all.



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Master of Agricultural Science. The members of the Supervisory Committee are as follows:

KAMARUZAMAN SIJAM, Ph.D.

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

INON SULAIMAN, Ph.D.

Faculty of Agriculture
Universiti Putra Malaysia
(Member)

MAHERAN ABDUL AZIZ, Ph.D.

Faculty of Agriculture
Universiti Putra Malaysia
(Member)

AINI IDERIS, Ph.D.

Professor/Dean,
School of Graduate Studies,
Universiti Putra Malaysia

Date:

TABLE OF CONTENTS

DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL SHEETS	xi
DECLARATION FORM	xiii
LIST OF TABLES	xvii
LIST OF FIGURES	xix
LIST OF ABBREVIATIONS/GLOSSARY OF TERMS	xx
CHAPTER	
1 INTRODUCTION	1.1
1.1. Economic Importance of Citrus	1.1
1.2. Citrus Tristeza Virus as a Major Disease of Citrus	1.3
1.3. Recent Strategy to Control CTV	1.7
1.4. Discussion	1.9
1.5. Research Objectives	1.11
2 LITERATURE REVIEW	2.1
2.1 Taxonomy and Property of CTV Virion	2.1
2.2 Cytopathology and Transmissibility	2.4
2.3 Biological and Serological Diversity of CTV Isolates	2.8
2.4 Immunodiagnosis Methods using Polyclonal Antiserum	2.11
2.4.1 Enzyme-linked Immunosorbent Assay	2.12
2.4.2 Electro-immuno Blot	2.16
2.5 Factors Influencing Virus Titer and Purified Virus	2.18
2.5.1 Effect of Host Plant Cultivars on Virus Titer	2.18
2.5.2 Effect of Seasonal Growth of Host Plant on Virus Titer	2.21
2.5.3 Distribution within Plants and Effect of Bark Tissue Sources on Virus Titer	2.22
2.5.4 Effect of Buffers and Additives on Virus Extraction	2.25
2.5.5 Effect of Extraction, Clarification and Concentration on Purified Virus	2.28
2.5.6 Methods for Final Purification	2.34
2.6 Immunization	2.36



3	MATERIALS AND METHODS	3.1
3.1	Biological Diversity of CTV Isolates from Peninsular Malaysia and West Sumatera	3.1
3.1.1	Survey	3.1
3.1.2	Selection Criteria	3.2
3.1.3	ELISA Test	3.3
3.2	Determination and Characterization of CTV Coat Protein	3.4
3.2.1	Sources of CTV Isolates	3.4
3.2.2	SDS-PAGE	3.5
3.3	Study on Virus Titer in Relation to Tissue Types, Annual Growth Flushes, and the Development of Procedure for Virus Purification	3.8
3.3.1	Sources of Plant Tissues	3.8
3.3.2	Development of Procedure for Virus Purification	3.9
3.3.3	Monitoring Antigen Recovery	3.10
3.4	Generation of Polyclonal Antiserum for CTV using Virion Coat Protein	3.10
3.4.1	Isolation of Coat Protein Band	3.10
3.4.2	Production and Purification of Immunoglobulin	3.13
3.4.3	Serological Test	3.15
4	RESULTS AND DISCUSSION	4.1
4.1	Biological Diversity of CTV Isolates from Peninsular Malaysia and West Sumatera	4.1
4.2	Determination and Characterization of CTV Coat Protein	4.11
4.3	Study on Virus Titer in Relation to Tissue Types, Annual Growth Flushes, and the Development for Procedure for Virus Purification	4.14
4.3.1	Effect of Tissue Types, Annual Growth Flushes on Virus Titer	4.14
4.3.2	Development of Purification Procedure for CTV	4.16
4.4	Generation of Polyclonal Antiserum for CTV using Virion Coat Protein	4.20
4.4.1	Isolation of Coat Protein as an Antigen for Immunization	4.20
4.4.2	Production and Purification of Immunoglobulin	4.21
4.4.3	Serological Test of the Specific Antiserum	4.24

5	GENERAL DISCUSSION	5.1
5.1	Biological Diversity of CTV Isolates from Peninsular Malaysia and West Sumatera	5.1
5.2	Determination and Characterization of CTV Coat Protein	5.3
5.3	Study on Virus Titer in Relation to Tissue Types, Annual Growth Flushes, and the Development of Procedure for Virus Purification	5.5
5.4	Generation of Monospecific Polyclonal Antiserum for CTV using Virion Coat Protein	5.7
6	CONCLUSION	6.1
	REFERENCES	R.1
	APPENDICES	A.1
A.1.	Preparation of Stock Solutions, Reagents and Chemicals for ELISA	A.1
A.2.	Preparation of Stock Solutions, Reagents and Chemicals for Electrophoresis	A.2
A.3.	Procedure of Virus Extraction, Clarification and Concentration of Antigen	A.5
A.4.	Procedure of Antiserum Purification	A.6
A.5.	Procedure of Conjugation of Alkaline Phosphatase with Immunoglobulin	A.7
A.6.	Preparation of Antiserum, Antigen and Conjugate for Serological Tests	A.8
	BIODATA OF THE AUTHOR	B.1

LIST OF TABLES

Table	Page
1.1. Trend of citrus production from 1985 to 1995.	1.1
1.2. Development of citrus areas in Malaysia from 1987 to 1995 (Ko, 1996).	1.3
1.3. Geographical distribution of citrus tristeza virus.	1.5
2.1. Properties of some currently recognized closterovirus.	2.3
2.2. Species of aphids known to be vectors of citrus tristeza virus.	2.6
2.3. Host plant reaction to strains of citrus tristeza virus (Rocha-Pena et al., 1995).	2.10
2.4. Comparison of 24 variations of enzyme-linked immunosorbent assay for the detection of citrus tristeza virus (Cambra et al., 1997).	2.15
2.5. Enzyme-linked immunosorbent assay absorbance values for various leaf tissue types from citrus tristeza virus infected trees (Matthews et al., 1997).	2.19
2.6. Virus titer as determined by enzyme-linked immunosorbent assay values for three individual sweet orange trees and four tissue types (Matthews et al., 1997).	2.22
2.7. Extraction buffer and additives used in virus purification for citrus plants.	2.26
3.1. Origins and biological properties of citrus tristeza virus isolates used in this study.	3.5
4.1. Citrus varieties and cultivars observed for citrus tristeza virus occurrence in Peninsular Malaysia and West Sumatera.	4.2
4.2. Immunoglobulin produced in different fraction of DE-52 column chromatography elution after ammonium sulphate precipitation and membrane dialysis.	4.23

4.3.	Absorbance values for purified preparation of citrus tristeza virus to the immunoglobulin and enzyme conjugate of various dilutions in double antibody sandwich-enzyme-linked immunosorbent assay.	4.26
4.4.	Absorbance values of citrus tristeza virus isolates to the chicken immunoglobulin generated from viral coat protein in double antibody sandwich-enzyme-linked immunosorbent assay	4.27

LIST OF FIGURES

Figure		Page
2.1.	Electron micrograph of citrus tristeza virus particles (Su, 1998).	2.2
2.2.	The distribution of citrus tristeza virus in different parts of young sour orange seedling at different times of post inoculation.	2.24
3.1.	Schematic flow chart on the purification of antigen for the production of monospecific polyclonal antiserum of citrus tristeza virus.	3.11
4.1.	Incidence of citrus tristeza virus in Peninsular Malaysia and West Sumatera.	4.5
4.2.	Four-year-old pomelo infected with citrus tristeza virus showing brittle branches; small leaves, corking, with spotted chlorotic mottle typical of boron deficiency and small-sized fruit found on branches.	4.6
4.3.	Citrus tristeza virus infected trees showing tree declining and vein clearing, dwarfing with small leaves, and moderate stem pitting when the bark was peeled off.	4.8
4.4.	Citrus Tristeza Virus infected <i>C. aurantifolia</i> tree showing severe stem pitting, severe vein clearing, and intense deep pits when the bark was peeled off (West Pasaman-Sumatera).	4.10
4.5.	SDS-PAGE pattern of viral proteins purified from citrus tristeza virus isolate.	4.11
4.6.	SDS-PAGE pattern of viral proteins purified from bark of actively growing flush, semi dormant flush and dormant flush samples.	4.12
4.7.	Effect of tissue types and annual flushes on virus titer.	4.15
4.8.	SDS-PAGE pattern of viral proteins purified before the first PEG precipitation, after the first PEG precipitation, and after the second PEG precipitation.	4.17
4.9.	Monitoring of antigens recovery by double antibody sandwich-enzymelinked immunosorbent assay.	4.19
4.10.	Reaction of chicken polyclonal antiserum generated from viral coat protein against purified virus in microprecipitine test.	4.25



LIST OF ABBREVIATIONS/NOTATIONS/GLOSSARY OF TERMS

APS	Ammonium peroxydisulphate
BIS	N, N'-Methylenbisacrylamide
BPB	Bromophenol blue
BSA	Bovine serum albumin
CTV	Citrus tristeza virus
DAS-ELISA	Double-antibody sandwich enzyme-linked immunosorbent assay
DE-52	Diethylaminoethyl cellulose
DNA	Deoxyribonucleic acid
hr	Hour
kDa	Kilodalton
μ l	Microliter
M	Molar
mg	Milligram
min	Minute
nm	Nanometer
OD	Optical density
PAGE	Polyacrylamide gel electrophoresis
PBS	Phosphate buffered saline
RNA	Ribonucleic acid
SDS	Sodium dodecyl sulfate
TEMED	N, N, N, N, -Tetramethylethylenediamine
Tris-HCl	Tris-(hydroxymethyl)-aminomethane hydrochloric acid



Alkaline phosphatase	An enzyme which hydrolyses certain phosphate-containing compounds under alkaline conditions; commonly obtained from calf intestine mucosa.
Antibody	A protein formed in blood serum in response to stimulation by an antigen. Antibodies are specific for their respective antigens, and antigens and antibodies are mutually attracted.
Antigen	A substance which, when introduced into the biological environment of a vertebrate animal, leads to the formation of antibodies directed specifically against it. All immunogens are antigenic, but not all antigens are immunogenic.
Antigenic Determinant	The small site on the antigen to which antibody is specifically able to become attached as determined by structural complementarity between antibody and antigen molecules.
Antiserum	Serum from any animal containing antibodies to a specified antigen.
Conjugate	The product of joining two or more dissimilar molecules by covalent bonds. In immunological contexts, one is usually a protein and the other either a hapten or a label such as fluorescein, ferritin, or enzyme.
Dialysis	A procedure using a membrane to separate various components in solution in accordance with their ability to pass through the membrane
Epitope	An antigenic determinant of defined structure, e.g., an identified oligosaccharide, or a chemical hapten.
Freund's Adjuvant.	A mixture of mineral oil and lanolin that enhances immune responses when emulsified with antigen for immunization. Freund's complete adjuvant includes killed mycobacteria; Freund's incomplete does not.

Immunization	Administration of antigen to an animal so as to produce an immune response to that antigen.
Immunogen	A substance that elicits an immune response when introduced into the tissues of an animal. To stimulate a response, immunogens must normally be foreign to the animal to which they are administered, of a molecular weight greater than 1,000, and of protein or polysaccharide nature.
Immunoglobulin (Ig).	Serum globular glycoprotein. There are five classes of immunoglobulin, IgA, IgD, IgE, IgG, and IgM. IgG is the major immunoglobulin class in the serum of man and in most species from amphibians upwards.
Polyclonal antibody	An heterologous antibody population derived from many clones.

CHAPTER 1

INTRODUCTION

1.1 Economic Importance of Citrus

Citrus is one of the most important fruit crops grown in all continents of the world. It contributes to the nourishment and refreshment of the people. Citrus products and by-products provide the basis for local agricultural industries, generate employment, raise income and, in many cases, constitute an important source of foreign revenue to developing and developed countries. Citrus production developed significantly during the last decade. From 1985 to 1995 world production increased at approximately 66 % from 47.8 Mt to 79.3 Mt. Oranges occupy the major part of the market, followed successively by mandarins, lemons, pomelos and other types (Table 1.1).

Table 1.1: Trend of citrus production from 1985 to 1995.

Citrus Types	Production campaigns			
	1985-1986		1995-1996	
	MT	%	MT	%
Oranges	31.0	65.7	58.4	73.0
Mandarins	7.9	16.7	8.8	11.0
Lemons	4.3	9.1	6.9	8.0
Pomelos and Grapes	4.0	8.5	5.2	8.0
Total	47.8	100.0	79.3	100.0

Source: FAO (1996)

MT: Metric tones



In Malaysia citrus has been cultivated since late 19th century and it was reported to be an economically viable plantation crop in the 1920's (Bunting, 1929). In the 1960's, citrus cultivation was popular among small land holders but diseases and pests caused a decline resulting in many growers abandoning their citrus plantings. Thus, citrus production decreased from 3670 ha in 1970 to 2487 ha in 1985. In 1995, the area cultivated throughout Peninsular Malaysia surpassed that of 1975 with 4291 ha planted with citrus. Out of these 1820 ha were planted with pomelo, 1295 ha with mandarins and the remaining 1176 ha consisting mainly of varieties of indigenous types such as sweet orange, limes and citrons (Table 1.2). With the increasing population, it is projected that Malaysia would need at least 12,500 ha of citrus to satisfy local demand (Ko, 1996).

Among commercial citrus varieties grown, oranges (*Citrus sinensis* L. Osb.), tangerines (*C. sinensis* x *C. reticulata*.), mandarins (*C. reticulata* Blanco), lemons (*C. limon* L. Burmf.), limes (*C. aurantifolia* Christm. Swing.), pomelos (*C. grandis* L. OSB.), and grapefruits (*C. paradisi* Macf.) are widely grown. Even though grapefruit was probably among the earliest citrus type introduced by the Department of Agriculture (Hawson, 1958), pomelos and especially mandarins were the most popular variety cultivated commercially. Limau Langkat (*Citrus suhuiensis* Hort. ex Tan), known locally as the popular loose peel citrus, can produce yields of up to 35 tones/ha/yr. yielding a gross income of RM 52,000. The recent introduction