



**UNIVERSITI PUTRA MALAYSIA**

**CROSSBREEDING OF THREE IRANIAN SHEEP BREEDS WITH  
RESPECT TO REPRODUCTIVE, GROWTH AND CARCASS  
CHARACTERISTICS**

**MOHAMMAD REZA KIYANZAD**

**FP 2002 37**

**CROSSBREEDING OF THREE IRANIAN SHEEP BREEDS WITH  
RESPECT TO REPRODUCTIVE, GROWTH AND  
CARCASS CHARACTERISTICS**

**By**

**MOHAMMAD REZA KIYANZAD**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia  
in Fulfilment of the Requirement for the Degree of Doctor of Philosophy**

**February 2002**



## *Dedication*

*To my parents, my dear wife and my children for their moral supports and encouragement.*



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**CROSSBREEDING OF THREE IRANIAN SHEEP BREEDS WITH  
RESPECT TO REPRODUCTIVE, GROWTH AND  
CARCASS CHARACTERISTICS**

**By**

**MOHAMMAD REZA KIYANZAD**

**February 2002**

**Chairman: Dr. Jothi Malar Panandam**

**Faculty: Agriculture**

In Iran, increasing demand for red meat, caused by increasing human population, pasture limitations and increasing cost of production, has become an impetus for sheep producers to change the system of farming and optimise meat production for more efficient output per unit livestock. One way to achieve the latter goal is by taking advantage of the effect of heterosis from crossbreeding.

Twenty seven rams and 473 ewes of three popular breeds of sheep in Iran, namely the Chal, Moghani and Zel, were crossed reciprocally. The reproductive performances of the ewes, and the growth performance, feed conversion ratio and carcass traits of the offspring, under an intensive system of management, were studied.

Zel ewes showed higher ( $P < 0.05$ ) overall apparent fertility, reproductivity and productivity compared Chal and Moghani ewes. However, when the problem



encountered by Zel rams in impregnating the bigger and fat-tailed breeds was taken into consideration, the breeds did not differ in ewe productivity and reproductivity. Zel ewes had relatively high pre-weaning lamb mortality, reducing their ewe efficiency.

Zel lambs had lower body weights and growth rates than Chal and Moghani. Their carcass had significantly ( $P<0.05$ ) lower total dissectible fat and more lean meat and bone percentages. Lambs sired by Chal and Moghani were heavier than those sired by Zel. Feed conversion ratio was not affected by crossbreeding. Chal, Moghani and Chal-Moghani crossbreeds showed lower prime-cut ( $P<0.05$ ) and higher fat-tail percent than the other breed groups. ZC (Zel ♂ x Chal ♀) had lower ( $P<0.05$ ) weaning and finishing weights than CC (Chal ♂ x Chal ♀). Their carcass fat percentages did not show significant difference. ZM (Zel ♂ x Moghani ♀) had significantly ( $P<0.05$ ) lower birth, weaning and finishing weights than MM (Moghani ♂ x Moghani ♀). MM, however, had higher carcass percent, but lower lean meat percent.

Zel ewes mated by Chal rams showed 26% higher ( $P<0.05$ ) ewe efficiency than those mated by Zel. CZ (Chal ♂ x Zel ♀) had heavier weaning and finishing weights and more fat percentage in fat-tail but less intramuscular and subcutaneous fat percentages compared to ZZ (Zel ♂ x Zel ♀). Carcass lean meat and fat percentages were similar. Zel ewes mated to Moghani rams weaned 28.5% more kilograms of lambs than those mated to Zel rams. Weaning and finishing weights of

MZ (Moghani ♂ x Zel ♀) were heavier than ZZ, but they had higher carcass fat percentages.

The males were, generally, heavier ( $P < 0.05$ ) than females from birth to finishing. Twins did not differ from singles for finishing weight. Females showed significantly ( $P < 0.05$ ) higher prime-cut and lower fat-tail percentages. Lean meat, fat and bone percents were not affected by sex or litter size.

It may be concluded that it would be beneficial to cross Chal rams with Zel ewes to produce CZ commercial lambs. However, to increase the ewe efficiency and to reduce their pre-weaning lamb mortality, supplemented feeding during pre-weaning period should be practised. Lambs of both sexes may be used in the feedlot.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doctor of Falsafah

**PEMBIAKBAKAAN KACUK TIGA BAKA BEBIRI IRAN  
MERUJUK KEPADA CIRI PEMBIAKAN, TUMBESARAN  
DAN KARKAS**

Oleh

**MOHAMMAD REZA KIYANZAD**

Februari 2002

**Pengerusi: Dr. Jothi Malar Panandam**

**Fakulti: Pertanian**

Di Iran, peningkatan permintaan untuk daging merah, disebabkan pertambahan populasi manusia, limitasi pastura dan peningkatan kos pengeluaran, telah menjadi satu pencetus kepada penternak-penternak bebiri untuk menukar sistem perladangan dan mengoptimasikan pengeluaran daging dengan pengeluaran yang lebih efisien daripada setiap unit ternakan. Salahsatu jalan untuk mencapai matlamat yang kedua ialah dengan menggunakan kesan heterosis hasil pembiakbakaan kacuk.

Dua puluh tujuh bebiri jantan dan 473 bebiri betina daripada tiga baka yang popular di Iran, iaitu Chal, Moghani dan Zel, telah dikacukkan secara silang. Prestasi pembiakan bebiri betina, dan prestasi tumbesaran, kadar penukaran makanan dan ciri-ciri karkas anak bebiri, di bawah sistem pengurusan intensif, telah dikaji.

Bebiri betina Zel telah menunjukkan kesuburan tampak keseluruhan, reproduktiviti dan produktiviti yang lebih tinggi ( $P < 0.05$ ) berbanding dengan bebiri betina Chal dan Moghani. Walau bagaimanapun, apabila masalah yang dihadapi oleh

bebiri jantan Zel membuntingkan baka yang lebih besar dan dengan ekor-lemak diambilkira, baka-baka tersebut tidak berbeza bagi produktiviti dan reproduksi betina. Bebiri betina Zel mempunyai mortaliti anak bagi tempoh pra-sapuh yang tinggi secara relatif, mengurangkan kecekapan mereka.

Anak bebiri Zel mempunyai berat badan dan kadar tumbesaran yang lebih rendah daripada Chal dan Moghani. Karkas mereka mengandungi lemak yang boleh diasingkan yang lebih rendah secara bererti ( $P < 0.05$ ) dan peratus daging lembut dan tulang yang lebih. Anak bebiri dengan Chal atau Moghani sebagai induk jantan adalah lebih berat daripada yang mempunyai Zel sebagai induk jantan. Kadar penukaran makanan tidak dipengaruhi pembiakbakaan kacuk. Chal, Moghani dan kacukan Chal-Moghani menunjukkan potongan prima yang lebih rendah ( $P < 0.05$ ) dan peratus lemak ekor yang lebih tinggi daripada kumpulan anak bebiri yang lain. ZC (Zel ♂ x Chal ♀) mempunyai berat sapuh dan berat akhir yang lebih kurang ( $P < 0.005$ ) daripada CC (Chal ♂ x Chal ♀). Peratus lemak karkas mereka tidak berbeza secara bererti. ZM (Zel ♂ x Moghani ♀) mempunyai berat lahir, berat sapuh dan berat akhir yang lebih rendah daripada MM (Moghani ♂ x Moghani ♀). MM mempunyai peratus karkas yang lebih tinggi, tetapi peratus daging lembut yang lebih rendah.

Zel betina yang dikawankan dengan jantan Chal menunjukkan kecekapan betina yang 26% lebih tinggi ( $P < 0.005$ ) daripada yang dikawankan dengan jantan Zel. CZ (Chal ♂ x Zel ♀) mempunyai berat sapuh, berat akhir dan peratus lemak pada ekor-lemak yang lebih tinggi tetapi kurang peratus lemak intramaskular dan



subkutanos berbanding dengan ZZ (Zel ♂ x Zel ♀). Peratus daging lembut dan lemak adalah serupa. Bebiri betina Zel yang dikawankan dengan jantan Moghani menyisipihkan 28.5% lebih kilogram anak berbanding dengan yang dikawankan dengan jantan Zel. Berat sapih dan berat akhir MZ (Moghani ♂ x Zel ♀) lebih berat daripada ZZ, tetapi mereka mempunyai peratus lemak pada karkas yang lebih tinggi.

Jantan, secara umumnya, adalah lebih berat ( $P < 0.05$ ) daripada betina sejak lahir hinggalah berat akhirnya. Anak kembar tidak berbeza ( $P < 0.05$ ) daripada anak tunggal untuk berat akhir. Betina menunjukkan potongan prima yang tinggi dan peratus lemak ekor yang rendah secara bererti. Peratus daging lembut, lemak dan tulang adalah tidak dipengaruhi oleh jantina atau saiz perindukan.

Bolehlah disimpulkan bahawa kacukan antara bebiri jantan Chal dan betina Zel untuk menghasilkan anak bebiri komersial CZ akan membawa manfaat. Walau bagaimanapun, untuk meningkatkan kecekapan betina Zel dan mengurangkan kematian anak bebiri pada tempoh pra-sapuh, pemberian makanan tambahan semasa tempoh pra-sapuh patut diamalkan. Anak bebiri kedua-dua jantina boleh digunakan dalam fidlot.

## ACKNOWLEDGEMENTS

All praise to Almighty Allah, the Merciful and the Benevolent. Had it not been due to His will and favour, the completion of this study would not have been possible.

I would like to express my sincere gratitude and appreciation to my supervisor, Dr. Jothi Malar Panandam for her invaluable advice, guidance, support, and patience and kind encouragement throughout the duration of this study.

I wish to express my sincere gratitude to my co-supervisors, Associate Professor Dr. Nasser Emamjomeh Kashan, Associate Professor Dr. Zainal Aznam Mohd Jelani and Professor Dr. Dahlan Ismail for their kind encouragement, guidance and sincere constructive comments.

Thanks are also due to Dr. Ali Kamali, Director of Animal Science Research Institute of Iran, Mr. Manochehr Monem, Dr. Ardeshir Nejati Javaremi for their advises, comments, support and friendly relationship.

I am especially grateful to the Ministry of Jihad-e-Sazandgi of Iran for providing the scholarship for my study.



I would like also to thank the skilful assistance by the staffs and workers of the Sheep and Goat Section of the Animal Science Research Institute of Iran for their kind hospitality during the breeding of the animals.

Most of all, I would like to acknowledge the support, understanding, and sacrifices of my dear wife and beloved children, Zahra, Sepideh and Mohammad, when I was unable to give them my attention.

Last, but not least, I would like to thank all other individuals who have contributed directly or indirectly to the successful completion of my study.



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement of the degree of Doctor of Philosophy.

---

AINI IDRIS, Ph.D.  
Professor  
Dean of School of Graduate Studies,  
Universiti Putra Malaysia

Date:



## TABLE OF CONTENTS

	Page
DEDICATION .....	ii
ABSTRACT .....	iii
ABSTRAK .....	v
ACKNOWLEDGEMENTS .....	vii
APPROVAL SHEETS .....	xi
DECLARATION FORM .....	xiii
LIST OF TABLE.....	xvi
LIST OF FIGURES.....	xix
LIST OF ABBREVIATIONS .....	xxi
 <b>CHAPTER</b>	
 1 INTRODUCTION.....	 1
1.1 Objectives of the Study .....	2
1.2 Research Hypothesis .....	3
1.3 Significance of the Study.....	3
 2 LITERATURE REVIEW.....	 5
2.1 Sheep Population and Mutton Production in Iran .....	5
2.1.1 Sheep Farming in Iran .....	6
2.1.2 Three Sheep Breeds of Iran.....	8
2.2 Economic Traits in Meat Sheep.....	10
2.2.1 Ewe Reproductive Performance.....	10
2.2.2 Birth Weight .....	15
2.2.3 Body Weight and Growth Rate .....	18
2.2.4 Feed Conversion Ratio .....	22
2.2.5 Body Conformation and Carcass Traits .....	23
2.3 Carcass Composition in Iranian and Exogenous Breeds .....	25
2.4 Fat Composition in Sheep .....	28
2.5 Crossbreeding .....	32
2.5.1 Heterosis.....	35
2.5.2 Genetic Basis for Heterosis .....	38
 3 MATERIALS AND METHODS.....	 39
3.1 Experimental Stocks and Breeding Design .....	39
3.2 General Management.....	40
3.2.1 Housing .....	40
3.2.2 Feeding .....	41
3.2.3 Hygiene and Health.....	44



3.3	Traits Studied.....	44
	3.3.1 Ewe Weights .....	44
	3.3.2 Ewe Reproductivity.....	44
	3.3.3 Growth Performance .....	47
	3.3.4 Body Measurements.....	48
	3.3.5 Growth of Animals in Feedlot.....	49
	3.3.6 Lamb Mortality .....	50
	3.3.7 Slaughter and Carcass Measurements .....	50
3.4	Statistical Analysis.....	55
	3.4.1 Statistical Model for Traits Measured.....	55
	3.4.2 Crossbreeding Parameters.....	59
4	RESULTS.....	61
4.1	Ewe Weights .....	61
4.2.	Reproductive Performance .....	62
	4.2.1 Apparent Fertility .....	62
	4.2.2 Prolificacy .....	62
	4.2.3 Litter Size .....	67
	4.2.4 Litter Weight .....	67
	4.2.5 Ewe Reproductivity.....	68
	4.2.6 Ewe Productivity .....	69
	4.2.7 Rearing Ability.....	69
	4.2.8 Ewe Efficiency .....	69
4.3	Growth Performance.....	70
	4.3.1 Birth Weight.....	70
	4.3.2 Weaning Weight.....	74
	4.3.3 Finishing Weight.....	80
4.4	Body Conformation .....	86
	4.4.1 Body Length.....	86
	4.4.2 Heart Girth.....	88
	4.4.3 Height at Wither .....	91
4.5	Feed Conversion Ratio.....	93
4.6	Lamb Mortality.....	95
4.7	Slaughter and Carcass Characteristics .....	96
	4.7.1 Carcass Measurements .....	104
	4.7.2 Prime-cut and Fat Content.....	107
	4.7.3 Chemical Composition of Carcass .....	110
4.8	Phenotypic Correlation.....	112
4.9	Regression Equations.....	114
4.10	Crossbreeding Parameters.....	115
5	DISCUSSION.....	119
5.1	Ewe Reproductive Performance .....	119
5.2	Growth Performance.....	123
	5.2.1 Body Weights .....	123



5.2.2	Body Conformation.....	128
5.2.3	Feed Conversion Ratio .....	129
5.2.4	Lamb Mortality .....	129
5.2.5	Carcass Characteristics.....	130
5.3	Predicting Live Weights and Carcass Components .....	133
5.4	Limitation of Study .....	135
5.5	Future Studies.....	135
6	GENERAL DISCUSSION AND CONCLUSION .....	137
	REFERENCES .....	143
	APPENDICES .....	153
	BIODATA OF THE AUTHOR.....	185



## LIST OF TABLES

Table		Page
2.1	Livestock population of Iran in the last 30 years.....	5
2.2	Production, importation and per capita consumption of red meat in Iran.....	6
2.3	Some characteristics of the Moghani, Chal and Zel sheep breeds...	9
2.4	Mean birth weight (kg) of some purebreed and crossbreed sheep..	17
2.5	Body weight (kg) of some purebreed and crossbreed sheep.....	20
2.6	Means and standard deviations of live measurements on ten Iranian fat-tail sheep breeds.....	24
2.7	Carcass composition of some Iranian fat-tailed sheep breeds.....	27
2.8	Least square means and standard deviations of total saturated and unsaturated fatty acid percentage and unsaturated/ saturated fatty acid ratio in three Iranian sheep breeds.....	30
2.9	Relative performance of <i>in-vivo</i> techniques and suitability for various uses.....	31
3.1	Sample size of each breed used in study.....	39
3.2	Mating plan used in the project.....	40
3.3	Ingredients and nutrient contents in diet of ewes .....	42
3.4	Dry matter (kg/day) offered to ewes of different body weight and stages of production .....	43
3.5	Ingredients and nutrient contents in lamb diet.....	43
3.6	Number of animals used in feedlot trial.....	49
3.7	Description of live and carcass traits measured .....	54
4.1	Least square analysis of variance for ewe reproductive traits.....	63
4.2	Means of reproductive performance of the three ewe breeds.....	66





4.3.	Least square means and standard errors of litter size and litter weight .....	68
4.4	Least square analysis of variance for birth weight of the lambs.....	71
4.5	Least square means and standard errors of lamb birth weight (kg)..	73
4.6	Least square analysis of variance for weaning weight.....	74
4.7	Least square analysis of variance for pre-weaning daily gain.....	77
4.8	Least square analysis of variance for finishing weight.....	80
4.9	Least square analysis of variance for average daily gain during feedlot period.....	83
4.10	Least square analysis of variance for body length.....	86
4.11	Least square means and standard errors of body length (cm).....	87
4.12	Least square analysis of variance for lamb heart girth.....	89
4.13	Least square means and standard errors of heart girth (cm) .....	90
4.14	Least square analysis of variance for lamb height at wither.....	91
4.15	Least square means and standard errors of height at wither (cm)...	92
4.16	Least square analysis of variance for feed conversion ratio.....	94
4.17	Lamb mortality with respect to dam breed and lamb breed group.	95
4.18	Least square analysis of variance for slaughter and carcass traits...	97
4.19	Least square means and standard errors for slaughter and carcass traits (kg).....	98
4.20	Least square analysis of variance for carcass component percentages.....	100
4.21	Least square means and standard errors of offal, carcass, meat, fat and bone percentages.....	102
4.22	Least square analysis for carcass measurements.....	105



4.23	Least square means and standard errors of carcass measurements..	106
4.24	Least square analysis of variance for prime-cuts, fat-tail, intramuscular fat and subcutaneous fat percents.....	107
4.25	Least square means and standard errors of prime-cuts, fat-tail, intramuscular and subcutaneous fat percentages .....	109
4.26	Least square analysis of variance for moisture, protein, chemical fat and ash percentages.....	110
4.27	Least square means and standard errors of chemical compositions of carcass.....	111
4.28	Estimates of heterosis (amount and percentage) for growth and carcass traits.....	116



## LIST OF FIGURES

Figure.		Page
3.1	Measurements on the live sheep.....	49
3.2	Longissimus dorsi sizes.....	52
3.3	Sites of cutting of lamb carcass and the parts produced .....	53
4.1	Ewe weights at purchase, before mating and after parturition.....	61
4.2.	Least square means and standard errors of lamb weaning weight	76
4.3	Mean pre-weaning average daily gain of the lambs.....	78
4.4.	Means of pre-weaning body weights of lambs.....	78
4.5	Overall means and standard errors of finishing weights for the nine lamb breed groups.....	82
4.6.	Means of body weights during feedlot period.....	82
4.7.	Means and standard errors of average daily gain during feedlot period for the nine lamb breed groups.....	85
4.8.	Means of average daily gain during feedlot period for the nine lamb breed groups.....	85
4.9.	Least square means of feed conversion ratios in the nine lamb breed groups.....	94



## LIST OF ABBREVIATIONS

A	Maximum width of the longissimus muscle at cross-section posterior to the last rib
B	Maximum depth of the longissimus muscle at cross-section posterior to the last rib
BL1	Body length at birth
BL2	Body length at weaning
BL3	Body length at finishing
BW	Birth weight
C	Chal
C <sub>1</sub>	Subcutaneous fat thickness over the point of greatest depth of the longissimus muscle posterior to the last rib
Ca	calcium
CC	Chal (♂) x Chal (♀)
CCW	Cold carcass weight
CF	Chemical fat
CM	Chal (♂) x Moghani (♀)
C-M	Reciprocal crosses of Chal and Moghani breeds
CZ	Chal (♂) x Zel (♀)
C-Z	Reciprocal crosses of Chal and Zel breeds
DE	Digestible energy
df	degree of freedom
DM	dry matter
EBW	Empty body weight

EE	Ewe efficiency
EE <sub>MWE</sub>	Ewe efficiency based on metabolic weight of ewe exposed
EE <sub>MWL</sub>	Ewe efficiency based on metabolic weight of lambed ewes
F <sub>1</sub>	first filial generation
FADG	Feedlot average daily gain
FCR	Feed conversion ratio
FW	Finishing weight
GR	True depth of soft tissue over 12 <sup>th</sup> rib, 11 cm from the dorsal midline on intact cold carcass.
HCW	Hot carcass weight
HG1	Heart girth at birth
HG2	Heart girth at weaning
HG3	Heart girth at Finishing
HW1	Height at wither at birth
HW2	Height at wither at weaning
HW3	Height at wither at finishing
M	Moghani
MC	Moghani (♂) x Chal (♀)
ME	Metabolizable energy
MM	Moghani (♂) x Moghani (♀)
MS	mean square
MZ	Moghani (♂) x Zel (♀)
M-Z	Reciprocal crosses of Moghani and Zel breeds
NFE	Nitrogen free extract

NGR	Measurements of GR with a 22-gauge hypodermic needle over the 12 <sup>th</sup> rib, 12-cm from the dorsal midline.
P	phosphorus
PADG	Pre-weaning average daily gain
SS	sum of square
SW	Slaughter weight
UGR	Ultrasonic Measurements of GR
WW	Weaning weight
Z	Zel
ZC	Zel (♂) x Chal (♀)
ZM	Zel (♂) x Moghani (♀)
ZZ	Zel (♂) x Zel (♀)

# CHAPTER 1

## INTRODUCTION

In Iran, red meat is one of the common sources of protein. The average red meat consumption per capita is 14 kg (Kiyanzad and Monem, 1999). The main sources of red meat are cattle, buffalo, sheep and goat. Meat from these sources, however, is not sufficient to meet the demand.

Sheep population in Iran is 50 million, comprising of 26 genetic groups. Meat production by sheep and goat amounts to 57% of the total red meat production in the country (FAO, 1998; Osfoori and Fesus, 1996). Sheep are kept for meat, wool, milk and pelt. Although there has been an increase in production over the years, the country aims only not for self-sufficiency, but also to be able to export mutton (Kiyanzad and Monem, 1999; FAO, 1997; Yalcin, 1979).

Periodical drought, overgrazing, lack of management of pasture, use of pasture plants by clans and villagers as fuel and use of pasture area for crop cultivation over the last four decades has caused the pasture capacity to be reduced. On the other hand, increasing demand for red meat, limitation faced by animal food producers (land use for other purposes, climatic conditions, etc.) and increasing cost of production, has become an impetus for livestock producers to change the system of production. The traditional sheep keeping practices are no longer viable or economical. It has become a necessity to adopt the intensive or semi-intensive system of farming and to use more productive animals or breeds so that sheep

