



**UNIVERSITI PUTRA MALAYSIA**

**CHARACTERIZATION, PURIFICATION AND ANTISERUM  
PRODUCTION OF SWEET POTATO FEATHERY  
MOTTLE POTYVIRUS (SPFMV)**

**KHAIRULMAZMI AHMAD**

**FP 2003 26**

**CHARACTERIZATION, PURIFICATION AND ANTISERUM PRODUCTION OF  
SWEET POTATO FEATHERY MOTTLE POTYVIRUS (SPFMV)**

**By**

**KHAIRULMAZMI AHMAD**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra  
Malaysia, in Fulfilment of the Requirements for the Degree of  
Master of Agricultural Science**

**August 2003**



# DEDICATION

Special dedication to my beloved..

Father..

Mother...

&

To all my family members..

Lastly

To all my best friends



Abstract of thesis presented to Senate of Universiti Putra Malaysia in Fulfilment  
of the Requirements for the Degree of Master of Agricultural Science

**CHARACTERIZATION, PURIFICATION AND ANTISERUM PRODUCTION OF  
SWEET POTATO FEATHERY MOTTLE POTYVIRUS (SPFMV)**

By

**KHAIRULMAZMI BIN AHMAD**

**August 2003**

**Chairman: Inon Sulaiman, Ph.D.**

**Faculty: Agriculture**

Characterization, purification and production of antiserum against sweet potato feathery mottle potyvirus (SPFMV) were carried out in this study. Characterization of the virus involved host range reaction, inclusion bodies determination, virion morphology and determination of coat protein molecular mass.

Initial serological detection of potyviruses infection was carried-out by Indirect ELISA using an Agdia® commercial kit for potyvirus on samples of sweet potato collected from major growing areas in Peninsular Malaysia, Sabah and Sarawak. Experimental results of this study showed, potyviruses are present in most of the



major sweet potato growing areas in Peninsular Malaysia and some areas in Sabah and Sarawak.

The SPFMV isolate used in this study was single probed by aphid and vectored to healthy *Ipomoea setosa* three times. The initial infection symptom on *I. setosa* was characterized by veinal chlorosis of actively growing leaves. In subsequent leaves, the veinal chlorosis was concentrated to major veins. Progressively the veinal chlorosis was restricted to midvein. Subsequent new foliar growth was symptomless.

Host reaction results showed, *I. nil* produced symptoms of faint vein clearing and crinkling after inoculation. Symptomless *I. purpurea* tested positive in Indirect ELISA, thus showing latent infection in this host. The rest of the indicator plants did not show any virus symptom and tested negative in the ELISA test. These results indicated that the virus isolate in this study has a narrow host range and restricted to the *Convolvulaceae* family only.

Cytoplasmic inclusions (CIs), a major characteristic of the poty group, were observed under light microscopy. In the infected cells, the CIs were stained dark blue (Azure A) and were usually located beside the plant cell nuclei.

Purification of SPFMV directly from infected sweet potato leaves by twice CsCl density gradient centrifugation yielded about 1.5 mg of virus per 100 g of fresh

tissue. The ratio of  $A_{260}/A_{280}$  nm was around 1.123 (without correction for light scattering). Electron microscopy revealed flexuous filamentous particles with a modal length of 750-849 nm. Protein analysis of denatured virus by SDS-PAGE revealed a major coat protein of 40 kDa.

Polyclonal antiserum in this study was produced against the purified local SPFMV isolate. The antiserum had a titer of 1:1536 in microprecipitin tests. The antibody reacted positively by DAS-ELISA with sap from SPFMV-field infected tissue and purified SPFMV preparation.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains Pertanian

**PENCIRIAN, PENULENAN DAN PENGHASILAN ANTISERUM TERHADAP  
SWEET POTATO FEATHERY MOTTLE POTYVIRUS (SPFMV)**

**Oleh**

**KHAIRULMAZMI BIN AHMAD**

**Ogos 2003**

**Pengerusi: Inon Sulaiman, Ph.D.**

**Fakulti: Pertanian**

Pencirian, penulenan dan penghasilan antiserum terhadap sweet potato feathery mottle potyvirus (SPFMV) telah dijalankan dalam kajian ini. Kajian pencirian melibatkan reaksi virus terhadap julat perumah, penentuan "inclusions bodies", morfologi virion dan penentuan berat molekul sarung-protein virus.

Ujian awal serologi terhadap jangkitan potyvirus telah dijalankan dengan menggunakan kaedah Indirect ELISA (Agdia® commercial kit for potyvirus) terhadap sampel-sampel keledak yang telah diambil daripada kawasan penanaman utama keledak di Semenanjung Malaysia, Sabah dan Sarawak. Keputusan ujian menunjukkan bahawa potyvirus telah menjangkiti kebanyakan



tanaman dikawasan penanaman utama keledak di Semenanjung Malaysia dan beberapa kawasan di Sabah dan Sarawak.

SPFMV dipencilkan melalui satu 'probe' vektor afid dan dijangkitkan kepada *Ipomoea setosa* sihat sebanyak tiga kali. Ciri simptom jangkitan awal pada *I. setosa*, adalah dengan pembentukan simptom urat jernih pada urat-urat daun yang masih aktif membentuk. Pada daun yang berikutnya, simptom urat jernih hanya tertumpu pada urat-urat utama daun. Seterusnya, simptom urat jernih hanya tertumpu pada urat tengah daun sahaja. Selepas itu, daun yang baru tidak lagi menunjukkan simptom.

Keputusan reaksi perumah, *I. nil* selepas diinokulasi adalah pembentukan simptom urat jernih yang samar dan daun berkedut. Tiada simptom jangkitan pada perumah *I. purpurea*, tetapi mencatat keputusan positif dalam ujian Indirect ELISA. Ini menunjukkan berlakunya jangkitan terpendam pada perumah tersebut. Perumah yang lain tidak menunjukkan simptom dan ia juga mencatat keputusan negatif terhadap ujian Indirect ELISA. Keputusan ujikaji ini menunjukkan bahawa virus yang telah dipencilkan mempunyai julat perumah yang terhad kepada tanaman jenis *Convolvulaceae* sahaja.

"Cytoplasmic inclusion"(CIs) iaitu pencirian utama kumpulan poty, telah dikesan melalui mikroskop cahaya. Pada daun berpenyakit, CIs dapat dibezakan apabila



ia mengambil warna biru gelap (Azure A) dan biasanya terdapat berdekatan dengan nukleus sel tumbuhan.

Penulenan SPFMV secara langsung dari daun keledek berpenyakit dengan menggunakan penggunaan emparan "CsCl density gradient" sebanyak dua kali telah menghasilkan 1.5 mg virus daripada 100 g daun segar. Manakala nisbah  $A_{260}/A_{280}$  nm adalah pada sekitar 1.123 (tanpa pembetulan terhadap biasan cahaya). Keputusan mikroskop elektron pula menunjukkan partikal virus berbentuk rod melentur (flexuous rod) dengan panjang, 750-849 nm berdasarkan model panjang virus. Analisis protein virus menggunakan SDS-PAGE telah menghasilkan satu jalur major sarung-protein yang berat molekulnya ialah 40 kDa.

Antiserum poliklonal dalam kajian ini telah dihasilkan dengan menggunakan SPFMV tempatan yang telah dituliskan. Antiserum yang dihasilkan mempunyai titer sehingga 1:1536 pencairan dengan menggunakan ujian mikroprinsipitin. Antibodi yang dihasilkan telah memberi reaksi positif terhadap sap dan daun yang dijangkiti oleh SPFMV dan juga SPFMV tulen dengan menggunakan ujian DAS-ELISA

## ACKNOWLEDGEMENTS

First of all, I thank Allah S.W.T for giving me the strength and ability to complete this study. I am also sincerely grateful and greatly indebted to my supervisor Dr. Inon Sulaiman for her guidance, understanding, support, compassion and sacrifices. Her invaluable advice and comments are highly appreciated.

My sincere appreciation and gratitude to members of my supervisory committee, En. Zakaria Sidek and En. Yaakob Doon for their critical comments and suggestions during the course of this project.

I also thank to Mrs Junaina, all the microbiology laboratory staffs and staffs of the Graduate School of UPM for their kind assistance and cooperation throughout my study. Special thanks to Mr. Nurhadi and Mr. Asaad for their encouragement and support throughout my study.

My utmost gratitude to my father; Ahmad b. Abdullah, my mother; Zaliha bt. Mohammad, my sisters; Nor Muzalni bt. Ahmad, Nor Asmizan bt. Ahmad, Nor Hazalzhiah, my brother; Khairul Naim b. Ahmad and Noraini Enrico for their boundless support and love. Finally, I wish to express my sincere thanks to all those who have in one way or another helped me in making this study succeed.



I certify that an Examination Committee met on 6 August 2003 to conduct the final examination of Khairulmazmi b. Ahmad on his Master of Agricultural Science thesis entitled "Characterization, Purification and Antiserum Production of Sweet Potato Feathery Mottle Potyvirus (SPFMV)" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulation 1981. The Committee recommends the candidate be awarded the relevant degree. Members of the Examination Committee are follows:

**Sariah Meon, Ph.D.**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
( Chairman )

**Inon Sulaiman, Ph.D.**


Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
( Member )

**Zakaria Sidek**

Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
( Member )

**Yaakob Doon**

Research Officer  
Institut Penyelidikan dan Kemajuan Pertanian Malaysia (MARDI)  
43400 Serdang, Selangor.  
( Member )



---

**GULAM RUSUL RAHMAT ALI, Ph.D.**  
Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 01 DEC 2003

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Agricultural Science. The members of the Supervisory Committee are as follows:

**Inon Sulaiman, Ph.D.**  
Faculty of Agriculture  
Universiti Putra Malaysia  
( Chairman )

**Zakaria Sidek**  
Faculty of Agriculture  
Universiti Putra Malaysia  
( Member )

**Yaakob Doon**  
Institut Penyelidikan dan Kemajuan Pertanian Malaysia (MARDI)  
43400 Serdang, Selangor  
( Member )




---

**AINI IDERIS, Ph.D.**  
Professor/Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 16 DEC 2003

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that is has not been previously or concurrently submitted for any other degree at UPM or other institutions.



**KHAIRULMAZMI B. AHMAD**

Date: 30.11.2003

## TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF ABBREVIATIONS	xix
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 General	4
2.1.1 Sweet Potato Production	5
2.1.2 Sweet Potato Diseases	9
2.1.3 Viral Problems Of Sweet Potato In Malaysia	10
2.2 General Properties Of Potyvirus	10
2.2.1 Sweet Potato Feathery Mottle Potyvirus (SPFMV)	14
2.2.2 Sweet Potato Virus Disease (SPVD)	16
2.2.3 Other Potyviruses Affecting Sweet Potato	17
2.2.3.1 Sweet Potato Mild Mottle Virus (SPMMV)	17
2.2.3.2 Sweet Potato Latent Virus (SPLV)	18
2.2.3.3 Sweet Potato Yellow Dwarf Virus (SPYDV)	18
2.2.3.4 Sweet Potato Vein Mosaic Virus (SPVMV)	19
2.3 Economic Loss	19
2.4 Detection Methods For Sweet Potato Virus	21
2.4.1 Indicator Plants	22
2.4.2 Serological Tests	23
2.4.3 Transmission Electron Microscopy (TEM)	24
2.5 Virus Purification	25
2.5.1 Virus Extraction	27
2.5.2 Concentration Of The Virus	28
2.5.3 Density Gradient Centrifugation	29
2.6 Enzyme-Linked Immunosorbent Assay (ELISA)	30
2.6.1 Test Samples	32
2.6.2 Antiserum	32
2.6.3 Enzyme-Antibody Conjugate	33
2.6.4 Substrate	33



3	<b>MATERIALS AND METHODS</b>	35
3.1	Assessment Of The Potyvirus Distribution in Sweet Potato Growing Areas in Malaysia	35
3.2	Sources Of Virus Isolate	37
3.3	Isolate Characterization	38
3.3.1	Virus Isolation And Maintenance	38
3.3.2	Host Range Test	39
3.3.3	Light Microscopy Examinations for Inclusions Bodies	40
3.4	Virus Purification	40
3.5	Characterization Of Purified Virus	42
3.5.1	Spectrophotometry	42
3.5.2	Transmission Electron Microscopy (TEM)	42
3.5.3	Estimation of Molecular Mass of Coat Protein	43
3.6	Production Of Polyclonal Antiserum	44
3.6.1	Homologous Antiserum Titer Test	45
3.6.2	Immuno-Globulin Purity Determination Test	46
3.6.3	ELISA Procedures	47
3.6.3.1	Immuno-Globulin Purification	47
3.6.3.2	Alkaline Phosphatase Conjugation	48
3.6.4	Double Antibody Sandwich-ELISA (DAS-ELISA)	49
3.6.4.1	Optimization of DAS-ELISA Test	49
3.6.4.2	DAS-ELISA Test Using Produced Antiserum	51
4	<b>RESULTS AND DISCUSSION</b>	53
4.1	Assessment Of Potyvirus Distribution In Sweet Potato Growing Areas In Malaysia	53
4.1.1	Peninsular Malaysia	53
4.1.2	Sabah and Sarawak	55
4.2	Isolates Characterization	58
4.2.1	Virus Isolation and Maintenance	58
4.2.2	Host Range	61
4.2.3	Light Microscopy	66
4.3	Virus Purification	68
4.4	Characterization Of The Purified Virus	71
4.4.1	Spectrophotometry	71
4.4.2	Transmission Electron Microscopy (TEM)	73
4.4.3	Coat Protein Molecular Mass	77
4.5	Serology	80
4.5.1	Microprecipitin Test	80
4.5.2	IgG Purification	83
4.5.3	Purity Of IgG	84
4.5.4	Optimization of DAS-ELISA Tests	85
4.5.4.1	Optimization Of Coating IgG	85
4.5.4.2	Optimization Of Incubation Period	87

4.5.4.3 Sensitivity Of DAS-ELISA	89
4.5.5 DAS-ELISA Test Using The Produced Antiserum	91
5 CONCLUSION	93
REFERENCES	97
APENDICES	106
BIODATA OF THE AUTHOR	113





## LIST OF TABLES

Table	Page	
2.1	World sweet potato production, 1989	6
2.2	Hectareage of land used for sweet potato in Peninsular Malaysia	8
2.3	Potyvirus proteins and their functions	12
3.1	List of sweet potato and <i>Ipomoea setosa</i> samples with their locations, symptoms and conditions for DAS-ELISA test	52
4.1	Results of Agdia® Indirect ELISA test on sweet potato samples collected from Terengganu, Johor and Kelantan	54
4.2	Results of Agdia® Indirect ELISA test on sweet potato samples collected from Perak, Kedah, Pahang and Kelantan	54
4.3	Results of Agdia® Indirect ELISA test on sweet potato samples collected from Sabah and Sarawak	57
4.4	Results of Agdia® Indirect ELISA test on sweet potato samples collected from Sarawak	57
4.5	Symptoms in plants mechanically inoculated with SPFMV 5 weeks after inoculation and ELISA result of Indirect-ELISA to detect latent infection	63
4.6	Determination of homologous antiserum titer and host protein specific antibodies by microprecipitin test	82
4.7	Purified IgG concentrations in different fractions of DE 52 column chromatography elution after ammonium sulphate precipitation and membrane dialysis	83
4.8	DAS-ELISA test against SPFMV field- infected sweet potato tissue, quantified at 405 nm 60 min. after incubation time.	92



## LIST OF FIGURES

Figures	Page
4.1 First stage symptom on <i>I. setosa</i> leaf two weeks after inoculation with SPFMV : vein clearing on actively growing leaf (arrow)	59
4.2 Second stage symptom on <i>I. setosa</i> leaf infected with SPFMV, 3 to 4 days after the appearance of the first stage : chlorotic banding of primary and secondary veins (arrow)	59
4.3 Third stage symptom on <i>I. setosa</i> leaf infected with SPFMV, 5 to 6 days after the appearance of the second stage symptom: leaf distortion (A) and vein banding of midrib (B)	60
4.4 <i>I. setosa</i> infected by SPFMV, 4 weeks after inoculation. The plant was severely stunted with deformed leaf (arrow)	60
4.5 Initial symptom on <i>I. nil</i> leaf infected with SPFMV, 8 to 9 days after sap inoculation: faint vein clearing on primary leaf restricted to major veins (arrow)	62
4.6 <i>I. nil</i> leaf infected with SPFMV: curling symptom on leaf as a result of epinasty	62
4.7 <i>I. nil</i> plant infected with SPFMV showing stunted symptom 5 weeks after inoculation. Healthy plant (left) and infected plant (right)	63
4.8 Epidermal cells of sweet potato infected with SPFMV. CIs stained dark blue with Azure A located beside the plant nuclei. S= Stomata, CI= Cytoplasmic inclusion and N= nucleus. (X400)	67
4.9 Epidermal cells of healthy sweet potato. CIs were absent from the epidermal cells. S= Stomata and N= nucleus. (X400)	67
4.10 Band of purified SPFMV (arrow) after the second centrifugation on step gradient CsCl. A single virus band located approximately 20 - 21 mm from the bottom of CsCl density gradient tube was seen	69
4.11 Ultraviolet absorption spectrum (230-320 nm) of purified preparation of SPFMV	72



4.12	Electron microscopy of purified SPFMV virions from CsCl step gradient negatively stained with 2% PTA. Bar represent 250 nm	75
4.13	Histogram of the length 100 purified SPFMV particles. The histogram indicated that the modal length of SPFMV particle length was 750 - 849 nm	76
4.14	SDS-PAGE analysis of SPFMV coat protein. The coat protein Molecular mass was estimated to be approximately 40 kDa. M= protein marker, S1 and S2=freshly purified SPFMV coat protein	78
4.15	Estimation of SPFMV by graphical comparison to standard markers migration distances. The migration distance of 3 cm (arrow) for SPFMV indicated that the molecular mass of the coat-protein is approximately 40 kDa	79
4.16	SDS-PAGE gel analysis: Lane A = major protein band (1) Agdia antiserum. Lane B= major protein band (1) and minor protein band (2) of purified IgG	84
4.17	Relationship between concentrations of coating IgG and OD value in DAS-ELISA test after 1 hour incubation period. SPFMV concentration: 500 µg/ml, enzymes conjugate: 1:500 dilutions	86
4.18	Effect of incubation times against OD value with 2 µg/ml of coating IgG, SPFMV concentration of 500 µg/ml and 1:500 dilution of enzyme labelled-IgG.	88
4.19	Effect of different concentrations of purified SPFMV with OD value in DAS-ELISA. SPFMV concentration: 500 µg/ml, Enzyme conjugate: 1:500 dilution: Coating IgG:2 µg/ml, Incubation period: 1 hour	90

## LIST OF ABBREVIATIONS

Adjuvant	Substance injected with antigens (usually mixed with them but sometimes given prior to or after the antigen) which non-specifically enhances or modifies the immune response to that antigen.
Alkaline phosphatase	An enzyme which hydrolyses certain phosphate-containing compounds under alkaline conditions; commonly obtained from calf intestine mucosa
Antibody	A protein formed in blood serum in response to stimulation by an antigen. Antibodies are specific for their respective antigens, and antigens and antibodies are mutually attracted
Antigen	A substance which, when introduced into the biological environment of a vertebrate animal, leads to the formation of antibodies directed specifically against it. All immunogens are antigenic, but not all antigens are immunogenic.
Antiserum	Serum from any animal containing antibodies to a specified antigen.
APS	Ammonium peroxydisulphate
BIS	N. N' –Methylenbisacrylamide
BPB	Bromophenol blue
BSA	Bovine serum albumin
Chromatography	The separation of mixtures of chemicals, compounds, proteins, macro-molecules etc. into their constituents or components by preferential adsorption by a solid such as a column of cellulose, or by filter paper or by gel
Conjugate	The product of joining two or more dissimilar molecules by covalent bonds. In immunological contexts, one is usually a protein and the other either a hapten or a label such as fluorescent, ferritin, or enzyme



Dialysis	A procedure using a membrane to separate various components in solution in accordance with their ability to pass through the membrane
DNA	Deoxyribonucleic acid
ELISA	Enzyme-Linked Immunosorbent Assay. An immunoassay system employing enzyme bound to antibody as the immunologic probe determining the extent of antigen-antibody reaction.
Immunogen	A substance that elicits an immune response when introduced into the tissues of an animal. To stimulate a response, immunogens must normally be foreign to the animal to which they are administered, of a molecular weight greater than 1,000, and of protein or polysaccharide nature.
Freund's Adjuvant.	A mixture of mineral oil and lanolin that enhances immune responses when emulsified with antigen for immunization. Freund's complete adjuvant includes killed mycobacteria; Freund's incomplete does not.
Immunoglobulin (Ig).	Serum globular glycoprotein. There are five classes of immunoglobulin, IgA, IgD, IgE, IgG, and IgM. IgG is the major immunoglobulin class in the serum of man and in most species from amphibians upwards.
kDa	Kilodalton
NC	Nitrocellulose membrane
OD	Optical density
PAGE	Polyacrylamide gel electrophoresis
PBS	Phosphate buffered saline
PBST	Phosphate buffered saline-Tween
PVP	Polyvinylpyrrolidone
RNA	Ribonucleic acid



SPFMV	Sweet potato feathery mottle potyvirus
SPCSV	Sweet potato chlorotic stunt crinivirus
SPVD	Sweet potato virus disease
SDS	Sodium dodecyl sulfate
TEMED	N, N ,N, N, -Tetramethylethyldiamine
Tris-HCl	Tris-(hydroxymethyl )-aminomethane



# CHAPTER 1

## INTRODUCTION

Sweet potato (*Ipomoea batatas* L.) is a dicotyledonous plant from the family *Convolvulaceae*. The crop originated from Central or South America (Karyeija et al., 1998b). It is grown exclusively in tropical and subtropical regions with low input and can produce good yield under marginal conditions (Farayi et al., 1997). It has been planted in Malaysia for more than 100 years and more than 100 varieties are available (Saad, 1994). Traditionally, during the time of food shortage, sweet potato became useful substitutes of the staple food crop, i.e. rice (Tan and Saad, 1994). This crop has the potential to become an important supplier of starch to the other industries.

As with other crops, sweet potato is also susceptible to plant pathogens. Sweet potato diseases, especially those caused by viral infections have been identified as the second most important biotic component after insect infestations (Moyer and Salazar, 1989 ; Farayi et al., 1997). Previous studies indicated that there are five major potyviruses affecting sweet potato production. One of these viruses is the sweet potato feathery mottle potyvirus (SPFMV), which has been reported to infect sweet potato wherever it is grown (Moyer and Salazar, 1989).



SPFMV has been detected by observing symptom expression patterns in infected plants in the field, as well as on sensitive indicator plants like *Ipomoea setosa* and *I. nil* in biological assays (Farayi et al., 1997 and Karyeija et al, 1998a). However, SPFMV indexing has been hampered by absence of rapid, sensitive and reliable diagnostic tool for viral pathogens (Farayi et al., 1997). Hence, an improved detection system is needed to detect and refine the diagnosis of SPFMV.

Double-antibody sandwich enzyme-linked immunosorbent assay (DAS-ELISA) introduced by Clark and Adams, (1977) was able to solve the limitation of the biological assay. The ELISA techniques are known to be sensitive, cheaper and faster than biological assay. ELISA techniques have a higher level of sensitivity and reproducibility, yet employs relatively simple and inexpensive equipments (Clark and Adams, 1977). Results of preliminary studies on viral problem of sweet potato in Malaysia, indicated the presence of putative SPFMV (Inon et al., 1998).

Information on the epidemiology, symptomatology, host range, virus-vector relationship and molecular aspect of the putative SPFMV are still incomplete or lacking. Important information is needed to develop strategies for SPFMV management. Research on sweet potato development in Malaysia is still lacking. Only two institutions (MARDI and UPM) are working on sweet potato problems (Saad, 1994). Most of the work reported has been in the



area of plant improvement and breeding. This study was conducted with the following objectives:

- 1) To assess the spread of the SPFMV infection in sweet potato in the major sweet potato growing areas in Malaysia.
- 2) To characterize a local SPFMV isolate.
- 3) To purify the local SPFMV isolate.
- 4) To produce local antiserum against local SPFMV isolate.