



**UNIVERSITI PUTRA MALAYSIA**

**POTENTIAL OF SELECTED ENTOMOPATHOGENIC  
HYPHOMYCETES FOR CONTROLLING THE RICE PESTS,  
SITOPHILUS ORYZAE (COLEOPTERA: CURCULIONIDAE) AND  
CORCYRA CEPHALONICA (LEPIDOPTERA: PYRALIDAE)**

**HENDRAWAN SAMODRA**

**FP 2003 29**



**POTENTIAL OF SELECTED ENTOMOPATHOGENIC HYPHOMYCETES  
FOR CONTROLLING THE RICE PESTS, *SITOPHILUS ORYZAE*  
(COLEOPTERA : CURCULIONIDAE) AND *CORCYRA CEPHALONICA*  
(LEPIDOPTERA : PYRALIDAE)**

**HENDRAWAN SAMODRA**

**MASTER OF SCIENCE  
UNIVERSITI PUTRA MALAYSIA  
2003**



**POTENTIAL OF SELECTED ENTOMOPATHOGENIC HYPHOMYCETES  
FOR CONTROLLING THE RICE PESTS, *SITOPHILUS ORYZAE*  
(COLEOPTERA : CURCULIONIDAE) AND *CORCYRA CEPHALONICA*  
(LEPIDOPTERA : PYRALIDAE)**

**By**

**HENDRAWAN SAMODRA**

**Thesis Submitted to the School of Graduate Studies,  
Universiti Putra Malaysia, in Fulfilment of the Requirements  
for the Degree of Master of Science**

**November 2003**



## DEDICATION

*For my parents Hendro Suyoko, my wife Takako, my daughter Aisha  
and my brother Danis*



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Master of Science

**POTENTIAL OF SELECTED ENTOMOPATHOGENIC HYPHOMYCETES FOR CONTROLLING THE RICE PESTS, *SITOPHILUS ORYZAE* (COLEOPTERA : CURCULIONIDAE) AND *CORCYRA CEPHALONICA* (LEPIDOPTERA : PYRALIDAE)**

By

Hendrawan Samodra

November 2003

**Chairman : Professor Yusof Bin Ibrahim, Ph.D**

**Faculty : Agriculture**

The rice weevil, *Sitophilus oryzae* (L.) and the rice moth, *Corcyra cephalonica* Stn. are serious stored grain pests worldwide. They attack a wide variety of stored products and are the two important pests of stored rice.

The use of entomopathogenic fungi (EF) is a novel approach to the control of insect pests of stored grains. The potential of dry conidia of three genera of EF, *Beauveria bassiana*, *Metarhizium anisopliae* and *Paecilomyces fumosoroseus* was examined in the laboratory against the adults of *S. oryzae* and the larvae of *C. cephalonica*.

The pathogenicities of nine selected isolates of the EF (BbGc, BbPs, BbPc, MaOrMaj, MaSc, MaGmC, MaOrMan, MaPs, PfPp) to *S. oryzae* adults and



*C. cephalonica* larvae were evaluated in the laboratory. All the isolates tested were pathogenic against *S. oryzae* adults and *C. cephalonica* larvae but pathogenicity varied among the isolates. Two isolates of *B. bassiana* (BbGc and BbPs) and one isolate of *M. anisopliae* (MaPs) were superior and caused high mortality against both these insects compared to other isolates. The median effective concentration (EC<sub>50</sub>) for isolates BbGc, BbPs and MaPs against *C. cephalonica* larvae were  $1.238 \times 10^6$ ,  $2.072 \times 10^6$ ,  $1.775 \times 10^6$  conidia g<sup>-1</sup> respectively. However, higher EC<sub>50</sub> values for these isolates were recorded against *S. oryzae* adult, namely  $9.491 \times 10^6$ ,  $1.377 \times 10^7$ ,  $1.120 \times 10^7$  conidia g<sup>-1</sup> respectively. Thus, it can be concluded that *C. cephalonica* larvae were more susceptible than *S. oryzae* adults to the three selected fungal isolates. The median lethal time (LT<sub>50</sub>) at concentration  $1 \times 10^9$  for these isolates against *C. cephalonica* larvae were 3.5 d for BbGc, 3.6 d for BbPs and 3.5 d for MaPs and against *S. oryzae* adults were 3.1d for BbGc, 3.3 d for BbPs and 2.5 d for MaPs.

The effectiveness of the EF, *B. bassiana* (BbGc; BbPs) and *M. anisopliae* (MaPs) as mycoinsecticide dusts against *S. oryzae* adults and *C. cephalonica* larvae were evaluated. Each admixture of the isolates with either kaolin, talc or tapioca flour (20% w/w a.i.) as the carriers was thoroughly mixed with long grain rice in a plastic cup (8 cm diameter x 5 cm) at the rate of 0.05 g a.i., 0.1 g a.i. and 0.15 g a.i. in 50 g rice grains. All dosages of these EF isolates in all dust formulations gave 100 % mortality to *C. cephalonica* larvae 12 days after introduction (DAI). At the dosage of 0.05 g a.i. *M. anisopliae* (MaPs)

formulated in tapioca flour provided only 82.5% mortality to *S. oryzae* adult recorded within 15 days and this differed significantly from the dosage of 0.10 g a.i. or 0.15 g a.i. In general, the mycoinsecticides in kaolin and talc were more efficacious and faster knock out effect to both insects compared to that in tapioca flour or unformulated control. The 0.1 g a.i. of isolate BbGc in kaolin was significantly the best effecting 100% mortality seven DAI against *C. cephalonica* larvae, while it gave 98.75% mortality against *S. oryzae* adults.

Isolate BbGc in kaolin and talc administered at 0.4 g a.i. in 200 g rice packed in plastic kept at room temperature provided protection in excess of 90% mortality at 15 DAI against *C. cephalonica* larvae up to four months of storage. A lower mortality ranging between 65-78% was recorded on adults of *S. oryzae*. Formulations of MaPs seemed to be effective against both insects only within the first month of storage beyond which infectivity rapidly declined.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan ijazah Master Sains

**POTENSI ENTOMOPATOGEN HYPHOMYCETES TERPILIH UNTUK  
PENGAWALAN PEROSAK BERAS, *SITOPHILUS ORYZAE*  
(COLEOPTERA : CURCULIONIDAE) DAN *CORCYRA CEPHALONICA*  
(LEPIDOPTERA : PYRALIDAE)**

Oleh

Hendrawan Samodra

November 2003

**Pengerusi : Profesor Dr. Yusof Bin Ibrahim**

**Fakulti : Pertanian**

Kutu beras, *Sitophilus oryzae* (L.) dan rama-rama beras, *Corcyra cephalonica* Stn. adalah perosak-perosak serius biji-bijian dalam simpanan di dunia. Perosak- perosak ini menyerang berbagai produk simpanan dan merupakan perosak penting pada simpanan beras.

Penggunaan kulat entomopatogen (EF) adalah sesuatu pendekatan baru untuk pengawalan serangga perosak pada biji-bijian dalam simpanan. Potensi konidia kering tiga genera EF, *Beauveria bassiana*, *Metarhizium anisopliae* dan *Paecilomyces fumosoroseus* telah dikaji di dalam makmal terhadap kumbang *S. oryzae* dan larva *C. cephalonica*.

Patogenisiti sembilan pencilan EF terpilih (BbGc, BbPs, BbPc, MaOrMaj,





MaSc, MaGmC, MaOrMan, MaPs, PfPp) terhadap kumbang *S. oryzae* dan larva *C. cephalonica* telah dinilai di dalam makmal. Kesemua pencilan telah didapati patogenik terhadap kumbang *S. oryzae* dan larva *C. cephalonica* tetapi patogenisiti di antara pencilan didapati berbeza. Dua pencilan *B. bassiana* (BbGc and BbPs) dan satu pencilan *M. anisopliae* (MaPs) didapati superior dan mengakibatkan kematian yang tinggi terhadap kedua-dua serangga tersebut berbanding pencilan yang lain. Median kepekatan berkesan ( $EC_{50}$ ) untuk pencilan BbGc, BbPs and MaPs terhadap larva *C. cephalonica* masing-masing adalah  $1.238 \times 10^6$ ,  $2.072 \times 10^6$ ,  $1.775 \times 10^6$  konidia  $g^{-1}$ . Bagaimanapun, nilai  $EC_{50}$  yang lebih tinggi bagi pencilan-pencilan tersebut telah direkodkan terhadap kumbang *S. oryzae*, masing-masing adalah  $9.491 \times 10^6$ ,  $1.377 \times 10^7$ ,  $1.120 \times 10^7$  konidia  $g^{-1}$ . Oleh itu, boleh dirumuskan bahawa larva *C. cephalonica* adalah lebih rentan dari pada kumbang *S. oryzae* kepada ketiga-tiga pencilan itu. Median masa maut ( $LT_{50}$ ) pada kepekatan  $1 \times 10^9$  bagi pencilan-pencilan tersebut terhadap larva *C. cephalonica* ialah 3.5 d bagi BbGc, 3.6 d bagi BbPs dan 3.5 d bagi MaPs dan terhadap kumbang *S. oryzae* ialah 3.1 d bagi BbGc, 3.3 d bagi BbPs dan 2.5 d bagi MaPs.

Keberkesanan EF, *B. bassiana* (BbGc; BbPs) dan *M. anisopliae* (MaPs) sebagai mikoinsektisid debu terhadap kumbang *S. oryzae* dan larva *C. cephalonica* telah dinilai. Pencampuran setiap pencilan EF dengan kaolin, talkum atau tepung ubi kayu (20 % b/b b.a.) sebagai bahan pembawa telah digaul serata dengan beras panjang di dalam bekas plastik (8 cm diameter



x 5 cm) pada kadar 0.05 g b.a., 0.1 g b.a. dan 0.15 g b.a. dalam 50 g biji beras. Kesemua dos pencilan EF ini di dalam semua formulasi debu menghasilkan 100% kematian terhadap larva *C. cephalonica* 12 hari selepas diperlakukan (DAI). Pada dos 0.05 g b.a. *M. anisopliae* (isolate : MaPs) yang diformulasikan dalam tepung ubi kayu hanya menghasilkan 82.5% kematian terhadap kumbang *S. oryzae* setelah direkodkan selama 15 hari dan ini nyata berbeza dari pada dos 0.10 b.a. atau 0.15 b.a. Pada amnya, efikasi mikoinsektisid dalam kaolin dan talkum adalah lebih cepat memberi kesan rebah ke atas kedua serangga berkenaan berbanding didalam formulasi tepung ubi kayu atau kawalan tidak terformulasi. Pada dos 0.1 g b.a. pencilan BbGc dalam kaolin nyata paling baik memberikan 100% kematian terhadap larva *C. cephalonica* tujuh DAI, sedangkan ia hanya menghasilkan 98.75% kematian terhadap kumbang *S. oryzae*.

Pencilan BbGc dalam kaolin dan talkum yang dimasukkan pada dos 0.4 g b.a. ke dalam 200 g beras yang dimuatkan di dalam plastik pada suhu bilik memberikan perlindungan melebihi 90% kematian pada 15 DAI terhadap larva *C. cephalonica* sehingga empat bulan simpanan. Kematian lebih rendah antara 65-78% direkodkan terhadap kumbang *S. oryzae*. Formulasi debu MaPs agak berkesan terhadap kedua serangga tersebut hanya dalam bulan pertama penyimpanan selepas mana kejangkitannya menurun dengan cepat.

## ACKNOWLEDGEMENTS

Above all, I would like to thank Allah S.W.T., Most Gracious, Most Merciful, for his Compassion and Mercy.

I would like to express my deep sense of gratitude and appreciation to my supervisor, Professor Dr. Yusof Ibrahim, for his advice, guidance and constructive criticisms in connection with the research and preparation and revisions of this manuscript. I also would like to extend my thank to my other committee members: Professor Dr. Dzolkhifli Omar and Puan Noorma Osman.

I gratefully acknowledge Integrated Pest Management of Smallholder Estate Crops Project – Plant Quarantine Component in Agricultural Quarantine Agency, Ministry of Agriculture in Indonesia for providing me the financial support which enables me to pursue this degree.

A special note of thanks goes to my friends, colleagues, and staff members of Department of Plant Protection, Faculty of Agriculture, UPM.

I would like to express my deepest thanks and appreciation to my father, Hendro Suyoko who passed away five years ago. Also my mother, Wiryani for encouragement, support and endless prayers during my study in Malaysia. This endeavour would not have been feasible without the sacrifice, patience, understanding and encouragement of my dearest wife Takako.



## TABLE OF CONTENTS

DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xv
LIST OF FIGURES	xvii
LIST OF APPENDICES	xx
CHAPTER	
I INTRODUCTION	1
II. LITERATURE REVIEW	6
2.1. Rice weevil and rice moth	6
2.2. Economic impact	11
2.3. Management of rice weevil and rice moth	12
2.3.1. Cultural control and sanitary methods	12
2.3.2. Mechanical control	13
2.3.3. Physical control	13
2.3.4. Chemical control	16
2.3.5. Biological control	18
2.4. Entomopathogenic fungi	20
2.4.1. <i>Metarhizium anisopliae</i>	23
2.4.2. <i>Beauveria bassiana</i>	24
2.4.3. <i>Paecilomyces fumosoroseus</i>	27
2.5. Isolation of entomopathogenic fungi	28
2.6. Mass production	29
2.5.1. Production in vivo	29
2.5.2. Submerged culture	30
2.5.3. Surface culture	31
2.7. Formulation of biological insecticide	33
2.8. Inert dust	34



III	METHODOLOGY	38
	3.1. Location of research	38
	3.2. Cultures <i>Sitophilus oryzae</i> and <i>Corcyra cephalonica</i>	38
	3.3. Selection of entomopathogenic fungal sporulation	38
	3.4. Passage through of conidia to insect and reisolatation from cadaver	40
	3.5. Production of air-dried conidia	41
	3.6. Preparation insect for experiment	42
	3.6.1. <i>Sitophilus oryzae</i> adult	43
	3.6.2. <i>Corcyra cephalonica</i> egg	43
	3.6.3. <i>Corcyra cephalonica</i> larvae	44
	3.6.3.1. Selection for the larval instars to be used in pathogenicity test	44
	3.7. First experiment : selection for virulent isolates and pathogenicity test	44
	3.8. Second experiment : admixture conidial formulation with rice grain	45
	3.9. Third experiment : Viability of <i>Beauveria bassiana</i> and <i>Metarhizium anispliae</i> in stored grain	47
IV	RESULTS AND DISCUSSION	50
	4.1. Selection fungal of isolates	50
	4.2. Determination for virulent isolates	51
	4.3. Admixture conidial formlation with rice grain	61
	4.4. Viability of <i>B. bassiana</i> and <i>M. anisopliae</i>	71
V.	CONCLUSION	78
	REFERENCES	80
	APPENDICES	93
	BIODATA OF THE AUTHOR	110



## LIST OF TABLES

Table 1.	Some common entomogenous fungus and their host.	22
Table 2.	Type of diluents and carriers used in agricultural formulations.	36
Table 3.	Isolates of <i>B. bassiana</i> , <i>M. anisopliae</i> , and <i>P. fumosoroseus</i> and their original hosts and countries of origin.	40
Table 4.	Sporulation of selected entomopathogenic fungal isolates	50
Table 5.	Effective concentration of nine isolates of entomopathogenic fungal isolates against <i>S. oryzae</i> adult.	52
Table 6.	Effective concentration of nine isolates of entomopathogenic fungal isolates against <i>C. cephalonica</i> larvae.	53
Table 7.	Lethal time of <i>C. cephalonica</i> larvae against different conidial concentration of selected entomopathogenic fungal isolates.	54
Table 8.	Lethal time of <i>S. oryzae</i> adult against different conidial concentration of selected entomopathogenic fungal isolates.	57
Table 9.	Mean percentage mortality of <i>C. cephalonica</i> larvae upon exposure to different formulations of selected isolates of entomopathogenic fungi in 50 g rice grain seven days after treatment.	61
Table 10.	Mean percentage mortality of <i>S. oryzae</i> adult upon exposure to different formulations of selected isolates of entomopathogenic fungi in 50 g rice grain seven days after treatment.	64



Table 11. Mean percentage mortality of *S. oryzae* adult and mean percentage grain weight loss upon exposure to different formulations of isolates of entomopathogenic fungi in 50 g rice grain after four month of storage.

67

## LIST OF FIGURES

Figure 1.	Rice grain damage by <i>S. oryzae</i> (A), larvae (B,C), pupa (D) and adult (E).	7
Figure 2.	Egg (A), larvae (B), pupae (C), adults (D) of <i>C. cephalonica</i> ; rice damage (E).	8
Figure 3.	Good sporulation of <i>B. bassiana</i> (A) and poor sporulation (B) on PDA	39
Figure 4.	Fungal isolates produced from single spore isolation technique.	41
Figure 5.	Air dried conidia obtained through 125 $\mu$ m size sieve. <i>Metarhizium anisopliae</i> (A), <i>B. bassiana</i> (B) and <i>P. fumosoroseus</i> (C).	42
Figure 6.	<i>Corcyra cephalonica</i> place in oviposition jar	43
Figure 7.	Powdered formulation of <i>M. anisopliae</i> (A) and <i>B. bassiana</i> (B) on 50 g rice grain	47
Figure 8.	Powdered formulation of <i>M. anisopliae</i> on 200 g rice grain packed in plastic.	48
Figure 9.	Treated rice grains packed in plastic stored between one to six months	49
Figure 10.	Dried cadaver of <i>C. cephalonica</i> caused <i>B. bassiana</i> (A) and <i>M. anisopliae</i> (B)	58
Figure 11.	<i>Beauveria bassiana</i> growth from cadaver of <i>C. cephalonica</i> larvae after placed in moist filter paper.	58



Figure 12.	Sporulation of <i>M. anisopliae</i> from cadaver of <i>C. cephalonica</i> larvae	59
Figure 13.	Pupa of <i>C. cephalonica</i> infected by entomopathogenic fungus (A). <i>Beauveria bassiana</i> (B) and <i>M. anisopliae</i> (C) growth from pupae of <i>C. cephalonica</i>	59
Figure 14.	<i>Sitophilus oryzae</i> adult infected by <i>B. bassiana</i> . Mycellium growth from : antenna and rostrum (A), intersegmental membrane of abdomen (B), joint between thorax and abdomen (C) and sporulation cover all the insect body.	60
Figure 15.	<i>Sitophilus oryzae</i> adult infected by <i>M. anisopliae</i> . The fungus already cover the insect body	60
Figure 16.	Mean persentage mortality of <i>C. cephalonica</i> larvae against different dust formulations of selected entomopathogenic fungal isolates	63
Figure 17.	Mean persentage mortality of <i>S. oryzae</i> adult against different dust formulations of selected entomopathogenic fungal isolates.	66
Figure 18.	Rice grains introduced with <i>S. oryzae</i> adult and stored for four month. Untreated (A) showing badly damaged grains and treated with <i>B. bassiana</i> (B) showing comparatively clean grains	69
Figure 19.	Cadaver <i>C. cephalonica</i> larvae on rice grain infected <i>B. bassiana</i> (A) and <i>M. anisopliae</i> (B). All larvae become on rice grain not treated with fungus (C).	69
Figure 20.	Condition of rice grain introduced with <i>C. cephalonica</i> larvae. Not treated with fungus (A) and treated with <i>B. bassiana</i> (B) stored for four months.	70

- Figure 21. Viability of selected isolates of entomopathogenic fungal formulations in 200 g rice grain in plastic bag stored up to six months as indicated by mean percent mortality of *C. cephalonica* larvae 15 days after introduction. 72
- Figure 22. Viability of selected isolates of entomopathogenic fungi in 200 g rice grain in plastic bag stored up to six months as indicated by mean percent mortality of *S. oryzae* adult 15 days after introduction. 74
- Figure 23 Dry conidia of *M. anisopliae* will float during washing. (A) before washing, (B) washing and (C) after washing 76

## LIST OF APPENDICES

Appendix 1.	Lethal time of 5 different instar larva of <i>C. cephalonica</i> against <i>B. bassiana</i> (Isolate : BbGc) at concentration $1 \times 10^9$	93
Appendix 2.	Mean percent mortality of <i>S. oryzae</i> adult against nine entomopathogenic fungal isolates at 15 days after treatment	94
Appendix 3.	Mean percent mortality of <i>C. cephalonica</i> larvae against nine entomopathogenic fungal isolates at 15 days after treatment.	94
Appendix 4.	Mean percentage egg of <i>C. cephalonica</i> infected against nine entomopathogenic fungal isolates at 7 days after treatment.	95
Appendix 5.	The trend of mean percentage mortality of <i>S. oryzae</i> adult against nine entomopathogenic fungal isolates at concentration $1 \times 10^9$	96
Appendix 6.	The trend of mean percentage mortality of <i>C. cephalonica</i> larvae against nine entomopathogenic fungal isolates at concentration $1 \times 10^9$	96
Appendix 7.	Mean percentage mortality of <i>S. oryzae</i> adult upon exposure to <i>B. bassiana</i> (Isolate : BbGc) in 50 g rice grain.	97
Appendix 8.	Mean percentage mortality of <i>S. oryzae</i> adult upon exposure to <i>B. bassiana</i> (Isolate : BbPs) in 50 g rice grain.	98
Appendix 9.	Mean percentage mortality of <i>S. oryzae</i> adult upon exposure to <i>M. anisopliae</i> (Isolate : MaPs) in 50 g rice grain.	99



Appendix 10.	Mean percentage mortality of <i>C. cephalonica</i> larvae upon exposure to <i>B. bassiana</i> (Isolate : Bb GC) in 50 g rice grain.	100
Appendix 11.	Mean percentage mortality of <i>C. cephalonica</i> larvae upon exposure to <i>B. bassiana</i> (Isolate : Bb Ps) in 50 g rice grain.	101
Appendix 12.	Mean percentage mortality of <i>C. cephalonica</i> larvae upon exposure to <i>M. anisopliae</i> (Isolate : Ma Ps) in 50 g rice grain..	102
Appendix 13.	Mean percentage mortality of <i>S. oryzae</i> adult, total population and mean percentage grain weight loss upon exposure to three isolates of entomopathogenic fungal formulations in 50 g rice grain after four month of storage.	103
Appendix 14.	Mean percentage grain weight loss of treated and untreated fungus against <i>C. cephalonica</i> larvae in 50 g rice grain after four month of storage.	105
Appendix 15.	Mean percentage mortality of <i>S. oryzae</i> adult and <i>C. cephalonica</i> larvae against three dust formulations and unformulated 0.1 g a.i. of <i>B. bassiana</i> (BbGc) in 50 g rice grain.	107
Appendix 16.	Mean percent mortality of <i>S. oryzae</i> adult against three selected fungal formulations in 200 g rice grain stored up to six month at room temperature	108
Appendix 17.	Mean percent mortality of <i>C. cephalonica</i> larvae against three selected fungal formulations in 200 g rice grain stored up to six month at room temperature.	109

## CHAPTER I

### INTRODUCTION

Stored product pests are still one of the causes of qualitative and quantitative losses of food crops during storage. Quantitative damage is due to grain weight loss caused by insect feeding. Qualitative damage is due to product alterations such as loss of nutritional and aesthetic value, increased levels of rejects in the grain mass and loss of industrial (baking) characteristics. They are also important pests of stored seeds by damaging the seed embryos and causing a decrease in germination (Pranata *et al.*, 1988; Baier and Webster, 1992; Thuy *et al.*, 1994; Moino *et al.*, 1998).

The rice weevil, *Sitophilus oryzae* (L), is one of the most serious stored grain pests worldwide. This pest of whole grain originated in India and has spread worldwide by commerce. It is now a cosmopolitan pest. *Sitophilus oryzae*, an ubiquitous pest of economic importance, is an internal feeding insect that bores into stored grain. Adult weevils feed mainly on the endosperm, reducing the carbohydrate content while the larvae feed preferentially on the germ of the grain, thus removing a large percentage of the protein and vitamins (Belloa *et al.*, 2000).

The rice moth, *Corcyra cephalonica* (Stainton) is believed to be of eastern origin but it is now a cosmopolitan pest. It has spread throughout the world with the transport of food stuffs. Beside rice, the rice moth is also a



major pest of stored grains of pearl millet and sorghum. It becomes established more readily in stored seeds that have been damaged. For this reason the species is regarded as a secondary pest (Hodges, 1979). The broken seeds have always been found to provide a more suitable medium than either whole seeds or flours. However, the favourability of whole seeds in comparison with flours depends upon the commodity under consideration. Its development in rice is both more rapid and more successful with whole grain than with the flour.

The control of arthropod pests on stored products has been primarily through the use of fumigants and residual insecticides to augment the more obvious approach of hygiene (Brooker *et al.*, 1992; Adane *et al.*, 1996). The excessive use of conventional insecticides has resulted in a number of serious problems, such as resistant to the chemical insecticides, elimination of economically beneficial insects, persistence in the environment, toxicity to humans and wildlife and higher cost of crop production (Khan and Selman, 1989).

Synthetic chemical pesticides have been the main stay of insect pest control for the past 50 years. The advent of insecticide resistance, pest resurgence and concern over the environmental impact of agricultural inputs are increasingly focusing attention on biologically based form of pest control (Esser and Lemket, 1997). Resistance to chlorpyrifos-methyl has been documented for a number of strains of the lesser grain borer, *R. dominica* in