



UNIVERSITI PUTRA MALAYSIA

**DISTRIBUTION AND CONCENTRATION OF HEAVY METALS AND
POLYAROMATIC HYDROCARBONS IN SURFACE SEDIMENTS OF
KUALA SEPETANG, PERAK, MALAYSIA**

ROSLAN BIN ISMAIL

FP 2007 34

**DISTRIBUTION AND CONCENTRATION OF HEAVY METALS AND
POLYAROMATIC HYDROCARBONS IN SURFACE SEDIMENTS OF
KUALA SEPETANG, PERAK, MALAYSIA.**

By

ROSLAN BIN ISMAIL

**submitted to the Faculty of Agriculture, University Putra
Malaysia in fulfillment of the requirement of SPS 5999 (Project) for the award
of the master of Master of Science in Environmental Soil Science**



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

**DISTRIBUTION AND CONCENTRATION OF HEAVY METALS AND
POLYAROMATIC HYDROCARBONS IN SURFACE SEDIMENTS OF
KUALA SEPETANG, PERAK, MALAYSIA.**

By

ROSLAN ISMAIL

APRIL 2007

Chairman : Samsuri Abd. Wahid, PhD

Faculty : Agriculture

A study was conducted to determine the concentration and levels of six heavy metals (Ni, Zn, Pb, Cu, Cr and Cd: total 65 samples) and polyaromatic hydrocarbons (PAHs) (16 USEPA listed PAHs: total 45 samples) in the sediment surface layer (0-10 cm) of Kuala Sepetang estuary. Sediment samples were freeze dried (-30⁰C for 12-16 hour) to preserve the sample prior to analysis. From the analysis, the sediment pH (range: pH 3.48-7.48), the total carbon (organic and inorganic sources) ranges 1.8-8.3%, higher than the organic carbon (sedimentation and decomposition of leaf litter) content that ranges from 1.3% to 4.2%. The highest CEC was found at Sg. Reba (51.04 cmol_c/kg) while other rivers averaged at 20 cmol_c/kg. X-ray diffractometry analysis showed dominant presence of kaolinite (7.26Å, 3.57 Å) and quartz (4.26 Å, 3.33 Å) in the sediment. On the other hand, sequential extraction showed that Ni, Zn, Pb, Cu, Cr and Cd were primarily present in the residual fraction (non-mobile); especially Ni, Pb and Cd up to 80%. Meanwhile, the heavy



metals concentration at S1, S2, S3, S4, S5 and S6 in Kuala Sepetang showed Ni (5.2-36 mg/kg), Zn (52.4-312.80 mg/kg), Pb (1.8-159.60 mg/kg), Cu (7.6-31.20 mg/kg), Cr (3.2-77.2 mg/kg) and Cd (0.32-5.04 mg/kg) in dry weight basis are comparable with other Malaysian studies. The enrichment factor (EF) value showed very high enrichment of Pb and Cd compared to the other heavy metals. Despite that, Ni, Zn, Cu and Cr concentrations showed deficiency to minimal enrichment, indicating lithogenous material origin rather than anthropogenic input. Meanwhile, total PAHs range from 72ng/g to 2606 ng/g dry weight of sediment. PAHs concentration decreases in the order of: Sg. Reba (S1) (8548 ng/g) > Sg. Selinsing (S4) (5462 ng/g) > Sg. Sepetang (S3) (4249 ng/g) > Sg. Sangga Besar (S5)(3216 ng/g) > Sg. Sepetang (S2) (2961 ng/g) > Sg. Sangga Kecil (S6) (1184 ng/g). The concentrations of PAHs reported in Kuala Sepetang were comparable to those reported by Malaysia (Zakaria et al., 2002) and other regional studies. Generally, Kuala Sepetang showed much lower concentration of total PAHs with mixed pattern of strong pyrolytic and low petrogenic sources compared to Pearl River Estuary, China (408-10811 ng/g) with high petrogenic origin contamination level. Sediment in Sg. Reba (close to town area) has the most significant correlation (at $p < 0.01$) with physico-chemical properties. This study can be used by environmental agencies and private sector as a baseline data for mangrove sediments.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**TABURAN DAN KANDUNGAN LOGAM BERAT DAN POLIAROMATIK
HIDROKARBON DI PERMUKAAN SAMPEL ENAPAN DI KUALA
SEPETANG, PERAK, MALAYSIA.**

Oleh

ROSLAN ISMAIL

APRIL 2007

Pengerusi : Samsuri Abd. Wahid, PhD

Fakulti : Pertanian

Kajian untuk menentukan taburan dan kandungan logam berat (Ni, Zn, Pb, Cu, Cr dan Cd) sebanyak 65 sampel dan poliaromatik hidrokarbon (PAH) sebanyak 45 sampel pada permukaan (0-10 cm) dalam enapan telah dijalankan disekitar kawasan paya laut, Kuala Sepetang. Sampel enapan disejuk bekukan (-30°C selama 12-16 jam) untuk memelihara sample sebelum analisa. Dari analisa pH, julat pH enapan sekitar 3.48 hingga 7.48. Julat bagi karbon organik berbanding jumlah karbon keseluruhan adalah sekitar 1.8% hingga 8.3%. Ini mungkin disebabkan oleh pereputan tumbuhan dan fauna yang termendak di dalam enapan. Kadar tukarganti kation didapati paling tinggi di kawasan Sg. Reba ($51.04 \text{ cmol}_c/\text{kg}$), manakala sungai yang lain mempunyai julat sekitar $20 \text{ cmol}_c/\text{kg}$. Data analisa difraktometri X-ray menunjukkan kehadiran mineral kaolinit (7.26\AA , 3.57\AA) dan kuarza (4.26\AA , 3.33\AA) adalah paling banyak dalam enapan. Manakala, pengekstrakan berperingkat menunjukkan kehadiran logam berat dalam bahagian sisa-baki adalah paling tinggi, terutamanya unsur Ni, Pb dan Cd. Julat logam berat dalam sample enapan di

kawasan Kuala Sepetang bagi Ni (5.2-36 mg/kg), Zn (52.4-312.80 mg/kg), Pb (1.8-159.60 mg/kg), Cu (7.6-31.20 mg/kg), Cr (3.2-77.2 mg/kg) dan Cd (0.32-5.04 mg/kg) berjaya dibandingkan dengan kajian berkaitan yang terdahulu di Malaysia. Kadar kandungan logam Pb dan Cd didapati paling tinggi berbanding logam berat lain, dikawasan kajian. Jumlah kandungan keseluruhan PAH dalam enapan adalah dari 72 ng/g hingga 2606 ng/g berdasarkan berat kering. Kandungan PAH di kawasan kajian adalah seperti berikut: Sg. Reba (S1) (8548 ng/g) > Sg. Selinsing (S4) (5462 ng/g) > Sg. Sepetang (S3) (4249 ng/g) > Sg. Sangga Besar (S5)(3216 ng/g) > Sg. Sepetang (S2) (2961 ng/g) > Sg. Sangga Kecil (S6) (1184 ng/g). Kandungan PAH setara dibandingkan dengan kajian terdahulu dari Malaysia. Didapati , kandungan PAH di kawasan paya laut di sekitar Kuala Sepetang dipengaruhi oleh gabungan pencemaran dari sumber pirolitik dan petrogenik. Secara umum, dilihat kandungan PAH dalam enapan kawasan paya laut lebih rendah berbanding enapan kawasan marin yang tercemar seperti Pearl River Estuary, China (408-10811 ng/g). Ini sejajar dengan hasil kajian yang mendapati kandungan PAH dalam enapan dikawasan kajian masih rendah dengan kesan sumber pirolitik yang tinggi dan sumber petrogenik yang rendah. Enapan di Sg. Reba mempunyai korelasi yang amat signifikan dengan sifat kimia tanah pada $p < 0.01$. Hasil dan data dari kajian ini, sesuai digunakan oleh pelbagai agensi kerajaan dan swasta yang terlibat dengan aktiviti paya bakau sebagai sumber rujukan untuk kajian-kajian di masa hadapan.

ACKNOWLEDGEMENTS

Assalamualaikum; As given the opportunity, I would like to record my deepest appreciation to my dearest parents, Mrs.Hajijah Sulaiman and Mr. Ismail Abdullah and family members, for always believing in me and given me the moral and physical support throughout my studies.

Special dedication with sincere gratitude to Ms. Norzilawati Razali, for always being there when I need the most, thank you so much.

Warm and heart full regards to my supervisors, Dr. Samsuri Abd.Wahid, for the guidance and understanding throughout the years, also not forgotten; Assoc.Prof. Dr. Siti Zauyah Darus, Assoc. Prof. Dr. Che Fauziah Ishak and Dr. Mustafa Kamal Abd. Satar; for giving me a great perspective in the conducted study. Thank you, also to the lab assistant and staffs at Department of Land Management (Universiti Putra Malaysia), Kolej Universiti Sains dan Teknologi (KUSTEM) and Universiti Teknologi Malaysia (UTM) for giving a helping hand in making the study possible. Last but not least, I hope to contribute more and more in future to the country development. Wassalam.



I certify that an Examination Committee has met on 30th April 2007 to conduct the final examination of Roslan bin Ismail on his Master of Science thesis entitled “Distribution and Concentration of Heavy Metals and Polyaromatic Hydrocarbons in Surface Sediments of Kuala Sepetang, Perak, Malaysia” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the student be awarded the degree of Master of Science.


Members of the Examination Committee were as follows:

Hamdan Jol, PhD
Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Puziah Abd. Latif, PhD
Lecturer
Faculty of Environmental Studies
Universiti Putra Malaysia
(Internal Examiner)

Mohd. Hanafi Musa, PhD
Professor
Institute of Tropical Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Kamaruzzaman Yunus, PhD
Associate Professor
Kulliyah of Science
Universiti Islam Antarabangsa
(External Examiner)



HASANA H. GHAZALI, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 24 October 2007

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment on the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

Samsuri Abd.Wahid, PhD

Lecturer
Faculty of Agriculture
University Putra Malaysia
(Chairman)

Siti Zauyah Hj.Darus, PhD

Associate Professor
Faculty of Agriculture
University Putra Malaysia
(Member)

Che Fauziah Ishak, PhD

Associate Professor
Faculty of Agriculture
University Putra Malaysia
(Member)

Mustafa Kamal Abd.Satar, PhD

Lecturer
Faculty of Agriculture
University Putra Malaysia
(Member)



AINI IDERIS, PhD
Professor and Dean
School of Graduate
Studies
Universiti Putra
Malaysia

Date: 24 October 2007



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

ROSLAN BIN ISMAIL

Date: 22 October 2007



LIST OF TABLES

| Table | | Page |
|-------|--|------|
| 1.0 | Economic significance of coastal resources and in Malaysia | 1.8 |
| 2.1a | Five contamination categories are recognized on the basis of the enrichment factor | 2.18 |
| 2.2 | Molecular Structure and Carcinogenic Evaluation of the PAHs of Primary Environmental Concern | 2.24 |
| 3.0a | Location of the sampling site for heavy metal sediment samples in Matang mangrove forest | 3.3 |
| 3.0b | Location of the sampling site for polyaromatic Hydrocarbons sediment samples in Matang mangrove forest | 3.7 |
| 3.0c | Location of the sampling site for background sediment samples | 3.10 |
| 3.1 | Validation of Accuphase® certified standard containing 16 USEPA PAHs using GC-FID instrument | 3.17 |
| 3.2 | Recovery percentage and detection limit of 16 PAHs by GC-FID | 3.18 |
| 3.3 | GC-FID temperature and pressure program for PAHs analysis | 3.21 |
| 4.1 | Sediment pH value in Kuala Sepetang estuary and Pasir Panjang background area | 4.3 |
| 4.2 | Total Carbon value of sediment in Kuala Sepetang estuary and Pasir Panjang background area | 4.5 |
| 4.3 | Organic Carbon value of sediment in Kuala Sepetang estuary and Pasir Panjang background area | 4.5 |
| 4.4 | Cation Exchange Capacity value of sediment in Kuala Sepetang estuary and Pasir Panjang background area | 4.7 |
| 4.5 | Electrical conductivity value of sediment in Kuala Sepetang estuary and Pasir Panjang background area | 4.8 |



| | | |
|-------|---|------|
| 4.6 | Value of d-spacing used to identify the mineral in different fraction | 4.15 |
| 4.7 | Relative intensity of sand fraction in the sediment sample from Kuala Sepetang mangrove | 4.17 |
| 4.8 | Relative intensity of silt fraction in the sediment sample from Kuala Sepetang mangrove | 4.18 |
| 4.9 | Relative intensity of clay fraction in the sediment sample from Kuala Sepetang mangrove | 4.20 |
| 4.10 | Comparison of heavy metal content in Kuala Sepetang estuary and other mangrove sediment studies | 4.23 |
| 4.11 | Total heavy metal concentration from the six rivers in Kuala Sepetang study area and the Pasir Panjang (background value) | 4.25 |
| 4.12 | Classification of sediments by heavy metal concentration (mg/kg) in fine grained-sized fraction (<63um) according to the background values and classification system from the Hong Kong Environmental Protection Department (EPD, 1992) | 4.26 |
| 4.12a | Enrichment factor (EF) value from sediment sample in Kuala Sepetang area | 4.48 |
| 4.13 | Correlation of total heavy metal concentration and chemical properties of sediment in Sg.Reba | 4.52 |
| 4.14 | Correlation of total heavy metal concentration and chemical properties of sediment in Sg.Sepetang (northern) | 4.52 |
| 4.15 | Correlation of total heavy metal concentration and chemical properties of sediment in Sg.Sepetang (southern) | 4.52 |
| 4.16 | Correlation of total heavy metal concentration and chemical properties of sediment in Sg.Selinsing | 4.53 |
| 4.17 | Correlation of total heavy metal concentration and chemical properties of sediment in Sg.Sangga Besar | 4.53 |
| 4.18 | Correlation of total heavy metal concentration and chemical properties of sediment in Sg.Sangga Kecil | 4.53 |



| | | |
|------|--|-------|
| 4.19 | Correlation of total heavy metal concentration and chemical properties of sediment in Pasir Panjang | 4.54 |
| 4.20 | Mean and standard deviation of available heavy metal concentration (mg/kg) in Kuala Sepetang estuary using EDTA method | 4.59 |
| 4.21 | Correlation between EDTA and soil properties of sediment in Kuala Sepetang area | 4.60 |
| 4.22 | Mean and standard deviation of available heavy metal concentration (mg/kg) in Kuala Sepetang estuary using DTPA method | 4.63 |
| 4.23 | Correlation between DTPA and soil properties of sediment in Kuala Sepetang area | 4.64 |
| 4.24 | Mean and standard deviation of available heavy metal concentration (mg/kg) in Kuala Sepetang estuary using HCl method | 4.67 |
| 4.25 | Correlation between HCl and soil properties of sediment in Kuala Sepetang area | 4.68 |
| 4.32 | Concentrations of total polyaromatic hydrocarbons (PAHs; ng/g) in other sediment studies | 4.89 |
| 4.33 | Concentrations of total polyaromatic hydrocarbons (PAHs; ng/g dry weight) and ratios of different individual PAH compound in Sg. Reba | 4.93 |
| 4.34 | Concentrations of total polyaromatic hydrocarbons (PAHs; ng/g dry weight) and ratios of different individual PAH compound in Sg. Sepetang (S2) | 4.99 |
| 4.35 | Concentrations of total polyaromatic hydrocarbons (PAHs; ng/g dry weight) and ratios of different individual PAH compound in Sg. Sepetang (S3) | 4.103 |
| 4.36 | Concentrations of total polyaromatic hydrocarbons (PAHs; ng/g dry weight) and ratios of different individual PAH compound in Sg. Selinsing (S4) | 4.107 |
| 4.37 | Concentrations of total polyaromatic hydrocarbons (PAHs; ng/g dry weight) and ratios of different individual PAH compound in Sg. Sangga Besar (S5) | 4.111 |



| | | |
|------|---|-------|
| 4.38 | Concentrations of total polycyclic aromatic hydrocarbons (PAHs; ng/g dry weight) and ratios of different individual PAH compound in Sg. Sangga Kecil (S6) | 4.115 |
| 4.39 | Median and range of concentrations (ng/g dry weight) of individual PAHs compounds in mangrove sediments in Matang mangrove forest and toxicity guidelines | 4.120 |
| 5.1 | The risk assessment code (RAC) and criteria for sediment | 5.5 |
| 5.2 | Risk Assessment Code criteria for sediment in sampling area | 5.6 |



LIST OF FIGURES

| Figure | | Page |
|--------|--|------|
| 1 | Peninsular Malaysia shows Perak state | 1.2 |
| 1.1a | Matang mangrove forest located in Larut Matang and Selama | 1.3 |
| 1.1b | Forest management zones in Matang Mangroves | 1.4 |
| 1.2 | Map sheet index and felling series | 1.6 |
| 2.1 | Relationship between Ph/Ant and Fluo/Pyr ratio | 2.33 |
| 3.1 | Sampling site for heavy metal (65 sample) in the Kuala Sepetang estuary | 3.6 |
| 3.2 | Sampling site for PAHs (45 sample) in the Kuala Sepetang estuary | 3.9 |
| 4.5a | Percentage of sand, silt and clay in Matang mangrove | 4.12 |
| 4.5b | Texture of the sediment in Matang mangrove | 4.13 |
| 4.6 | Mean total heavy metal in Sg.Reba sediment | 4.27 |
| 4.7 | Mean total heavy metal in Sg.Sepetang (S2) sediment | 4.30 |
| 4.8 | Mean total heavy metal in Sg.Sepetang (S3) sediment | 4.30 |
| 4.9 | Mean total heavy metal in Sg.Selinsing sediment | 4.31 |
| 4.10 | Mean total heavy metal in Sg.Sangga Besar sediment | 4.34 |
| 4.11 | Mean total heavy metal in Sg.Sangga Kecil sediment | 4.34 |
| 4.12 | Distribution of nickel (mg/kg) in the sediment of Matang mangrove forest | 4.36 |
| 4.13 | Distribution of zinc (mg/kg) in the sediment of Matang mangrove forest | 4.38 |
| 4.14 | Distribution of lead (mg/kg) in the sediment of Matang mangrove forest | 4.40 |
| 4.15 | Distribution of copper (mg/kg) in the sediment of Matang mangrove forest | 4.42 |



| | | |
|------|--|-------|
| 4.16 | Distribution of cadmium (mg/kg) in the sediment of Matang mangrove forest | 4.44 |
| 4.17 | Distribution of chromium (mg/kg) in the sediment of Matang mangrove forest | 4.46 |
| 4.18 | Mean concentration value of available heavy metals (EDTA method) | 4.59 |
| 4.19 | Mean concentration value of available heavy metals (DTPA method) | 4.63 |
| 4.20 | Mean concentration value of available heavy metals (HCl method) | 4.67 |
| 4.21 | Fractionation of nickel (Ni) in sediment | 4.73 |
| 4.22 | Fractionation of zinc (Zn) in sediment | 4.75 |
| 4.23 | Fractionation of cadmium (Cd) in sediment | 4.77 |
| 4.24 | Fractionation of lead (Pb) in sediment | 4.79 |
| 4.25 | Fractionation of copper (Cu) in sediment | 4.82 |
| 4.26 | Fractionation of chromium (Cr) in sediment | 4.84 |
| 4.27 | Distribution of total PAHs (ng/g) in the Matang mangrove forest | 4.87 |
| 4.28 | Concentration of PAHs in Sg. Reba (S1) | 4.97 |
| 4.29 | Concentration of PAHs in Sg. Sepetang (S2) | 4.101 |
| 4.30 | Concentration of PAHs in Sg. Sepetang (S3) | 4.105 |
| 4.31 | Concentration of PAHs in Sg. Selinsing (S4) | 4.108 |
| 4.32 | Concentration of PAHs in Sg. Sangga Besar (S5) | 4.112 |
| 4.33 | Concentration of PAHs in Sg. Sangga Kecil (S6) | 4.117 |



LIST OF ABBREVIATIONS

| | |
|-------|---|
| PAHs | polyaromatic hydrocarbons |
| HM | heavy metals |
| HMW | High Molecular Weight |
| LMW | Low Molecular Weight |
| EDTA | Ethylenediaminetetraaceticacid |
| DTPA | Diaminetetraphenylaceticacid |
| mg/kg | concentration in part per million (ppm) |
| ng/g | concentration in part per billion (ppb) |
| S1 | Sg.Reba |
| S2 | Sg.Sepetang (northern part) |
| S3 | Sg.Sepetang (southern part) |
| S4 | Sg.Selinsing |
| S5 | Sg.Sangga Besar |
| S6 | Sg.Sangga Kecil |
| Naph | Naphthalene |
| Acy | Acenaphylene |
| Ace | Acenaphthene |
| Fl | Fluorene |
| Ant | Anthracene |
| Phe | Phenanthrene |
| Fluo | Fluoranthene |
| Pyr | Pyrene |
| BaA | Benzo(a)anthracene |



| | |
|-------|----------------------------------|
| Chry | Chrysene |
| BbF | Benzo(b)fluoranthene |
| BkF | Benzo(k)fluoranthene |
| BaP | Benzo(a)pyrene |
| BghiP | Benzo(g,h,i)perylene |
| IP | Indeno(1,2,3-cd)perylene |
| dBahA | Dibenz(a,h)anthracene |
| CEC | Cation exchange capacity |
| OM | Organic matter |
| EC | Electrical conductivity |
| ** | highly significant at $p < 0.05$ |
| * | significant at $p < 0.05$ |



TABLE OF CONTENTS

| | Page |
|--|-------------|
| ABSTRACT | iii |
| ABSTRAK | v |
| ACKNOWLEDGEMENTS | vii |
| APPROVAL | viii |
| DECLARATION | x |
| LIST OF TABLES | xi |
| LIST OF FIGURES | xv |
| LIST OF ABBREVIATIONS | xvii |
| | |
| CHAPTER | |
| | |
| 1 INTRODUCTION | |
| 1.0 Introduction to Mangrove Forest | 1.1 |
| 1.1 Sources of contaminants in mangrove area | 1.9 |
| 1.2 Objectives of the study | 1.10 |
| | |
| 2 LITERATURE REVIEW | |
| 2.0 Introduction to contaminants | 2.1 |
| 2.1 Contamination input in mangrove ecosystem | 2.2 |
| 2.1.1 Land based input | 2.2 |
| 2.1.2 Atmospheric precipitation | 2.3 |
| 2.1.3 Sea based input | 2.4 |
| 2.2 Classification of heavy metals | 2.5 |
| 2.3 Source of heavy metals | 2.7 |
| 2.4 Effects of heavy metals | 2.8 |
| 2.4.1 Effects of heavy metals on human health | 2.8 |
| 2.4.2 Effects of heavy metals on plants | 2.11 |
| 2.4.3 Effects of heavy metals on fauna | 2.11 |
| 2.5 Heavy metals in the sediment | 2.12 |
| 2.5.1 Soil pH | 2.12 |
| 2.5.2 Organic Matter | 2.13 |
| 2.5.3 Cation Exchange Capacity (CEC) | 2.14 |
| 2.5.4 Electrical conductivity | 2.14 |
| 2.5.5 Organic Carbon and Total Carbon | 2.15 |
| 2.5a Enrichment factors (EF) of sediment | 2.16 |
| 2.6 Mineralogy of heavy metals | 2.18 |
| 2.7 Analysis of heavy metals | 2.19 |
| 2.8 Speciation analysis of heavy metals | 2.19 |
| 2.8.1 Operationally and functionally defined species characterization | 2.21 |
| 2.8.2 Effect upon sample pretreatment upon speciation | 2.21 |
| 2.9 Polyaromatic hydrocarbons (PAHs) in the environment | 2.22 |



| | |
|---|------|
| 2.10 Sources of polyaromatic hydrocarbons in the environment | 2.26 |
| 2.10.1 Pyrogenic PAHs | 2.27 |
| 2.10.2 Petrogenic PAHs | 2.27 |
| 2.10.3 Diagenic PAHs | 2.28 |
| 2.10.4 Biogenic PAHs | 2.28 |
| 2.11 Effects of polyaromatic hydrocarbons | 2.28 |
| 2.12 Bioavailability of polyaromatic hydrocarbons | 2.30 |
| 2.13 Fate of polyaromatic hydrocarbons in the sediments | 2.31 |
| 2.14 Polyaromatic hydrocarbons ratio | 2.32 |
| 2.14.1 Phenanthrene/Anthracene (Phe/Ant) ratio | 2.32 |
| 2.14.2 Fluoranthene/Pyrene (Fluo/Pyr) ratio | 2.34 |
| 2.14.3 Low Molecular Weight/High Molecular Weight (LMW/HMW) ratio | 2.35 |
| 2.14.4 Other PAHs ratios | 2.35 |
| 2.15 Effects Range-Low (ER-L) and Effects Range Median (ER-M) | 2.36 |
| 2.16 Extraction of polyaromatic hydrocarbons | 2.36 |
| 2.17 Analysis of polyaromatic hydrocarbons | 2.36 |
| 2.18 Problems related to PAH analysis | 2.37 |
| 2.18.1 Sampling | 2.37 |
| 2.18.2 Analysis | 2.38 |
| 2.18.3 Interpretation | 2.38 |
| 2.19 Summary | 2.39 |

3 MATERIALS AND METHODS

| | |
|--|------|
| 3.0 Study Area | 3.1 |
| 3.1 Sampling on the sediment | 3.2 |
| 3.1.1 Sample preparation | 3.2 |
| 3.1.2 Sample pre-treatment | 3.11 |
| 3.2 Chemicals and apparatus | 3.11 |
| 3.2.1 Chemicals | 3.11 |
| 3.2.2 Apparatus | 3.12 |
| 3.2.3 Cleaning glassware | 3.12 |
| 3.3 Characterizations of sample | 3.13 |
| 3.3.1 Soil pH | 3.13 |
| 3.3.2 Total Carbon | 3.13 |
| 3.3.3 Organic carbon | 3.13 |
| 3.3.4 Cation exchange capacity | 3.14 |
| 3.3.5 Electrical conductivity of sediment solution | 3.14 |
| 3.3.6 Particle size analysis | 3.14 |
| 3.3.7 X-ray diffractometry (XRD) | 3.14 |
| 3.4 Heavy metals analysis | 3.15 |
| 3.4.1 Total heavy metals analysis (Aqua-Regia method) | 3.15 |
| 3.4.2 Available heavy metal analysis (EDTA, DTPA and HCl method) | 3.15 |
| 3.5 Sequential extraction | 3.16 |



| | |
|---|------|
| 3.6 Polyaromatic hydrocarbons analysis | 3.16 |
| 3.6.1 External standard stock solution preparation | 3.16 |
| 3.6.2 Determination of polyaromatic hydrocarbons detection limit | 3.18 |
| 3.6.3 Analyte recovery test | 3.19 |
| 3.6.4 Quality control | 3.20 |
| 3.7 Polyaromatic hydrocarbon extraction | 3.20 |
| 3.8 Analysis of PAHs in sediment | 3.21 |
| 3.8.1 Gas Chromatography Flame Ionization Detector (GC-FID) analysis | 3.21 |
| | |
| 4 RESULTS AND DISCUSSION | |
| 4.0 Introduction | 4.1 |
| 4.1 Characterization of sediment | 4.2 |
| 4.1.1 Sediment pH | 4.2 |
| 4.1.2 Total Carbon and Organic Carbon | 4.4 |
| 4.1.3 Cation Exchange Capacity | 4.7 |
| 4.1.4 Electrical conductivity (EC) | 4.8 |
| 4.1.5 Texture of sediments | 4.9 |
| 4.1.6 X-ray diffractometry | 4.14 |
| 4.2 Concentration of heavy metals in Kuala Sepetang estuary | 4.22 |
| 4.3 Distribution of heavy metals in the Kuala Sepetang estuary | 4.35 |
| 4.3.1 Nickel | 4.35 |
| 4.3.2 Zinc | 4.37 |
| 4.3.3 Lead | 4.39 |
| 4.3.4 Copper | 4.41 |
| 4.3.5 Cadmium | 4.43 |
| 4.3.6 Chromium | 4.45 |
| 4.3a Enrichment factor (<i>EF value</i>) in sediment | 4.47 |
| 4.4 Relationship between total heavy metal and chemical properties | 4.50 |
| 4.4.1 Nickel | 4.51 |
| 4.4.2 Zinc | 4.55 |
| 4.4.3 Lead | 4.55 |
| 4.4.4 Copper | 4.56 |
| 4.4.5 Chromium | 4.56 |
| 4.4.6 Cadmium | 4.57 |
| 4.5 Available heavy metal | 4.58 |
| 4.5.1 EDTA method | 4.58 |
| 4.5.2 DTPA method | 4.62 |
| 4.5.3 HCl method | 4.66 |
| 4.6 Fractionation study of heavy metals in sediment | 4.70 |
| 4.6.1 Nickel | 4.72 |
| 4.6.2 Zinc | 4.74 |
| 4.6.3 Cadmium | 4.77 |
| 4.6.4 Lead | 4.79 |
| 4.6.5 Copper | 4.81 |
| 4.6.6 Chromium | 4.83 |

| | | |
|----------|--|------------|
| 4.7 | Distribution of PAHs in sediment | 4.86 |
| 4.7.1 | Distribution of PAHs in Sg. Reba (S1) | 4.92 |
| 4.7.2 | Distribution of PAHs in Sg. Sepetang (S2) | 4.98 |
| 4.7.3 | Distribution of PAHs in Sg. Sepetang (S3) | 4.102 |
| 4.7.4 | Distribution of PAHs in Sg. Selinsing (S4) | 4.106 |
| 4.7.5 | Distribution of PAHs in Sg. Sangga Besar (S5) | 4.110 |
| 4.7.6 | Distribution of PAHs in Sg. Sangga Kecil (S6) | 4.114 |
| 4.8 | Relationship between PAHs and sediment texture | 4.118 |
| 4.9 | Toxicity of PAHs in sediment | 4.119 |
| | | |
| 5 | CONCLUSION | |
| 5.1 | Heavy metals in Kuala Sepetang | 5.1 |
| 5.1a | Enrichment factor (EF value) in Kuala Sepetang | 5.3 |
| 5.2 | Sequential extraction in Kuala Sepetang | 5.5 |
| 5.3 | Polyaromatic hydrocarbons (PAHs) in Kuala Sepetang | 5.7 |
| 5.3a | Summary | 5.11 |
| | REFERENCES | R.1 |
| | APPENDICES | A.1 |
| | BIODATA OF THE AUTHOR | B.1 |



CHAPTER ONE

INTRODUCTION

1.0 Introduction to Mangrove Forest

At present, Malaysia (Figure 1) still has 566,856 hectare (ha) of mangroves that 99,767 ha in the Peninsular Malaysia, and nearly 44,000 ha are situated in Perak under the Matang Mangrove Forest Reserve area (Figure 1.1a); 340,689 ha in Sabah, and the rest, 126,400 ha in Sarawak. From year 1980 to 1990, Malaysia has lost nearly 12% of its mangrove forest, with more than 1% rates of loss per year. In respond to rate of the loss, ecological restoration is being carried out throughout South East Asia, by simple management plan based on clear felling small patches and replanting with seedlings (Figure 1.2). Generally, mangrove soils are found along intertidal shores and estuarine such as Brazil, Australia, Malaysia, China and many other tropical countries. Mangroves forest help to protect the coastline from erosion, storm damage, and wave action by acting as buffers zone. It catches alluvial materials and deposits them in sediment, thus stabilizing land elevation. In recent years, many concerns have been raised regarding the threat of contamination on the mangrove forest ecosystem around the world. Human activities, such as industrialization and urbanization have led to various source of effluent being channeled into the mangrove ecosystem. The effluents may contain contaminants such as heavy metals and PAHs. The heavy metals and PAHs can precipitate and settle into sediments as studied elsewhere (Budzinski et al., 1997; Kucklik et al., 1997; Hartmann et al., 2000).



Figure 1: Peninsular Malaysia shows Perak state
Source: taken online from www.ipoh-online.com.my



Figure 1.1a: Matang mangrove forest located in Larut Matang and Selama

Note: The red circle in the map shows the study area

Source: taken online from www.ipoh-online.com.my

