

Efficacy of a bacteriophage isolated from chickens as a therapeutic agent for colibacillosis in broiler chickens

ABSTRACT

The efficacy of bacteriophage EC1, a lytic bacteriophage, against *Escherichia coli* O78:K80, which causes colibacillosis in poultry, was determined in the present study. A total of 480 one-day-old birds were randomly assigned to 4 treatment groups, each with 4 pens of 30 birds. Birds from the control groups (groups I and II) received PBS (pH 7.4) or 10(10) pfu of bacteriophage EC1, respectively. Group III consisted of birds challenged with 10(8) cfu of *E. coli* O78: K80 and treated with 10(10) pfu of bacteriophage EC1 at 2 h postinfection, whereas birds from group IV were challenged with 10(8) cfu of *E. coli* O78: K80 only. All the materials were introduced into the birds by intratracheal inoculation. Based on the results of the present study, the infection was found to be less severe in the treated *E. coli*-challenged group. Mean total viable cell counts of *E. coli* identified on eosin methylene blue agar (designated EMB + *E. coli*) in the lungs were significantly lower in treated, *E. coli*-challenged birds than in untreated, *E. coli*-challenged birds on d 1 and 2 postinfection. The EMB + *E. coli* isolation frequency was also lower in treated birds; no *E. coli* was detectable in blood samples on any sampling day, and *E. coli* were isolated only in the liver, heart, and spleen of treated chickens at a ratio of 2/6, 1/6, and 3/6, respectively, at d 1 postinfection. The BW of birds from the *E. coli*-challenged group treated with bacteriophage EC1 were not significantly different from those of birds from both control groups but were 15.4% higher than those of the untreated, *E. coli*-challenged group on d 21 postinfection. The total mortality rate of birds during the 3-wk experimental period decreased from 83.3% in the untreated, *E. coli*-challenged birds (group IV) to 13.3% in birds treated with bacteriophage EC1 (group III). These results suggest that bacteriophage EC1 is effective *in vivo* and could be used to treat colibacillosis in chickens.

Keyword: bacteriophage, *Escherichia coli*, colibacillosis, broiler