



UNIVERSITI PUTRA MALAYSIA

**EFFECTS OF *ANDROGRAPHIS PANICULATA*, *CARICA PAPAYA*
AND *CASSIA ALATA* EXTRACTS ON THE REPRODUCTIVE
SYSTEMS OF MICE**

AZMAHANI ABDULLAH

FPSK (M) 2003 7

EFFECTS OF *ANDROGRAPHIS PANICULATA*, *CARICA PAPAYA* AND *CASSIA ALATA* EXTRACTS ON THE REPRODUCTIVE SYSTEMS OF MICE

By

AZMAHANI ABDULLAH

**Thesis Submitted to the School of Graduates Studies, Universiti Putra Malaysia,
In Fulfillment of the Requirement for the Degree of Master of Science**

November 2003



DEDICATION

"Dedicated to my supervisor Associate Professor Dr. Mohd Nazrul Hakim Abdullah, my father (Abdullah), my mom (Azizah) and my brothers and sisters (Arzura, Azlinda, Al-Fatihah, Adibah, Al-Fadilah, Mohd. Asyraf, Alia Shazwani and Aiman Adli & Friends"



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

EFFECTS OF *ANDROGRAPHIS PANICULATA*, *CARICA PAPAYA* AND *CASSIA ALATA* EXTRACTS ON THE REPRODUCTIVE SYSTEMS OF MICE

By

AZMAHANI ABDULLAH

November 2003

Chairman : Associate Professor Mohammad Nazrul Hakim bin Abdullah, Ph. D.

Faculty : Medicine and Health Sciences

In facing the high rise of population especially in developing countries, a continuing need is felt for oral and non-surgical methods of contraception. Plants extracts have been used widely in traditional medical practice especially as an anti-fertility agent. The *Andrographis paniculata*, *Cassia alata* and *Carica papaya* extracts have been reported to poses multifunctional properties. In the present studies, the effects of 20 mg/kg body weight ethanol extract of *Andrographis paniculata*, *Carica papaya* and *Cassia alata* on the reproductive systems was studied after 30 days and 90 days of treatment in female and male mice. Clomiphene citrate, given daily at 50 mg/kg body weight, was used for the purposed comparison. Results obtained showed that the female genital organs (ovaries and uteri) weight was reduced significantly after treatment with *Carica papaya* and *Andrographis paniculata* respectively. All females groups showed changes in fertility and reduction in litter number. However, after treatment with the same extracts,



the litter number for *Cassia alata* treatment was found to increase as compared to control group. A significant decrease in the number of Graffian follicles and corpora lutea with significant increase in number of atretic follicles in all treatment groups was observed except in the group treated with *Cassia alata*. There were no changes in the weight of the body, epididymides and seminal vesicles. A reduction in epididymal sperm count, sperm motility, number of fertile males, litter number, and serum testosterone levels were also observed after manipulation of all extracts duration of treatment except *Cassia alata*. Testis histology study revealed spermatogenetic arrest at the spermatocyte level. In mice treated with clomiphene citrate, the body weight, testis, epididymis, and seminal vesicle weight, sperm counts, and sperm motility were found to reduce. Sperm morphology revealed abnormalities of the head, neck and tail. Progressive regress of the seminiferous tubules with most tubules devoid of spermatozoa and spermatid were also seen. However, these effects were reversible upon cessation of treatment. The findings therefore suggest that *Carica papaya* and *Andrographis paniculata* have mild inhibitory effects on male and female reproductive systems as compared to control group (clomiphene citrate). However, *Cassia alata* extracts aid in promoting fertility by improving male and female reproductive functions.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

KESAN EKSTRAK *ANDROGRAPHIS PANICULATA*, *CARICA PAPAYA* DAN *CASSIA ALATA* KE ATAS SISTEM PEMBIAKAN MENCIT

Oleh

AZMAHANI ABDULLAH

November 2003

Pengerusi : Profesor Madya Mohammad Nazrul Hakim Abdullah, Ph. D.

Fakulti : Perubatan dan Sains Kesihatan

Dengan meningkatnya populasi penduduk terutama di negara membangun, kaedah perancang keluarga tanpa pembedahan yang berterusan diperlukan. Ekstrak tumbuhan telah digunakan dengan meluas dalam perubatan tradisional terutama sebagai agen anti-kesuburan. Ekstrak *Andrographis paniculata*, *Carica papaya* dan *Cassia alata* telah dilaporkan mempunyai pelbagai kegunaannya. Dalam kajian ini, kesan 20 mg/kg berat badan ekstrak etanol *Andrographis paniculata*, *Carica papaya* dan *Cassia alata* dikaji pada sistem pembiakan mencit betina dan jantan selepas 30 dan 90 hari rawatan. Clomiphene sitrat, di beri setiap hari pada 50 mg/kg berat badan digunakan untuk tujuan perbandingan. Keputusan kajian mendapati berat alat kelamin betina (ovari dan uteri) telah menurun secara bererti selepas rawatan *Carica papaya* dan *Andrographis paniculata*. Kesemua kumpulan betina menunjukkan perubahan dalam kesuburan dan penurunan dalam bilangan anak. Walaubagaimanapun, bilangan anak bagi rawatan *Cassia alata* didapati bertambah berbanding kumpulan kawalan. Penurunan bererti pada

bilangan folikel Graffian dan corpora lutea dan peningkatan bererti pada bilangan folikel atretik pada semua rawatan melainkan ekstrak *Cassia alata*. Berat badan dan berat epididimis dan vesikel seminal tidak menunjukkan sebarang perubahan. Pengurangan dalam bilangan sperma epididimis, pergerakan sperma, bilangan jantan yang subur, bilangan anak dan aras testosterone serum juga dilihat selepas manipulasi jangka masa rawatan kesemua ekstrak kecuali *Cassia alata*. Kajian histologi pada testis menunjukkan terhentinya proses spermatogenesis pada peringkat spermatosit. Dalam mencit yang dirawat dengan clomiphene sitrate, berat badan, berat testis, epididimis, dan vesikel seminal, serta bilangan dan pergerakan sperma didapati berkurangan. Morfologi sperma menunjukkan kecacatan pada bahagian kepala, leher dan ekor. Dari kajian histologi testis pula menunjukkan tanda-tanda regrasi pada tubul seminiferous dengan ketidakhadiran spermatozoa dan spermatid pada kebanyakan tubul. Namun, kesan ini hilang selepas rawatan dihentikan. Dengan itu, penemuan ini mencadangkan bahawa ekstrak *Andrographis paniculata* dan *Carica papaya* memberi sedikit kesan perencatan pada sistem pembiakan betina dan jantan berbanding kumpulan kawalan (clomiphene citrate). Sebaliknya ekstrak *Cassia alata* membantu menambah kesuburan dengan meningkatkan fungsi sistem pembiakan mencit jantan dan betina.

ACKNOWLEDGEMENTS

In the Name of Allah, the Most Benevolent and the Most Merciful....

Alhamdulillah, first of all I would like to express my utmost thanks and gratitude to Almighty Allah S.W.T. who has given me the capability to complete this thesis and my salawat and salam to His righteous messenger, prophet Muhammad s.a.w.

Deepest gratitude and appreciation is expressed to my supervisor, Associate Professor Dr. Mohd. Nazrul Hakim Abdullah, for his dedication, guidance, support, endless encouragement and patience towards completing the research. I would like also to extend my thanks to other supervisory committee members, Dr. Sabrina Sukardi, Prof. Dato' Dr. Abdul Salam Abdullah and Dr. Wan Nordin Wan Mahmud for their guidance, advice and encouragement throughout the duration of this project.

Heartfelt thanks to Puan Sapiah Ismail, En. Kufli Che Nor and all staff members of the Histology Laboratory and Physiology Laboratory, Faculty of Veterinary Medicine for technical assistance.

My greatest appreciation to the very important persons in my life, Abah and Ma, (En. Abdullah Said and Azizah Awang) whose support and inspiration have sustained me and help me succeed in my study. And not forgetting my brothers and sisters for their support. Last but not least, to all others not mentioned but whose help has been tremendous, I express my sincere thanks and appreciation.



I certify that an Examination Committee met on 19th November, 2003 to conduct the final examination of Azmahani Abdullah on her Master of Science thesis entitled "Effects of *Andrographis paniculata*, *Carica papaya* and *Cassia alata* Extracts on the Reproductive Systems of Mice" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

JOHNSON STANSLAS, Ph.D.

Lecturer
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

MUHAMMAD NAZRUL HAKIM ABDULLAH, Ph.D.

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Member)

DATO' ABDUL SALAM ABDULLAH, Ph.D.

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

SABRINA SUKARDI, Ph.D.

Lecturer
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Member)

WAN NORDIN WAN MAHMUD, Ph.D.

Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)



GULAM RUSUL RAHMAT ALI, Ph.D.
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 05 MAR 2004

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as partial fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

MOHD NAZRUL HAKIM ABDULLAH, Ph. D

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

DATO' ABDUL SALAM ABDULLAH, Ph. D

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

SABRINA SUKARDI, Ph. D

Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

WAN NORDIN WAN MAHMUD, Ph. D.

Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)



AINI IDERIS, Ph.D.

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 12 MAR 2004



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



AZMAHANI ABDULLAH

Date: 13 MAR 2004

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xix
CHAPTER	
1 INTRODUCTION	
1.1 Population	1
1.2 Contraception	2
1.2.1 Contraceptive Methods for Women	2
1.2.2 Contraceptive Methods for Men	4
1.3 Herbs, Fertility and Contraception	5
1.4 Objectives of Study	6
2 LITERATURE REVIEW	
2.1 The Experimental Plants	7
2.1.1 <i>Andrographis paniculata</i>	7
2.1.1.1 Description	7
2.1.1.2 Chemistry	8
2.1.1.3 Folk Medicine	10
2.1.2 <i>Carica papaya</i>	12
2.1.1.1 Description	12
2.1.1.2 Chemistry	12
2.1.1.3 Folk Medicine	13
2.1.3 <i>Cassia alata</i>	15
2.1.1.1 Description	15
2.1.1.2 Chemistry	15
2.1.1.3 Folk Medicine	16
2.2 Regulation of Female Fertility	18
2.2.1 The Female Reproductive Tract of the Mouse	18
2.2.2 Oogenesis	20
2.2.3 Hormonal Regulation of the Ovarian Activity	22
2.3 Regulation of Male Fertility	24
2.3.1 The Male Reproductive Tract of the Mouse	24
2.3.2 Spermatogenesis	27
2.3.3 Hormonal Control of Spermatogenesis	30



2.4	Plants as Sources of Anti-fertility Agents	33
2.4.1	Mechanism of Action	33
2.5	Clomiphene Citrate	37
3	MATERIALS AND METHODS	
3.1	Plant Material	39
3.2	Preparation of the Extract	39
3.3	Fertility Testing on Female Mice	40
3.3.1	Experimental Animals	40
3.3.2	Experimental Design	40
3.3.3	Fertility Test	42
3.3.4	Body and Organ Weight	42
3.3.5	Histology	43
3.4	Fertility Testing on Male Mice	43
3.4.1	Experimental Animals	43
3.4.2	Experimental Design	44
3.4.3	Fertility Test	46
3.4.4	Blood Hormone Level	46
3.4.4.1	Blood Collection	46
3.4.4.2	ELISA	47
3.4.5	Body and Organ Weight	48
3.4.6	Sperm Profile	49
3.4.6.1	Sperm Collection	49
3.4.6.2	Concentration of Spermatozoa	49
3.4.6.3	Sperm Morphology	50
3.4.7	Histology	50
3.4.7.1	Tissue Processing	50
3.4.7.2	Quantitative Histology	51
3.5	Statistical Analysis	53
4	THE EFFECTS OF <i>ANDROGRAPHIS PANICULATA</i>, <i>CARICA PAPAYA</i> AND <i>CASSIA ALATA</i> ON THE REPRODUCTIVE SYSTEM OF FEMALE MICE	
4.1	Introduction	54
4.2	Results	55
4.3.1	Body and Organ Weight	55
4.3.2	Fertility Testing	57
4.3.3	Histological Examination	58
4.3	Discussion	63
4.4	Conclusion	67
5	THE EFFECTS OF <i>ANDROGRAPHIS PANICULATA</i>, <i>CARICA PAPAYA</i> AND <i>CASSIA ALATA</i> ON THE REPRODUCTIVE SYSTEM OF MALE MICE	
5.1	Introduction	68
5.2	Results	69



5.2.1	Body Weight	69
5.2.2	Organs Weight	72
5.2.3	Fertility Testing	75
5.2.4	Sperm profile	80
5.2.5	Testosterone	87
5.3	Discussion	88
5.4	Conclusion	92
6	HISTOLOGICAL EXAMINATIONS OF THE MALE REPRODUCTIVE SYSTEM TREATED WITH <i>ANDROGRAPHIS PANICULATA</i>, <i>CARICA PAPAYA</i> AND <i>CASSIA ALATA</i>	
6.1	Introduction	93
6.2	Results	95
6.2.1	Histological Examination	95
6.2.2	Quantitative Histology	101
6.3	Discussion	111
6.4	Conclusion	114
7	GENERAL DISCUSSION, CONCLUSION AND SUGGESTION FOR FURTHER RESEARCH	
7.1	General Discussion and Conclusion	115
7.2	Suggestion for Further Research	120
	BIBLIOGRAPHY	122
	APPENDICES	129
	BIODATA OF THE AUTHOR	138



LIST OF TABLES

Table	Page
1. Chemical constituents of <i>Andrographis paniculata</i>	8
2. Chemical constituents of <i>Carica papaya</i>	12
3. Chemical constituents of <i>Cassia alata</i>	15
4. Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on body and ovary organ weights during treatment	55
5. Fertility test on female mice for 30 days treatment	57
6. Effect on male body weights (g) of all treated groups during treatment	69
7. Organ weight (mg/100 g body weight) of male mice in all treatment groups	71
8. Fertility test and litter numbers of female mice during treatment	79
9. Effect of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on sperm count and motility during treatment	80
10. Serum testosterone (ng/ml) of all treatment groups	87
11. Mean volumetric proportion (%) of tubular elements, lumen and testicular space during treatment	101
12. Mean volumetric proportion (%) of spermatogonium, spermatocyte spermatid and spermatozoa during treatment	105
13. Mean volumetric proportion (%) of Sertoli cell, degenerated cells, membrane and interstitial cells	109



LIST OF FIGURES

Figure		Page
1.	Chemical structures of andrographolide and neoandrographolide	9
2.	<i>Andrographis paniculata</i>	11
3.	<i>Carica papaya</i>	14
4.	Anthraquinone (C ₁₅ H ₁₀ O ₅)	15
5.	<i>Cassia alata</i>	17
6.	Reproductive organs of female mice	19
7.	Events during oogenesis	21
8.	Regulation of ovarian function	23
9.	Reproductive organs of male mice	26
10.	Flowchart of events of spermatogenesis, showing the relative positioning of the various spermatogenic cells	29
11.	Hormonal regulation of testicular function (the brain-testicular axis)	32
12.	Structural formula of Clomiphene citrate	37
13.	Diagrammatic representation of experimental design for females	41
14.	Diagrammatic representation of experimental design for males	45
15.	Body weight (g) changes of female mice during treatment	56
16.	Section of the ovary of a mice (control) showing corpus luteum and preovulatory follicles	59
17.	Section of the ovary of a mice (Clomiphene treated) showing granuloma nodules in interstitium and atretic follicles	61
18.	Section of the ovary of a mice (Clomiphene treated) showing atretic follicles	61
19.	Section of the ovary of a mice (treated with extract <i>Carica papaya</i>) showing atretic follicles	62



20.	Section of the ovary of a mice (treated with extract <i>Andrographis paniculata</i>) showing atretic follicles	62
21.	Section of the ovary of a mice (treated with extract <i>Cassia alata</i>) showing normal developing follicles	63
22.	Body weight (g) changes of male mice during treatment	69
23.	Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on testis weight (mg/100g body weight) during treatment	73
24.	Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on epididymis weight (mg/100g body weight) during treatment	74
25.	Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on seminal vesicle weight (mg/100g body weight) during treatment	74
26.	Estrous stage showing large cornified, a nucleated epithelial cells	75
27.	Metestrous stage showing a few cornified epithelial cells and leucocytes	76
28.	Diestrus stage showing many leucocytes	77
29.	Proestrous stage showing many nucleated epithelial cells	78
30.	Effect of all extracts treated groups on sperm count during treatment	83
31.	Effect of all extracts treated groups on sperm count during treatment (clomiphene citrate treated group not included)	83
32.	Normal sperm from control mice	84
33.	Sperm showing defects in morphology abnormalities after Clomiphene treatment	84
34-37	Showing abnormalities in neck of sperm (clomiphene treated groups)	85
38.	Sperm showing defects in morphology after Clomiphene citrate treatment. SH, separated head	86
39.	Sperm showing defects in morphology after Clomiphene citrate treatment. ST, separated tail	86



40.	Cross section (C.S.) of normal mice testis	95
41.	C.S. of testis from Clomiphene citrate treated group after 60 days. Most seminiferous tubules are devoid of spermatozoa and degenerated spermatids are more abundant with some slouching into the lumen	96
42.	C.S. of testis from Clomiphene citrate treated group after 90 days. Most seminiferous tubules are left with spermatogonium and spermatocyte only and left with large lumens. Note the more abundant testicular space between the seminiferous tubules and the atrophied appearance of the seminiferous tubules	97
43.	C.S. of testis from <i>Carica papaya</i> treated group after 90 days showing some degenerated cells which are mostly round spermatids. Note the more abundant testicular space between the seminiferous tubules	98
44.	C.S. of testis from <i>Andrographis paniculata</i> extract	99
45.	C.S. of testis from <i>Cassia alata</i> extract	100
46.	Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on mean volumetric proportion (%) of tubular elements during treatment	102
47.	Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on mean volumetric proportion (%) of lumen during treatment	103
48.	Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on mean volumetric proportion (%) of testicular space during treatment	104
49.	Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on mean volumetric proportion (%) of spermatogonium during treatment	106
50.	Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on mean volumetric proportion (%) of spermatocyte during treatment	106
51.	Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on mean volumetric proportion (%) of spermatid during treatment	107
52.	Effects of <i>Andrographis paniculata</i> , <i>Carica papaya</i> and <i>Cassia alata</i> ethanol extracts on mean volumetric proportion (%) of spermatozoa during treatment	108



53. Effects of *Andrographis paniculata*, *Carica papaya* and *Cassia alata* ethanol extracts on mean volumetric proportion (%) of degenerated cell during treatment 110



LIST OF ABBREVIATIONS

b.w.	body weight
$^{\circ}\text{C}$	Degress Celcius
g	gram
M	molar
mg	miligram
mm	millimeter
ml	militers
μl	microliters
μm	micrometer
μg	microgram
mw	miliwolt
s.e.m	standard error of mean
pH	$-\log_{10}[\text{H}^+]$
rpm	rotation per minute
ng	nanogram
p	probability of an event due to chance alone
ELISA	enzyme-linked immunosorbent assay
Min	minute
G.I.	Gastrointestinal
FSH	follicle-stimulating hormone



CHAPTER 1

INTRODUCTION

1.1 Population Explosion

The single outstanding fact about world population during the past two hundred years has been the rapidity of its growth. Since 1750, there has been a rapid and accelerating expansion in population and more recently termed 'population explosion'. Currently, according to the World Health Organization (WHO), 3.7 human beings are born every second: the net increment works out to be at 222 per minute, 13 260 per hour, 318 240 per day, or nearly 2.25 million per week (Robinson, 1981).

The limits to growth have been analysed recently by a team of scientists led by Professor Dennis Meadows from the Massachusetts Institute of Technology, with the help of models and computers. These models were constructed to investigate 'the five major trends of global concern – accelerating industrialization, rapid population growth, widespread malnutrition, depletion of non-renewable resources, and a deteriorating environment' (Llewellyn-Jones, 1975). The recent and unprecedented population explosion is of concern not only to the demographer but to the politician, economist, geographer, sociologist, and medical scientist among many others, all of whom are very much concerned with ensuring that the sheer quantity of human beings will not diminish the quality of human life (Robinson, 1981).



Global search of anti-fertility agents is going to tackle the problem of 'Population Explosion'. Many hormonal drugs are available for the purpose but they are not free from side effects. Hence, the search for a suitable product from indigenous medicinal plants is proposed which could be effectively used in place of the synthetic or hormonal (Hoffman, 1990).

1.2 Contraceptive

There are several methods of contraception today. However, an ideal contraceptive that is safe, inexpensive, totally effective, easily reversible and without side effects has yet to be developed. Oral contraceptives, intrauterine devices, male and female sterilizations, barrier contraceptives, medical termination of pregnancy, rhythm and symptothermal methods and *coitus interruptus* are presently being used to control births (Salunkhe *et al.*, 1989).

1.2.1 Contraceptive Methods for Women

The pill is a very simple and easy to use method of birth control. The female sexual partner simply swallows one pill each day for 20 or 21 days. During that month she will not get pregnant. The pill works by keeping the woman's ovaries from releasing an egg as long as she is taking the pills according to the prescribed regimen. The pill contains the female hormones estrogen and progesterone which a woman's body produces when she is pregnant. These hormones keep her body from releasing eggs for as long as she takes birth control pills. The eggs simply do not mature and thus are not released by the ovaries. For a few women, the pills sometimes produce temporary and mildly unpleasant

side effects. Some women experience mild nausea or stomach upsets when taking the first few pills. This goes away after the pills have been used for a month or so (Zawacki, 1971). Other problems that need further study to establish their association with the pill are an increased risk of urinary tract infections, gall bladder disease, liver disease and tumors, and birth defects if pill use is continued into pregnancy. Prolonged infertility is also possible after the pill is discontinued, but whether it is caused by the pill itself or masked by the artificial menses during use is unclear. Evidence so far shows pill users to have a lower risk of benign breast disease, fewer ovarian cysts, and unchanged risk of cervical cancer (Hauser, 1979).

Chemical impregnated intra-vaginal plugs have been used for contraceptive purpose by many primitive tribes. Currently, there are over 50 suppositories, pessaries, gels, or foams available. The solid vaginal suppositories contain spermicide in a base of soap, gelatin or cocoa-butter which is designed to melt at body temperature. The vagitory is inserted just prior to coitus. The gels which are made up in a water-soluble gelatinous base disperse easily in the vagina, and usually serve as an additional protection if the male uses a condom, or in conjunction with a vaginal diaphragm. The gel or cream is usually introduced high into the vagina through a plastic plunger applicator which is provided in the package. The foams are either in foaming tablet form or in containers from which the appropriate amount can be introduced into the vagina (Llewellyn-Jones, 1975). Disadvantages are the leakage of fluid from the vagina, the need to wait while the content of the vaginal suppositories melts, and poor effectiveness, usually due to poor quality, inadequate quantity, or disregard for the required waiting period (Hauser, 1979).



1.2.2 Contraceptive Methods for Men

Apart from abortion and infanticide, coitus interruptus is probably the oldest method of birth control (Llewellyn-Jones, 1975). The method requires no supplies, expense, preparation, or assistance from physicians. On the other hand, it demands practice, male-control, and considerable motivation. It may be reliable in highly motivated couples. Failures can result from the escape of semen before ejaculation, from semen deposited externally near the vagina, or from simple delay in withdrawal (Hauser, 1979). Its greatest disadvantage is that it is difficult to practice because it interrupts the act of sex just at a point when human nature desires no interruption (Zawacki, 1971).

A cylindrical sheath that envelops the penis, the condom has the advantage of being cheap, simple to use and can be obtained without prescription from a physician. However, it has the disadvantages of being distracting and dulling sensation. Moreover in some cultures, it is associated with prostitution because of its prophylactic role in venereal diseases. It is effective if properly used and carefully manufactured. Rubber condoms deteriorate with time, particularly in sunlight and heat, and therefore have a limited shelf life, especially in tropical countries (Hauser, 1979). Male sterilization (called vasectomy) is quite a simple operation that can be done in a doctor's office or a clinic. The man merely goes home and rests for a few of days. The procedure involves the removal of a small section of the two tubes that carry the male sperm to the penis. This procedure will not change in his nature or his sexual desire (Zawacki, 1971).



1.3 Herbs, Fertility and Contraception

Several approaches are being used in different parts of the world to control human fertility. Most of the currently used methods of birth control are associated with certain risks and side effects. As such, an ideal contraceptive has yet to be developed. Many plants have been used in traditional medicine as contraceptive agents.

New and improved methods of contraception are being developed. For example, long-acting injectable preparations, postcoital drugs, low-dose oral contraceptives, biphasic and triphasic pills, paper pills, sperm and ova pills. The currently used methods of contraception are associated with risks and side effects. For example, the use of oral contraceptives is associated with some circulatory system diseases including thromboembolism, myocardial infarction and hypertension. Endometrial bleeding, pelvic infections, pelvic inflammatory disease and ectopic pregnancy are frequently observed with use of the intrauterine device (IUD) (Stephen *et al.*, 1994).

Great attention is being given to plants with potential anti-fertility properties. These may act through effects on sperm motility and viability, implantation of the fertilized egg or a rejection effect within the uterus. The biochemistry of these pathways is complex, and the study of plants having such effects is revealing new mechanisms all the time. The planetary crisis that is upon us has the population explosion as a major component, and the World Health Organization has put great emphasis on the search for a safe, cheap and socially acceptable form of contraception (Hoffman, 1990).