



**EVALUATION OF *IN VITRO* ANTIDIABETIC MECHANISMS OF
METHANOLIC EXTRACTS FROM FOUR *Ficus deltoidea* Jack VARIETIES**

By

NURSHIEREN YAHAYA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

March 2020

FBSB 2020 11

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia





*I dedicate this thesis for my families, future generations
and the enthusiasts of Malaysia's herbal medicines*

@COPYRIGHT UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

EVALUATION OF *IN VITRO* ANTIDIABETIC MECHANISMS OF METHANOLIC EXTRACTS FROM FOUR *Ficus deltoidea* Jack VARIETIES

By

NURSHIEREN YAHAYA

March 2020

Chairman: Muhajir Hamid, PhD

Faculty: Biotechnology and Biomolecular Sciences

Diabetes Mellitus has been a global metabolic disorder and affected people of various ages. In this research, four varieties of methanolic *Ficus deltoidea* extracts containing natural products that possess anti-diabetic properties has been studied. The study was carried out because antidiabetic synthetic drugs have been reported to have negative side effects on diabetes patients. Therefore, the study was done to elucidate insulin-secreting activity towards the beta pancreatic cell, to elucidate glucose uptake activity in insulin targeted cells, to determine the potential of insulin-sensitizing activity in adipocytes cells and to determine the potential of antiglycation activities of four methanolic *Ficus deltoidea* varieties. The viability study which manifests viability assay has shown that methanolic *Ficus deltoidea* variety *deltoidea*, *angustifolia* and *motleyana* does not toxic towards the treated cells with the highest viability were portrayed by the highest viability can be seen when treated with 500 µg/ml of methanolic extract *Ficus deltoidea* variety *angustifolia* with 151.42 ± 0.08 percent ($p < 0.001$) on beta pancreatic cell, 1000 µg/ml of methanolic extract *Ficus deltoidea* variety *angustifolia* with 122.66 ± 1.92 percent ($p < 0.01$) on L6 muscle cell and the highest viability can be seen at 100 µg/ml when treated with methanolic *Ficus deltoidea* variety *motleyana* at 110.58 ± 3.45 percent on 3T3F442A adipocytes cell but not significant. Besides, insulin secretion study conducted with insulin secretion assay showed methanolic extracts of *Ficus deltoidea* varieties can stimulate insulin release from pancreatic cells through the K_{ATP} independent pathway. The highest insulin secretion was shown by methanolic extract *Ficus deltoidea* variety *deltoidea* with 6.24 fold ($p < 0.001$). In addition, the extracts able to stimulate glucose uptake in basal and insulin-mediated state where the highest uptake can be seen in insulin-mediated state of L6 muscle cell treated with methanolic extract *Ficus deltoidea* variety *motleyana* with 3.65 fold ($p < 0.001$) when compared to control and 2.90 fold ($p < 0.001$) when compared to insulin 100 nM. Apart from that the highest uptake can be seen in insulin-mediated state of 3T3F442A adipocyte cell treated with methanolic extract *Ficus deltoidea* variety *motleyana* with 1.39 fold ($p < 0.001$) when compared to control and 1.27 fold ($p < 0.001$) when compared to insulin 100 nM. All of the extracts follow the phosphatidylinositol-3-kinase-independent

pathway except for *Ficus deltoidea* variety *motleyana* where when treated to 3T3F442A cell, the extract does not follow both of phosphatidylinositol-3-kinase-dependent and independent pathway. While in gene expression study through using Quantigene assay kit, showed methanolic *Ficus deltoidea* variety *bilobata* as the highest expression for phosphatidylinositol-3-kinase-dependent pathway with 1.24 fold in 6 hours while *Ficus deltoidea* variety *motleyana* showed the highest expression for phosphatidylinositol-3-kinase-independent pathway with 1.21 fold in 6 hours. All four varieties at a concentration of 100 μ M in insulin-mediated state shown a significant glucose uptake activity in insulin resistance cell with the best secretion of leptin and inhibition of tumor necrosis factor- α was significantly recorded by methanolic extract *Ficus deltoidea* variety *angustifolia* with 2.56 fold ($p < 0.001$) and 0.73 fold ($p < 0.001$). Last but not least, the highest antiglycation activity was exhibited by methanolic *Ficus deltoidea* variety *bilobata* with 78.92 percent of advanced glycation endproducts inhibition formation along with the highest detection of fructosamine inhibition with 86.40 percent reduction. While *Ficus deltoidea* variety *deltoidea* showed significant results with protein thiol preservation of 6.14 fold ($p < 0.001$) and protein carbonyl reduction with 1.76 fold ($p < 0.001$). Total phenolic content assay showed that methanolic *Ficus deltoidea* variety *bilobata* to have highest phenolic content of 299.78 ± 2.88 mg GAE/ g of extract ($p < 0.001$). The varieties also showed significant antioxidant activities. In conclusion, methanolic *Ficus deltoidea* extract showed that most extract follow independently of ATP-sensitive potassium channel in insulin secretion activity. Apart from that, the extracts follow glucose uptake phosphatidylinositol-3-kinase independent pathway, able to sensitize insulin and inhibit advanced glycation formation. Therefore, it is suggested that methanolic *Ficus deltoidea* extracts can be a promising candidate to be used as phytopharmaceutical agents to manage diabetes mellitus.

Keywords : Antidiabetic, *Ficus deltoidea*, insulin, glucose uptake, antiglycation

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

KAJIAN *IN VITRO* MEKANISMA ANTIDIABETIK OLEH EMPAT JENIS METANOLIK EKSTRAK VARIATI *Ficus deltoidea* Jack

Oleh

NURSHIEREN YAHAYA

Mac 2020

Pengerusi: Muhajir Hamid, PhD

Fakulti: Bioteknologi dan Sains Biomolekul

Kencing manis telah menjadi gangguan metabolisme global dan boleh didapati dalam pelbagai peringkat umur. Dalam kajian ini, empat jenis ekstrak standard metanolik *Ficus deltoidea* mengandungi produk semula jadi yang mempunyai sifat anti-kencing manis. Kajian telah dijalankan kerana ubat sintetik antidiabetes telah dilaporkan mempunyai kesan sampingan yang negatif kepada pesakit kencing manis. Oleh itu, kajian ini telah dilakukan untuk menjelaskan aktiviti insulin ke arah sel pankreas beta, mendapatkan semula aktiviti pengambilan glukosa dalam sel sasaran insulin, mengkaji potensi aktiviti insulin-sensitif dalam sel adiposit dan mengkaji potensi aktiviti antiglikasi empat jenis ekstrak standard metanolik *Ficus deltoidea*. Kajian daya maju telah menunjukkan bahawa ekstrak standard metanolik *Ficus deltoidea* jenis *deltoidea*, *angustifolia* dan *motleyana* tidak toksik terhadap sel yang dirawati. Daya maju yang paling tinggi direkodkan ialah 500 µg/ml ekstrak metanolik *Ficus deltoidea* variati *angustifolia* dengan peratusan 151.42 ± 0.08 persen ($p < 0.001$) ke atas sel pankreatik beta, 1000 µg/ml ekstrak metanolik *Ficus deltoidea* variati *angustifolia* dengan peratusan 122.66 ± 1.92 persen ($p < 0.01$) ke atas sel otot L6 dan 100 µg/ml ekstrak metanolik *Ficus deltoidea* variati *motleyana* dengan peratusan 110.58 ± 3.45 persen ke atas 3T3F442A sel adiposit tetapi tidak menunjukkan kesan signifikan. Selain itu, kajian rembesan insulin menggunakan metod kajian perembesan insulin menunjukkan ekstrak metanolik *Ficus deltoidea* boleh merangsang pembebasan insulin dari sel pankreas melalui laluan bergantung K_{ATP} . Penghasilan insulin yang paling tinggi ditunjukkan oleh ekstrak metanolik *Ficus deltoidea* variati *deltoidea* dengan 6.24 kali ganda ($p < 0.001$). Selain itu, ekstrak metanolik daripada semua variati *Ficus deltoidea* mampu meningkatkan pengambilan glukosa di peringkat basal dan pengantaraan insulin di mana pengambilan glukosa yang paling tinggi adalah dalam keadaan pengantaraan glukosa di dalam sel otot L6 oleh metanolik ekstrak *Ficus deltoidea* variati *motleyana* dengan 3.65 kali ganda ($p < 0.001$) apabila dibandingkan dengan kumpulan kawalan dan 2.90 kali ganda ($p < 0.001$) apabila dibandingkan dengan insulin 100 nM. Selain daripada itu, pengambilan glukosa tertinggi di dalam sel 3T3F442A adiposit ditunjukkan oleh ekstrak metanolik *Ficus deltoidea* variati *motleyana* dengan 1.36

kali ganda ($p < 0.001$) dibandingkan dengan kumpulan kawalan dan 1.27 kali ganda ($p < 0.001$) apabila dibandingkan dengan insulin 100 nM. Semua ekstrak mampu merangsang pengambilan glukosa di dalam keadaan basal dan pengantaraan insulin dan mengikuti laluan bebas phosphatidylinositol-3-kinase kecuali untuk *Ficus deltoidea* variati *motleyana* di mana apabila diuji sel 3T3F442A, ekstrak tidak mengikuti kedua-dua laluan bebas phosphatidylinositol-3-kinase. Dalam kajian ekspresi gen, ekstrak standard metanolik *Ficus deltoidea* variati *bilobata* menunjukkan keputusan tertinggi untuk laluan phosphatidylinositol-3-kinase dengan keputusan 1.24 kali ganda dalam masa 6 jam manakala *Ficus deltoidea* jenis *motleyana* menunjukkan keputusan tertinggi bagi laluan bebas phosphatidylinositol-3-kinase dengan keputusan 1.21 kali ganda dalam masa 6 jam. Kesemua empat jenis pada kepekatan 100 μM dalam keadaan bersama insulin telah menunjukkan aktiviti pengambilan glukosa yang signifikan dalam sel rintangan insulin dengan rembesan leptin dan perencatan faktor tumor nekrosis- α telah direkodkan dengan ketara dan terbaik oleh ekstrak metanolik *Ficus deltoidea* variati *angustifolia* dengan 2.56 kali ganda ($p < 0.001$) serta 0.73 kali ganda ($p < 0.001$). Akhir sekali, aktiviti antiglikasi tertinggi telah dipamerkan oleh standard metanolik *Ficus deltoidea* jenis *bilobata* dengan 78.92 persen penghalangan kepada pembentukan produk glikasi lanjutan manakala pengesanan tertinggi perencatan fruktosamin menunjukkan 86.40 persen penurunan. Manakala *Ficus deltoidea* jenis *deltoidea* menunjukkan keputusan yang signifikan dengan pengesanan protein thiol bersama keputusan 6.14 kali ganda ($p < 0.001$) dan peratusan penghalangan pembentukan protin karbonil iaitu sebanyak 1.76 kali ganda ($p < 0.001$). Ujian Jumlah Kandungan Fenolik pula mencatatkan ekstrak metanolik *Ficus deltoidea* variati *bilobata* mempunyai kandungan fenolik tertinggi iaitu 299.78 ± 2.88 mg GAE/ g ekstrak ($p < 0.001$). Semua ekstrak juga menunjukkan signifikansi di dalam aktiviti antioksidan. Sebagai kesimpulan, kebanyakan ekstrak metanolik daripada variati *Ficus deltoidea* menunjukkan laluan bebas daripada saluran potasium ATP-sensitif di dalam aktiviti perembesan insulin. Selain itu, aktiviti pengambilan glukosa menunjukkan kebanyakan ekstrak metanolik *Ficus deltoidea* mengikuti laluan bebas fosfatidilinositol-3-fosfat. Ekstrak tersebut juga dapat menjadikan sel lebih sensitif terhadap insulin dan mampu menghalang pembentukan produk glikasi lanjutan. Oleh itu, adalah dicadangkan bahawa ekstrak metanolik *Ficus deltoidea* boleh menjadi calon yang menjanjikan kesan positif sebagai agen fitofarmaseutikal bagi membendung penyakit kencing manis.

Kata kunci : Antidiabetik, *Ficus deltoidea*, insulin, pengambilan glukosa, antiglikasi

ACKNOWLEDGEMENTS

I would like to thank Allah SWT for the opportunities given to me to embark on the doctoral study and ease my research journey. There are no words to describe how delighted I am to be able to complete this research journey. In addition to that, I would like to thank my supervisor, Associate Professor Dr. Muhajir Hamid and co-supervisor Dr. Zainah Adam as well as Professor Dr. Norazizah Shafee for their advice, guidance, reference, and moral support during the study period. Their kind words and technical resources have made me thrive and persevere in this journey. Apart from that, I also would like to thank Universiti Malaysia Perlis and the Ministry of Higher Education for the scholarship given which made the study possible. I also would like to thank the Ministry of Agriculture for providing grants and Universiti Putra Malaysia for providing financial aid to postgraduate students. Without all these financial resources I will probably not be able to finish the study in stipulated time. In addition, I would like to thank Nur Sumirah Mohd Dom for sharing knowledge and for being a helpful colleague during my study years. Besides that, I would like to thank my family and friends for their endless support and prayers during my study particularly to my husband Mohamad Helmi Ismail, my parents Yahaya Mat Ali and Zakiah Abdul Rahman as well as my mother in law, Yusaidar M. Azhar. Not to forget my son Muhammad Harraz Shafiy bin Mohamad Helmi whom always there for me during the ups and downs of my study. Last but not least, I would like to thank my siblings Naeiranie Yahaya and Nazlizulidris Yahaya for helping hands during my final years of completing the study. Thank you, everyone, for believing in me.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Muhajir Hamid, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Norazizah Shafee, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Zainah Adam, PhD

Research Officer
Medical Technology Department
Malaysian Nuclear Agency
(Member)

ZALILAH MOHD SHARIFF, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 16 July 2020

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xx
CHAPTER	
1 INTRODUCTION	
1.1 Background of study	1
1.2 Statement of problem	2
1.3 Justification of the study	2
1.4 Hypothesis of the study	3
1.5 Objectives of the study	4
2 LITERATURE REVIEW	
2.1 Definition of diabetes	5
2.1.1 Type 1 diabetes	5
2.1.2 Type 2 diabetes	6
2.1.3 Hybrid form of diabetes	6
2.1.4 Diabetes in pregnancy	7
2.2 Insulin resistance	7
2.3 Glucose homeostasis	9
2.4 Diabetes complications	9
2.5 Present drugs to treat diabetes	10
2.6 Antidiabetic herbal plants	11
2.7 Compounds that exhibit antidiabetic properties	11
2.8 Insulin secretion and glucose uptake mechanisms	12
2.8.1 Phosphoinositol-3-kinase pathway	15
2.9 Advanced glycation endproducts	15
2.10 <i>Ficus deltoidea</i> as antidiabetic herbal medicine	17
3 CELL VIABILITY EFFECT AND INSULINOTROPIC ACTIVITIES OF FOUR METHANOLIC <i>FICUS DELTOIDEA</i> VARIETIES	
3.1 Introduction	21
3.2 Materials and methods	23
3.2.1 Plant extract preparation	23
3.2.2 Cell culture	23
3.2.3 Cell viability assay	23

	3.2.4	Insulin secretion assay	24
	3.2.5	Elucidation of insulin secretion mechanisms	24
	3.2.6	Statistical analysis	25
3.3	Results		26
	3.3.1	Cell viability effect on beta pancreatic cell	26
	3.3.2	Cell viability effect on L6 muscle cell	28
	3.3.3	Cell viability effect on 3T3F442A adipocytes cell	30
	3.3.4	Insulin secretion activity of methanolic <i>Ficus deltoidea</i> extracts from four varieties	32
	3.3.5	Elucidation of insulin secretion mechanisms of methanolic <i>Ficus deltoidea</i> extracts from four varieties	37
3.4	Discussion		41
3.5	Conclusion		43
4	GLUCOSE UPTAKE MECHANISMS AND INSULIN SENSITIZING EFFECTS OF FOUR METHANOLIC <i>FICUS DELTOIDEA</i> VARIETIES		
	4.1	Introduction	44
	4.2	Materials and methods	46
	4.2.1	Glucose uptake into L6 muscle cells and 3T3F442A adipocytes cells	46
	4.2.2	Glucose uptake inhibition assay	47
	4.2.3	Quantification of genes using Quantigene	47
	4.2.4	Insulin sensitizing activity of methanolic <i>Ficus deltoidea</i> varieties in adipocytes cells	49
	4.2.5	Statistical analysis	49
	4.3	Results	49
	4.3.1	Glucose uptake into L6 muscle cells and 3T3F442A adipocytes cells	49
	4.3.2	Glucose uptake inhibition assay	59
	4.3.3	Quantification of genes using quantigene	75
	4.3.4	Glucose uptake in insulin resistance cells	77
	4.3.5	Determination of Leptin in insulin resistance cells	81
	4.3.6	Determination of TNF- α in insulin resistance cells	85
	4.4	Discussion	89
	4.5	Conclusion	94

5	INHIBITION OF ADVANCED GLYCATION ENDPRODUCTS BY FOUR METHANOLIC <i>FICUS DELTOIDEA</i> VARIETIES	
5.1	Introduction	95
5.2	Materials and methods	96
	5.2.1 In vitro antiglycation of bovine serum albumin (24 hour)	96
	5.2.2 In vitro antiglycation of bovine serum albumin (14 and 28 days)	96
	5.2.3 Determination of fructosamine	96
	5.2.4 Determination of thiol group	97
	5.2.5 Determination of protein carbonyl content	97
	5.2.6 Total phenolic content assay	97
	5.2.7 DPPH radical scavenging activity of <i>Ficus deltoidea</i> varieties	97
	5.2.8 Power reducing assay	98
	5.2.9 Statistical analysis	98
5.3	Results	99
	5.3.1 In vitro antiglycation of bovine serum albumin (24 hours)	99
	5.3.2 In vitro antiglycation of bovine serum albumin (14 and 28 days)	100
	5.3.3 Determination of fructosamine	105
	5.3.4 Determination of thiol group	110
	5.3.5 Determination of protein carbonyl content	115
	5.3.6 Total phenolic content assay	120
	5.3.7 DPPH radical scavenging activity of <i>Ficus deltoidea</i> varieties	121
	5.3.8 Power reducing assay	122
5.4	Discussion	123
5.5	Conclusion	126
6	SUMMARY, GENERAL CONCLUSION, AND RECOMMENDATION FOR FUTURE RESEARCH	
6.1	Summary of the study	127
6.2	Limitation and recommendation of future research	129
6.3	Conclusion of the study	129
	REFERENCES	130
	APPENDICES	165
	BIODATA OF STUDENT	174
	LIST OF PUBLICATIONS	175

LIST OF TABLES

Table		Page
3.1	Effect of methanolic <i>Ficus deltoidea</i> extracts and glybenclamide on BRIN BD11 beta pancreatic cell	27
3.2	Effect of methanolic <i>Ficus deltoidea</i> extracts and metformin on L6 muscle cell	29
3.3	Effect of methanolic <i>Ficus deltoidea</i> extracts and metformin on 3T3F442A adipocytes cell	31
4.1	The expression of mRNA level of four genes when treated with control, Metformin, Insulin 100 nM and four methanolic <i>Ficus deltoidea</i> varieties	76
5.1	Antiglycation activity of bovine serum albumin in 24 hours when treated with quercetin and methanolic <i>Ficus deltoidea</i> varieties	99
5.2	Total phenolic content of Quercetin and four varieties of methanolic <i>Ficus deltoidea</i> extracts	120
5.3	DPPH radical scavenging activity of methanolic <i>Ficus deltoidea</i> varieties.	121

LIST OF FIGURES

Figure		Page
2.1	Insulin receptor signaling cascade (Carnagarin <i>et. al.</i> 2015)	14
2.2	Four varieties of <i>Ficus deltoidea</i> . a) <i>Ficus deltoidea</i> variety <i>deltoidea</i> b) <i>Ficus deltoidea</i> variety <i>angustifolia</i> c) <i>Ficus deltoidea</i> variety <i>bilobata</i> d) <i>Ficus deltoidea</i> variety <i>motleyana</i>	20
3.1	Effect of glybenclamide in insulin secretion on BRIN BD11 pancreatic beta cells for 1 hour incubation.	32
3.2	Effect of methanolic <i>Ficus deltoidea</i> variety <i>deltoidea</i> in insulin secretion on BRIN BD11 pancreatic beta cells for 1 hour incubation.	33
3.3	Effect of methanolic <i>Ficus deltoidea</i> variety <i>angustifolia</i> in insulin secretion on BRIN BD11 pancreatic beta cells for 1 hour incubation.	34
3.4	Effect of methanolic <i>Ficus deltoidea</i> variety <i>bilobata</i> in insulin secretion on BRIN BD11 pancreatic beta cells for 1 hour incubation	35
3.5	Effect of methanolic <i>Ficus deltoidea</i> variety <i>motleyana</i> in insulin secretion on BRIN BD11 pancreatic beta cells for 1 hour incubation.	36
3.6	Effect of four modulators on BRIN BD 11 treated with 500 µg/ml methanolic <i>Ficus deltoidea</i> var. <i>deltoidea</i> .	37
3.7	Effect of four modulators on BRIN BD 11 treated with 500 µg/ml methanolic <i>Ficus deltoidea</i> var. <i>angustifolia</i> .	38
3.8	Effect of four modulators on BRIN BD 11 treated with 500 µg/ml methanolic <i>Ficus deltoidea</i> var. <i>bilobata</i> .	39
3.9	Effect of four modulators on BRIN BD 11 treated with 500 µg/ml methanolic <i>Ficus deltoidea</i> var. <i>motleyana</i> .	40
4.1	Effect of metformin on basal and the insulin-mediated glucose uptake in L6 muscle cell	49
4.2	Effect of methanolic <i>Ficus deltoidea</i> var. <i>deltoidea</i> extract on basal and the insulin-mediated glucose uptake in L6 muscle cell	50
4.3	Effect of methanolic <i>Ficus deltoidea</i> var. <i>angustifolia</i> extract on basal and the insulin-mediated glucose uptake in L6 muscle cell	51
4.4	Effect of methanolic <i>Ficus deltoidea</i> var. <i>bilobata</i> extract on basal and the insulin-mediated glucose uptake in L6 muscle cell	52
4.5	Effect of methanolic <i>Ficus deltoidea</i> var. <i>motleyana</i> extract on basal and the insulin-mediated glucose uptake in L6 muscle cell	53

4.6	Effect of Rosiglitazone maleate on basal and the insulin-mediated glucose uptake in 3T3F442A adipocyte cell	54
4.7	Effect of methanolic <i>Ficus deltoidea</i> var. <i>deltoidea</i> extract on basal and the insulin-mediated glucose uptake in 3T3F442A adipocyte cell	55
4.8	Effect of methanolic <i>Ficus deltoidea</i> var. <i>angustifolia</i> extract on basal and the insulin-mediated glucose uptake in 3T3F442A adipocyte cell	56
4.9	Effect of methanolic <i>Ficus deltoidea</i> var. <i>bilobata</i> extract on basal and the insulin-mediated glucose uptake in 3T3F442A adipocyte cell	57
4.10	Effect of methanolic <i>Ficus deltoidea</i> var. <i>motleyana</i> extract on basal and the insulin-mediated glucose uptake in 3T3F442A adipocyte cell	58
4.11	The inhibitory effects of ALLN with methanolic <i>Ficus deltoidea</i> var. <i>deltoidea</i> extract in L6 muscle cell.	59
4.12	The inhibitory effects of LY294002 with methanolic <i>Ficus deltoidea</i> var. <i>deltoidea</i> extract in L6 muscle cell.	60
4.13	The inhibitory effects of ALLN with methanolic <i>Ficus deltoidea</i> var. <i>angustifolia</i> extract in L6 muscle cell.	61
4.14	The inhibitory effects of LY294002 with methanolic <i>Ficus deltoidea</i> var. <i>angustifolia</i> extract in L6 muscle cell.	62
4.15	The inhibitory effects of ALLN with methanolic <i>Ficus deltoidea</i> var. <i>bilobata</i> extract in L6 muscle cell.	63
4.16	The inhibitory effects of LY294002 with methanolic <i>Ficus deltoidea</i> var. <i>bilobata</i> extract in L6 muscle cell.	64
4.17	The inhibitory effects of ALLN with methanolic <i>Ficus deltoidea</i> var. <i>motleyana</i> extract in L6 muscle cell.	65
4.18	The inhibitory effects of LY294002 with methanolic <i>Ficus deltoidea</i> var. <i>motleyana</i> extract in L6 muscle cell.	66
4.19	The inhibitory effects of ALLN with methanolic <i>Ficus deltoidea</i> var. <i>deltoidea</i> extract on 3T3F442A adipocytes cell.	67
4.20	The inhibitory effects of LY294002 with methanolic <i>Ficus deltoidea</i> var. <i>deltoidea</i> extract on 3T3F442A adipocytes cell.	68
4.21	The inhibitory effects of ALLN with methanolic <i>Ficus deltoidea</i> var. <i>angustifolia</i> extract on 3T3F442A adipocytes cell.	69
4.22	The inhibitory effects of LY294002 with methanolic <i>Ficus deltoidea</i> var. <i>angustifolia</i> extract on 3T3F442A adipocytes cell.	70

4.23	The inhibitory effects of ALLN with methanolic <i>Ficus deltoidea</i> var. <i>bilobata</i> extract on 3T3F442A adipocytes cell.	71
4.24	The inhibitory effects of LY294002 with methanolic <i>Ficus deltoidea</i> var. <i>bilobata</i> extract on 3T3F442A adipocytes cell.	72
4.25	The inhibitory effects of ALLN with methanolic <i>Ficus deltoidea</i> var. <i>motleyana</i> extract on 3T3F442A adipocytes cell	73
4.26	The inhibitory effects of LY294002 with methanolic <i>Ficus deltoidea</i> var. <i>motleyana</i> extract on 3T3F442A adipocytes cell.	74
4.27	Effect of Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>deltoidea</i> extract treated on 3T3F442A insulin-resistant cells in glucose uptake activity.	77
4.28	Effect of Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>angustifolia</i> extract treated on 3T3F442A insulin-resistant cells in glucose uptake activity.	78
4.29	Effect of Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>bilobata</i> extract treated on 3T3F442A insulin-resistant cells in glucose uptake activity.	79
4.30	Effect of Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>motleyana</i> extract treated on 3T3F442A insulin-resistant cells in glucose uptake activity.	80
4.31	Effect of leptin secretion by 3T3F442A insulin-resistant cell when treated with Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>deltoidea</i> extract	81
4.32	Effect of leptin secretion by 3T3F442A insulin-resistant cell when treated with Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>angustifolia</i> extract	82
4.33	Effect of leptin secretion by 3T3F442A insulin-resistant cell when treated with Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>bilobata</i> extract.	83
4.34	Effect of leptin secretion by 3T3F442A insulin-resistant cell when treated with Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>motleyana</i> extract	84
4.35	Effect of TNF- α secretion by 3T3F442A insulin-resistant cell when treated with Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>deltoidea</i> extract	85
4.36	Effect of TNF- α secretion by 3T3F442A insulin-resistant cell when treated with Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>angustifolia</i> extract.	86

4.37	Effect of TNF- α secretion by 3T3F442A insulin-resistant cell when treated with Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>bilobata</i> extract.	87
4.38	Effect of TNF- α secretion by 3T3F442A insulin-resistant cell when treated with Rosiglitazone Maleate and methanolic <i>Ficus deltoidea</i> variety <i>motleyana</i> extract.	88
5.1	The effects of Quercetin on inhibition of advanced glycation endproducts formation (%) in BSA incubated with fructose.	100
5.2	The effects of methanolic <i>Ficus deltoidea</i> variety <i>deltoidea</i> extracts on inhibition of advanced glycation endproducts	101
5.3	The effects of methanolic <i>Ficus deltoidea</i> variety <i>angustifolia</i> extracts on inhibition of advanced glycation endproducts formation (%) in BSA incubated with fructose.	102
5.4	The effects of methanolic <i>Ficus deltoidea</i> variety <i>bilobata</i> extracts on inhibition of advanced glycation endproducts formation (%) in BSA incubated with fructose.	103
5.5	The effects of methanolic <i>Ficus deltoidea</i> variety <i>motleyana</i> extracts on inhibition of advanced glycation endproducts formation (%) in BSA incubated with fructose.	104
5.6	The effects of Quercetin on the level of fructosamine (mmol/ L) in BSA incubated with fructose.	105
5.7	The effects of methanolic <i>Ficus deltoidea</i> variety <i>deltoidea</i> extracts on the level of fructosamine (mmol/ L) in BSA incubated with fructose.	106
5.8	The effects of <i>Ficus deltoidea</i> variety <i>angustifolia</i> extracts on the level of fructosamine (mmol/ L) in BSA incubated with fructose.	107
5.9	The effects of <i>Ficus deltoidea</i> variety <i>bilobata</i> extracts on the level of fructosamine (mmol/ L) in BSA incubated with fructose.	108
5.10	The effects of <i>Ficus deltoidea</i> variety <i>motleyana</i> on the level of fructosamine (mmol/ L) in BSA incubated with fructose.	109
5.11	The effects of Quercetin on thiol group content (nmol/ mg protein) in BSA incubated with fructose.	110
5.12	The effects of methanolic <i>Ficus deltoidea</i> variety <i>deltoidea</i> extract on thiol group content (nmol/ mg protein) in BSA incubated with fructose.	111
5.13	The effects of methanolic <i>Ficus deltoidea</i> variety <i>angustifolia</i> extract on thiol group content (nmol/ mg protein) in BSA incubated with fructose.	112

5.14	The effects of methanolic <i>Ficus deltoidea</i> variety <i>bilobata</i> extract on thiol group content (nmol/ mg protein) in BSA incubated with fructose.	113
5.15	The effects of methanolic <i>Ficus deltoidea</i> variety <i>motleyana</i> extract on thiol group content (nmol/ mg protein) in BSA incubated with fructose.	114
5.16	The effects of <i>Ficus deltoidea</i> varieties extracts and Quercetin on the inhibition of protein carbonyl formation (%) in BSA incubated with fructose.	115
5.17	The effects of <i>Ficus deltoidea</i> variety <i>deltoidea</i> extract and Quercetin on the inhibition of protein carbonyl formation (%) in BSA incubated with fructose.	116
5.18	The effects of <i>Ficus deltoidea</i> variety <i>angustifolia</i> extract on the inhibition of protein carbonyl formation (%) in BSA incubated with fructose.	117
5.19	The effects of <i>Ficus deltoidea</i> varieties extracts and Quercetin on the inhibition of protein carbonyl formation (%) in BSA incubated with fructose.	118
5.20	The effects of <i>Ficus deltoidea</i> variety <i>motleyana</i> extract on the inhibition of protein carbonyl formation (%) in BSA incubated with fructose.	119
5.21	Power reducing assay of methanolic <i>Ficus deltoidea</i> varieties and data was compared with ascorbic acid	122

LIST OF ABBREVIATIONS

AMPK	Adenosine 5' Monophosphate-activated Protein Kinase
AGEs	Advanced Glycation Endproducts
AKT	Protein Kinase B
ALLN	N-acetyl-Leu-Leu-Norleu-al
ANOVA	Analysis of Variance
ATP	Adenosine Triphosphate
cAMP	Cyclic Adenosine Monophosphate
CAP	Catabolite Activator Protein
cfDNA	Cell-free Deoxyribonucleic Acid
DAG	Diacylglycerol
DMEM	Dulbecco's Modified Eagle Medium
DNA	Deoxyribonucleic Acid
DPP-4	Dipeptidyl Peptidase-4
DPPH	2,2-diphenyl-1-picrylhydrazyl
ELISA	Enzyme-Linked Immunosorbent Assay
GDM	Gestational Diabetes Mellitus
GDP	Guanosine diphosphate
GLP-1	Glucagon-like Peptide 1
GTP	Guanosine Triphosphate
IDF	International Diabetes Federation
IFN- γ	Interferon-gamma
IL-6	Interleukin 6
IP3	Inositol Triphosphate
IRS-1	Insulin receptor Substrate 1
JNK	c-Jun N-terminal Kinase

KATP	ATP-sensitive potassium
KRB	Kreb's Ringer Buffer
MAPK	Mitogen-Activated Protein Kinase
mRNA	Messenger Ribonucleic Acid
mTOR	Mechanistic Target of Rapamycin
MTT	3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
NF-kB	Nuclear Factor kappa B
PBS	Phosphate Buffer Saline
PI3K	Phosphatidylinositol 3-kinase
PPAR	Peroxisome proliferator-activated receptor
RAGE	Receptor Advanced Glycation Endproducts
ROS	Reactive Oxygen Species
RPMI 1640	Roswell Park Memorial Institute 1640
SDS	Sodium Dodecyl Sulfate
SGLT-2	Sodium-glucose co-transporter-2
SNP	Single Nucleotide Polymorphism
STAT3	Signal Transducer and Activator of Transcription 3
TLR-4	Toll-Like Receptor 4
TNF- α	Tumor Necrosis Factor Alpha

CHAPTER 1

INTRODUCTION

1.1 Background of Study

Diabetes has occurred worldwide. This metabolic disease has affected all age stages and occurs etiologically in several types. The first type of diabetes mellitus occurs when the beta cell is impaired and ultimately results in the need for insulin injection. The second type of diabetes mellitus is due to insulin resistance, where the insulin-targeting cell does not respond to insulin and is therefore unable to regulate blood glucose levels. Part of that other form of diabetes is the hybrid type in which type 1 diabetes has been detected in adult life preceding type 2 diabetes mellitus. Another form of diabetes is gestational diabetes, which developed during pregnancy due to placental activity. The new estimate in adults with diabetes mellitus showed that large burdens were recorded, particularly in developing countries, where almost 382 million people were affected by metabolic disorder in 2013, and 592 million people were expected to have this disease by 2035 (Guariguata *et. al.*, 2014). In addition, IDF diabetes atlas has shown that the prevalence of this disease is estimated to be 451 million diabetics in 2017 and an additional 242 million in 28 years (Cho *et. al.*, 2018). Metabolic disorder is the top ten cause of death worldwide in the 2013 Global Burden of Disease Study (Naghavi *et. al.*, 2015) while the 2013 Malaysian study reported that diabetes has become one of the leading causes of death in women (Omar *et. al.*, 2019).

Diabetes mellitus has had a negative impact on several aspects. The most common impact is the cost burden where higher costs are needed for catering patients with diabetes mellitus complications in health care (Domeikiene *et. al.*, 2014). It has been estimated that by 2030 the cost of the economic burden of diabetes is more than USD 2.1 trillion compared to 2015, where the cost of the burden is USD 1.3 trillion (Bommer *et. al.*, 2018). IDF diabetes atlas also reported that the average cost per person worldwide in 2014 ranged from USD 1.583 to USD 2.842, while North America and the Caribbean region had the highest annual expenditure per person, while South East Asia had the lowest annual cost per person for diabetes (Da Rocha Fernandes *et. al.*, 2016). Apart from that, looking at the national economic burden in Iran, patients with type 2 diabetes need higher costs compared to other health care costs, while in Malaysia, diabetic patients with anemia require higher costs to be managed (Javanbakht *et. al.*, 2011; Azmi *et. al.*, 2018). In addition, diabetes mellitus has an impact on the economy by decreasing the employment rate of men diagnosed with diabetes (Seuring *et. al.*, 2015). In a study in Kuala Terengganu, unemployment has been a major contributor to atrocious blood glucose levels, apart from social support and belief where diabetes interferes with daily activities (Mat H. *et. al.*, 2019). In addition, metabolic disorder also has a psychological impact on its patients (Jones *et. al.*, 2016). Diabetes patients in Malaysia are two times likely to have been diagnosed with anxiety and depression. Anxiety was associated with age, ethnicity and ischaemic heart disease, while depression is significantly associated with age, ethnicity and household income (Ganasegeran *et. al.*, 2014).

1.2 Statement of Problem

As for now, there is no record of an absolute cure for diabetes mellitus. Metformin was prescribed as a first-line oral drug to reduce hyperglycaemia through its ability to absorb glucose through the AMPK pathway (Zhou *et al.*, 2001). If metformin has not been used to control satisfactory blood glucose levels, a combination of other drugs has been prescribed for diabetic patients to control blood glucose levels (Charbonnel *et al.*, 2006). The combination of drugs is usually effective compared to initial monotherapy (Tosi *et al.*, 2003). For example, when combined with metformin, glimepiride will give better results in blood glucose levels, as this drug has been acting faster to control glycaemic levels (Umpierrez *et al.*, 2006). However, the most common combination of two or more drugs causes various adverse reactions (Bennett *et al.*, 2011). For example, sulphonylureas, an add-on drug to metformin therapy, increases the risk of cardiovascular disease in diabetic patients (Phung *et al.*, 2013). Several drugs have also been reported to have an adverse effect on older patients (Kahn *et al.*, 2006). The drugs included glybenclamide or also known as glyburide and tolbutamide. Glybenclamide in combination with metformin increases the mortality rate compared to drug monotherapy (Fisman *et al.*, 2001). While tolbutamide has led to a deterioration in cardiovascular health (Douros *et al.*, 2017).

In addition, thiazolidinediones have been reported for bone breakage in elderly women (Mabilleau *et al.*, 2010). TZD effect in bone fracture through Erk1/2 and p38 leading to osteocyte apoptosis and activation of the glitazone receptor, leading to the upregulation of sclerostin, an inhibitor of bone formation (Mieczkowska *et al.*, 2012). Pioglitazone, another TZD drug, has increased the incidence of bladder cancer shown in the previous study in French diabetic patients (Neumann *et al.*, 2012). While rosiglitazone increases the risk of myocardial infarction, it causes a higher incidence of death in patients with cardiovascular problems (Nissen and Wolski, 2007). In addition, saxagliptin, which is a DPP-4 inhibitor, increases the risk of hospitalization due to cardiac arrest (Scirica *et al.*, 2013). In addition, a study in Malaysia has shown that diabetic patients in Malaysia have poor adherence to prescribed diabetic and self-care drugs, such as blood glucose testing (Ahmad *et al.*, 2013; Jannoo and Mamode Khan, 2019). Therefore, another alternative treatment for diabetes mellitus is needed to control blood glucose levels without adverse effects.

1.3 Justification of the Study

There are a number of alternative ways to manage diabetes and chronic diseases. Complementary and alternative medicine (CAM) has been another option for the treatment of chronic diseases. CAM can take a number of forms, including prayers, natural products, chiropractic, yoga and meditation (Barnes *et al.*, 2009). The study showed that diabetic patients have a high preference for alternative treatment (Kumar *et al.*, 2006). In fact, the patient who opts for the method still uses ordinary treatment (Donald and Egede, 2006). In the United States, complementary and alternative medicines are chosen because users feel that they have more control over their health

and that the method has improved their health (Rhee *et. al.*, 2018). In addition, many users were satisfied with the impact of the method (Singh *et. al.*, 2004). Apart from that, the methods also include naturopathy and homeopathy. Homeopathy is a therapeutic method that uses natural sources and is believed to improve blood glucose levels (Mourão *et. al.*, 2019). In Malaysia, the use of herbal medicines is favorably accepted and the prevalence is high (Aziz and Tey, 2009). CAM has been reported to be used in patients with chronic diseases in Malaysia (Hasan *et. al.*, 2009). In addition, the use of CAM is high in diabetic patients in Malaysia as a herbal medicine is thought to be safe, of good quality and an additional method for conventional treatment (Hasan *et. al.*, 2011). The use of *Momordica charantia* (bitter melon) and *Orthosiphon stamineus*, *Benth* (Misai Kucing) is most common in diabetic patients in Malaysia (Ching *et. al.*, 2013). *Momordica charantia* has been reported to be able to regulate blood glucose levels and increase glucose disposal in muscle cells (Ahmed *et. al.*, 2004). In addition, herbal medicine capable of overcoming insulin resistance by inhibiting NF- κ B and JNK pathways (Yang *et. al.*, 2015). Natural compounds isolated from *Momordica charantia* have been reported to be capable of inhibiting the activity of 5-007-amylase and 5-007-glucosidase (Shivanagoudra *et. al.*, 2019). The hypoglycaemic effect of *Momordica charantia* is reported due to its cucurbitan constituents and is capable of enhancing the effect of rosiglitazone, a glucose-lowering drug (Harinantenaina *et. al.*, 2006; Nivitashekam *et. al.*, 2009). While *Orthosiphon stamineus*, *Benth* (Misai Kucing) also had a blood glucose-reducing effect (Mohamed *et. al.*, 2011) and was able to inhibit 5-007-amylase and 3-glucosidase (Mohamed *et. al.*, 2012).

Ficus deltoidea is another potential herbal for diabetes mellitus. In Malaysia, the herbal plant is provincially known as Mas Cotek or Serapat Angin and has been used as a post-birth decoction plant. A study in Singapore reported that *Ficus deltoidea* leaves have been used to promote health while its fruit is used for diabetes (Siew *et. al.*, 2014). In addition, it has been reported that the herbal has the ability to reduce pain, to heal wounds, to have an anti-hyperglycemic, anti-inflammatory and anti-melanogenic effect and to reduce blood pressure (Sulaiman *et. al.*, 2008; Abdulla *et. al.*, 2010; Adam *et. al.*, 2011; Oh *et. al.*, 2011; Zakaria *et. al.*, 2012; Kamal *et. al.*, 2019). In addition, the natural compounds responsible for the antioxidant effect of this herbal product are vitexin and isovitexin (Farsi *et. al.*, 2011). In fact, vitexin and isovitexin have been isolated from the leaves of *Ficus deltoidea* and have been reported to have inhibited 5-007-glucosidase (Choo *et. al.*, 2012). Four methane strains of *Ficus deltoidea* have therefore been selected to further elucidate antidiabetic mechanisms in cells.

1.4 Hypothesis of the study

It is assumed that methanolic extracts of *Ficus deltoidea* varieties have antidiabetic activities and mechanisms. Specifically, methanolic *Ficus deltoidea* extracts are capable of enhancing insulin secretion through the K_{ATP} independent pathway, enhancing glucose uptake through the PI3-Kinase pathway, increasing insulin sensitivity in insulin resistance cells and inhibiting the formation of glycated proteins.

1.5 Objectives of the Study

The main objectives of this research are to elucidate antidiabetic mechanisms by using four methanolic extracts of *Ficus deltoidea* varieties. The specific objectives are:

1. To evaluate the insulin-secreting activity of methanolic extracts of *Ficus deltoidea* varieties on β -pancreatic cells and the molecular mechanisms underlying such activity.
2. To determine the potential of extracts of *Ficus deltoidea* varieties to enhance glucose uptake into insulin targeting cells and molecular mechanisms underlying such enhancement.
3. To assess insulin-sensitizing activity of extracts of *Ficus deltoidea* varieties in adipocytes cells.
4. To determine the potential of extracts of *Ficus deltoidea* varieties to prevent diabetes complications by inhibiting the production of Advanced Glycation End Products (AGE).

REFERENCES

- Abdulla MA, Abdul-Aziz Ahmed K, Abu-Luhood FM, Muhanid M. 2010. Role of *Ficus deltoidea* extract in the enhancement of wound healing in experimental rats. *Biomedical Research*. 21(3):241–245.
- Abdullah A, Stoelwinder J, Shortreed S, Wolfe R, Stevenson C, Walls H, De Courten M, Peeters A. 2011. The duration of obesity and the risk of type 2 diabetes. *Public Health Nutrition*. 14(1):119–126.
- Abdullah FI, Chua LS, Rahmat Z, Soontornngun N, Somboon P. 2018. Trypsin Hydrolysed Protein Fractions as Radical Scavengers and Anti-bacterial Agents from *Ficus deltoidea*. *International Journal of Peptide Research and Therapeutics*. 24(2):279–290.
- Abolmaesoomi M, Aziz AA, Junit SM, Ali JM. 2019. Effects of solvent polarity on antioxidant and anti-proliferative activities in breast and colon cancer cells. *European Journal of Integrative Medicine*. 28(October 2018):57–67.
- Abraham NN, Abdul-rahman PS, Aminudin N. 2018. The antioxidant activities, cytotoxic properties, and identification of water-soluble compounds of *Ficus deltoidea* leaves. *PeerJ*:1–20.
- Abu Bakar MF, Ahmad NE, Suleiman M, Rahmat A, Isha A. 2015. *Garcinia dulcis* Fruit Extract Induced Cytotoxicity and Apoptosis in HepG2 Liver Cancer Cell Line. *BioMed Research International*. 2015:1–10.
- Adam Z, Ismail A, Khamis S, Muhammad MH, Hamid M. 2011. Antihyperglycemic activity of *F. deltoidea* ethanolic extract in normal rats. *Sains Malaysiana*. 40(5):489–495.
- Adam Z, Khamis S, Ismail A, Hamid M. 2012. *Ficus deltoidea*: A potential alternative medicine for diabetes mellitus. *Evidence-based Complementary and Alternative Medicine*. 2012(2012):632763.
- Adeyemo AA, Tekola-Ayele F, Doumatey AP, Bentley AR, Chen G, Huang H, Zhou J, Shriner D, Fasanmade O, Okafor G, *et. al.* 2015. Evaluation of genome wide association study associated type 2 diabetes susceptibility loci in sub-Saharan Africans. *Frontiers in Genetics*. 6(335):1–8.
- Adisakwattana S, Thilavech T, Chusak C. 2014. *Mesona Chinensis Benth* extract prevents AGE formation and protein oxidation against fructose-induced protein glycation in vitro. *BMC complementary and alternative medicine*. 14(1):130.
- Aggarwal S, Shailendra G, Ribnicky DM, Burk D, Karki N, Qingxia Wang MS. 2015. An extract of *Artemisia dracuncululus* L. stimulates insulin secretion from β cells, activates AMPK and suppresses inflammation. *Journal of Ethnopharmacology*. 170:98–105.

- Aguirre V, Uchida T, Yenush L, Davis R, White MF. 2000. The c-Jun NH2-terminal kinase promotes insulin resistance during association with insulin receptor substrate-1 and phosphorylation of Ser307. *Journal of Biological Chemistry*. 275(12):9047–9054.
- Ahmad H, Khan I, Wahid A. 2012. Antiglycation and antioxidation properties of *juglans regia* and *calendula officinalis* : possible role in reducing diabetic complications and slowing down ageing. *Journal of Traditional Chinese Medicine*. 32(3):411–414.
- Ahmad NS, Ramli A, Islahudin F, Paraidathathu T. 2013. Medication adherence in patients with type 2 diabetes mellitus treated at primary health clinics in Malaysia. *Patient Preference and Adherence*. 7:525–530.
- Ahmad S, Khan MS, Akhter F, Khan MS aji., Khan A, Ashraf JM, Pandey RP at., Shahab U. 2014. Glycooxidation of biological macromolecules: a critical approach to halt the menace of glycation. *Glycobiology*. 24(11):979–990.
- Ahmed I, Adeghate E, Cummings E, Sharma AK, Singh J. 2004. Beneficial effects and mechanism of action of *Momordica charantia* juice in the treatment of streptozotocin-induced diabetes mellitus in rat. *Molecular and Cellular Biochemistry*. 261(1):63–70.
- Ahmed MF, Kazim SM, Ghori SS, Mehjabeen SS, Ahmed SR, Ali SM, Ibrahim M. 2010. Antidiabetic activity of *vinca rosea* extracts in alloxan-induced diabetic rats. *International Journal of Endocrinology*. 2010(2010):841090.
- Ahmed N. 2005. Advanced glycation endproducts - Role in pathology of diabetic complications. *Diabetes Research and Clinical Practice*. 67(1):3–21.
- Al-Abed Y, Bucala R. 1995. N ϵ -carboxymethyllysine formation by direct addition of glyoxal to lysine during the Maillard reaction. *Bioorganic and Medicinal Chemistry Letters*. 5(18):2161–2162.
- Al-Abed Y, Bucala R. 2000. Structure of a synthetic glucose derived advanced glycation end product that is immunologically cross-reactive with its naturally occurring counterparts. *Bioconjugate Chemistry*. 11(1):39–45.
- Alam MM, Ahmad I, Naseem I. 2015. Inhibitory effect of quercetin in the formation of advance glycation end products of human serum albumin: An in vitro and molecular interaction study. *International Journal of Biological Macromolecules*. 79:336–343.
- Alberto OC, Molina-Carrion M, Abdul-Ghani MA, Folli F, DeFronzo RA, Tripathy D. 2009. Circulating Fibroblast Growth Factor-21 Is Elevated in Impaired Glucose Tolerance and Type 2 Diabetes and Correlates With Cardiovascular and Metabolic Risk. 323(June):1542–1546.
- Al Ali M, El hajj Chehadeh S, Osman W, Almansoori K, Abdulrahman M, Tay G, Alsafar H. 2019. Investigating the association of rs7903146 of TCF7L2 gene, rs5219 of KCNJ11 gene, rs10946398 of CDKAL1 gene, and rs9939609 of FTO gene with type 2 diabetes mellitus in Emirati population. *Meta Gene*. 21(July):100600.

- Alkhalidy H, Moore W, Zhang Y, McMillan R, Wang A, Ali M, Suh KS, Zhen W, Cheng Z, Jia Z, *et. al.* 2015. Small Molecule Kaempferol Promotes Insulin Sensitivity and Preserved Pancreatic β -Cell Mass in Middle-Aged Obese Diabetic Mice. *Journal of Diabetes Research*. 2015(15):138.
- Alvaro C De, Teruel T, Hernandez R, Lorenzo M. 2004. Tumor Necrosis Factor α Produces Insulin Resistance in Skeletal Muscle by Activation of Inhibitor κ B Kinase in a p38 MAPK-dependent Manner. *The Journal of Biological Chemistry*. 279(17):17070–17078.
- An D, Toyoda T, Taylor EB, Yu H, Fujii N, Hirshman MF, Goodyear LJ. 2010. TBC1D1 regulates insulin- and contraction-induced glucose transport in mouse skeletal muscle. *Diabetes*. 59(6):1358–1365.
- Angius F, Floris A. 2015. Liposomes and MTT cell viability assay: An incompatible affair. *Toxicology in Vitro*. 29(2):314–319.
- Anusree SS, Nisha VM, Priyanka A, Raghu KG. 2015. Insulin resistance by TNF- α is associated with mitochondrial dysfunction in 3T3-L1 adipocytes and is ameliorated by puniceic acid, a PPAR γ agonist. *Molecular and Cellular Endocrinology*. 413:120–128.
- Apelqvist J, Elgzyri T, Larsson J, Lndahl M, Nyberg P, Thrne J. 2011. Factors related to outcome of neuroischemic/ischemic foot ulcer in diabetic patients. *Journal of Vascular Surgery*. 53(6):1582-1588.e2.
- Ardestani A, Yazdanparast R. 2007. *Cyperus rotundus* suppresses AGE formation and protein oxidation in a model of fructose-mediated protein glycooxidation. *International Journal of Biological Macromolecules*. 41(5):572–578.
- Arif S, Gibson VB, Nguyen V, Bingley PJ, Todd JA, Guy C, Dunger DB, Dayan CM, Powrie J, Lorenc A, *et. al.* 2017. β -cell specific T-lymphocyte response has a distinct inflammatory phenotype in children with Type 1 diabetes compared with adults. *Diabetic Medicine*. 34(3):419–425.
- Ashraf JM, Shahab U, Tabrez S, Ju E, Choi I, Ahmad S. 2015. International Journal of Biological Macromolecules Quercetin as a finer substitute to aminoguanidine in the inhibition of glycation products. *International Journal of Biological Macromolecules*. 77:188–192.
- Attia RT, Abdel-mottaleb Y, Abdallah DM, El-abhar HS, El-maraghy NN. 2019. Raspberry ketone and *Garcinia Cambogia* rebalanced disrupted insulin resistance and leptin signaling in rats fed high fat fructose diet. *Biomedicine & Pharmacotherapy*. 110(September 2018):500–509.
- Azis NA, Agarwal R, Ismail NM, Ismail NH, Kamal MSA, Radjeni Z, Singh HJ. 2019. Blood pressure lowering effect of *Ficus deltoidea* var. *kunstleri* in spontaneously hypertensive rats: possible involvement of renin–angiotensin–aldosterone system, endothelial function and anti-oxidant system. *Molecular Biology Reports*. 46(3):2841–2849.
- Aziz Z, Tey NP. 2009. Herbal medicines: Prevalence and predictors of use among Malaysian adults. *Complementary Therapies in Medicine*. 17(1):44–50.

- Azmi S, Goh A, Muhammad NA, Tohid H, Rashid MRA. 2018. The Cost and Quality of Life of Malaysian Type 2 Diabetes Mellitus Patients with Chronic Kidney Disease and Anemia. *Value in Health Regional Issues*. 15:42–49.
- Babenko AP, Gonzalez G, Bryan J. 1999. The tolbutamide site of SUR1 and a mechanism for its functional coupling to KATP channel closure. *FEBS Letters*. 459(3):367–376.
- Bae U, Jung E, Jung S, Chae S, Park B. 2018. Mulberry leaf extract displays antidiabetic activity in db/db mice via Akt and AMP-activated protein kinase phosphorylation. *Food & Nutrition Research*. 62:1473.
- Baker JR, Zyzak D V, Thorpe R, Baynes JW. 1994. Chemistry of the Fructosamine Assay: D-Glucosone Is the Product of Oxidation of Amadori Compounds. 40(10):1950–1955.
- Ball AJ, McCluskey JT, Flatt PR, McClenaghan NH. 2004. Chronic exposure to tolbutamide and glibenclamide impairs insulin secretion but not transcription of KATP channel components. *Pharmacological Research*. 50(1):41–46.
- Banerjee S. 2017. Formation of Pentosidine Cross-Linking in Myoglobin by Glyoxal: Detection of Fluorescent Advanced Glycation End Product. *Journal of Fluorescence*. 27(4):1213–1219.
- Barnes PM, Bloom B, Nahin RL. 2009. Complementary and alternative medicine use among adults and children: United States, 2007. *National health statistics reports*.(12):1–23.
- Basu S, McKee M, Galea G, Stuckler D. 2013. Relationship of soft drink consumption to global overweight, obesity, and diabetes: A cross-national analysis of 75 countries. *American Journal of Public Health*. 103(11):2071–2077.
- Beh JE, Khoo LT, Latip J, Abdullah MP, Alitheen NBM, Adam Z, Ismail A, Hamid M. 2013. SDF7, a group of *Scoparia dulcis* Linn. derived flavonoid compounds, stimulates glucose uptake and regulates adipocytokines in 3T3-F442a adipocytes. *Journal of Ethnopharmacology*. 150(1):339–352.
- Beisswenger PJ, Howell SK, O'Dell RM, Wood ME, Touchette AD, Szwegold BS. 2001. alpha -Dicarbonyls Increase in the Postprandial Period and Reflect the Degree of Hyperglycemia. *Diabetes Care*. 24(4):726–732.
- Bennett WL, Maruthur NM, Singh S, Segal JB, Wilson LM, Chatterjee R, Marinopoulos SS, Pahan MA, Ranasinghe P, Block L, *et al.* 2011. Comparative effectiveness and safety of medications for type 2 diabetes: An update including new drugs and 2-drug combinations. *Annals of Internal Medicine*. 154(9):602–618.
- Berdichevsky A, Guarente L, Bose A. 2010. Acute oxidative stress can reverse insulin resistance by inactivation of cytoplasmic JNK. *Journal of Biological Chemistry*. 285(28):21581–21589.

- Berti L, Kellerer M, Capp E, Häring HU. 1997. Leptin stimulates glucose transport and glycogen synthesis in C2C12 myotubes: Evidence for a PI3-kinase mediated effect. *Diabetologia*. 40(5):606–609.
- Bhat M, Zinjarde SS, Bhargava SY, Kumar AR, Joshi BN. 2011. Antidiabetic Indian plants: A good source of potent amylase inhibitors. *Evidence-based Complementary and Alternative Medicine*. 2011(2011):810207.
- Bhavesh VD, Nayak Y, Bs J. 2013. In Vitro Antioxidant and Antiglycation Activity of *Zingiber zerumbet* (Wild Zinger) Rhizome Extract. *International Journal of Research in Pharmaceutical*. 4(4):482–489.
- Bhuiyan MNI, Mitsuhashi S, Sigetomi K, Ubukata M. 2017. Quercetin inhibits advanced glycation end product formation via chelating metal ions, trapping methylglyoxal, and trapping reactive oxygen species. *Bioscience, Biotechnology and Biochemistry*. 81(5):882–890.
- Bidasee KR, Zhang Y, Shao CH, Wang M, Patel KP, Dincer ÜD, Besch HR. 2004. Diabetes Increases Formation of Advanced Glycation End Products on Sarco(endo)plasmic Reticulum Ca²⁺-ATPase. *Diabetes*. 53(2):463–473.
- Bommer C, Sagalova V, Heesemann E, Manne-Goehler J, Atun R, Bärnighausen T, Davies J, Vollmer S. 2018. Global economic burden of diabetes in adults: Projections from 2015 to 2030. *Diabetes Care*. 41(5):963–970.
- Bonora M, Patergnani S, Rimessi A, de Marchi E, Suski JM, Bononi A, Giorgi C, Marchi S, Missiroli S, Poletti F, *et al.* 2012. ATP synthesis and storage. *Purinergic Signalling*. 8(3):343–357.
- Boon J, Hoy AJ, Stark R, Brown RD, Meex RC, Henstridge DC, Schenk S, Meikle PJ, Horowitz JF, Kingwell BA, *et al.* 2013. Ceramides contained in LDL are elevated in type 2 diabetes and promote inflammation and skeletal muscle insulin resistance. *Diabetes*. 62(2):401–410.
- Braun JD, Pastene D o., Breedijk A, Rodriguez A, Hofmann BB, Sticht C, Ochsenstein elke von, Allgayer H, Born J van den, Bakker S, *et al.* 2019. Methylglyoxal down-regulates the expression of cell cycle associated genes and activates the p53 pathway in human umbilical vein endothelial cells. *Scientific Reports*. 9(1152):1–14.
- Bray GA, Hu FB, Willett WC, Popkin BM, Despres J-P, Malik VS. 2010. Sugar-Sweetened Beverages and Risk of Metabolic Syndrome and Type 2 Diabetes: A meta-analysis. *Diabetes Care*. 33(11):2477–2483.
- Brouwers O, de vos-Houben MJM, Niessen PMG, Miyata T, van Nieuwenhoven F, Janssen BJA, Hageman G, Stehouwer CDA, Schalkwijk CG. 2013. Mild oxidative damage in the diabetic rat heart is attenuated by glyoxalase-1 overexpression. *International Journal of Molecular Sciences*. 14(8):15724–15739.
- Bunawan H, Amin NM, Bunawan SN, Baharum SN, Mohd Noor N. 2014. *Ficus deltoidea* jack: A review on its phytochemical and pharmacological importance. *Evidence-based Complementary and Alternative Medicine*. 2014(4):902734.

- Bursal E, Gülçin I. 2011. Polyphenol contents and in vitro antioxidant activities of lyophilised aqueous extract of kiwifruit (*Actinidia deliciosa*). *Food Research International*. 44(5):1482–1489.
- Cai W, Ramdas M, Zhu L, Chen X, Striker GE, Vlassara H. 2012. Oral advanced glycation endproducts (AGEs) promote insulin resistance and diabetes by depleting the antioxidant defenses AGE receptor-1 and sirtuin 1. *Proceedings of the National Academy of Sciences of the United States of America*. 109(39):15888–15893.
- Calabrese V, Cornelius C, Leso V, Trovato-Salinaro A, Ventimiglia B, Cavallaro M, Scuto M, Rizza S, Zanolli L, Neri S, *et. al.* 2012. Oxidative stress, glutathione status, sirtuin and cellular stress response in type 2 diabetes. *Biochimica et Biophysica Acta - Molecular Basis of Disease*. 1822(5):729–736.
- Campbell-Thompson M, Fu A, Kaddis JS, Wasserfall C, Schatz DA, Pugliese A, Atkinson MA. 2016. Insulinitis and β -cell mass in the natural history of type 1 diabetes. *Diabetes*. 65(3):719–731.
- Cao H, Hininger-Favier I, Kelly MA, Benaraba R, Dawson HD, Coves S, Roussel AM, Anderson RA. 2007. Green tea polyphenol extract regulates the expression of genes involved in glucose uptake and insulin signaling in rats fed a high fructose diet. *Journal of Agricultural and Food Chemistry*. 55(15):6372–6378.
- Carmichael J, Degraff WG, Gazdar AF, Minna JD, Mitchell JB. 1987. Evaluation of a Tetrazolium-based Semiautomated Colorimetric Assay : Assessment of Chemosensitivity Testing Evaluation of a Tetrazolium-based Semiautomated Colorimetric Assay : Assessment. *Cancer Research*. 47:936–942.
- Carnagarin R, Dharmarajan AM, Dass CR. 2015. Molecular aspects of glucose homeostasis in skeletal muscle – A focus on the molecular mechanisms of insulin resistance. *Molecular and Cellular Endocrinology*. 417:52–62.
- Cassese A, Esposito I, Fiory F, Barbagallo APM, Paturzo F, Mirra P, Ulianich L, Giacco F, Iadicicco C, Lombardi A, *et. al.* 2008. In skeletal muscle advanced glycation end products (AGEs) inhibit insulin action and induce the formation of multimolecular complexes including the receptor for AGEs. *Journal of Biological Chemistry*. 283(52):36088–36099.
- Chang L, Chiang S-H, Saltiel AR. 2004. Insulin signaling and the regulation of glucose transport. *Molecular medicine (Cambridge, Mass)*. 10(7–12):65–71.
- Charbonnel B, Karasik A, Liu J, Wu M, Meininger G. 2006. Efficacy and safety of the dipeptidyl peptidase-4 inhibitor sitagliptin added to ongoing metformin therapy in patients with type 2 diabetes inadequately controlled with metformin alone. *Diabetes Care*. 29(12):2638–2643.
- Chellan P, Nagaraj RH. 1999. Protein crosslinking by the Maillard reaction: Dicarbonyl-derived imidazolium crosslinks in aging and diabetes. *Archives of Biochemistry and Biophysics*. 368(1):98–104.

- Chen CC, Hsiang CY, Chiang AN, Lo HY, Li CI. 2008. Peroxisome proliferator-activated receptor gamma transactivation-mediated potentiation of glucose uptake by Bai-Hu-Tang. *Journal of Ethnopharmacology*. 118(1):46–50.
- Chen SA, Chen HM, Yao Y Der, Hung CF, Tu CS, Liang YJ. 2012. Topical treatment with anti-oxidants and Au nanoparticles promote healing of diabetic wound through receptor for advance glycation end-products. *European Journal of Pharmaceutical Sciences*. 47(5):875–883.
- Chen SC, Brooks R, Houskeeper J, Bremner SK, Dunlop J, Viollet B, Logan PJ, Salt IP, Ahmed SF, Yarwood SJ. 2017. Metformin suppresses adipogenesis through both AMP-activated protein kinase (AMPK)-dependent and AMPK-independent mechanisms. *Molecular and Cellular Endocrinology*. 440:57–68.
- Chen Y, Carlessi R, Walz N, Cruzat VF, Keane K, John AN, Jiang F-X, Carnagarin R, Dass CR, Newsholme P. 2016. Pigment epithelium-derived factor (PEDF) regulates metabolism and insulin secretion from a clonal rat pancreatic beta cell line BRIN-BD11 and mouse islets. *Molecular and Cellular Endocrinology*. 426:50–60.
- Cheng KKY, Lam KSL, Wang B, Xu A. 2014. Signaling mechanisms underlying the insulin-sensitizing effects of adiponectin. *Best Practice & Research Clinical Endocrinology & Metabolism*. 28(1):3–13.
- Cheng WE, Su LT, Chen SC, Li TC, Lin HW. 2014. Economic burden of diabetes mellitus on patients with respiratory failure requiring mechanical ventilation during hospitalizations. *Value in Health Regional Issues*. 3(1):33–38.
- Cheung N, Wang JJ, Rogers SL, Brancati F, Klein R, Sharrett AR, Wong TY. 2008. Diabetic Retinopathy and Risk of Heart Failure. *Journal of the American College of Cardiology*. 51(16):1573–1578.
- Chiang S, Baumann CA, Kanzaki M, Thurmond DC, Watson RT, Neudauer CL, Macara IG, Pessin JE, Saltiel AR. 2001. Insulin-stimulated GLUT4 translocation requires the CAP-dependent activation of TC10. *Nature*. 410(April):944–948.
- Ching SM, Zakaria ZA, Paimin F, Jalalian M. 2013. Complementary alternative medicine use among patients with type 2 diabetes mellitus in the primary care setting: A cross-sectional study in Malaysia. *BMC Complementary and Alternative Medicine*. 2013(13):148.
- Cho GJ, Hong SC, Oh MJ, Kim HJ. 2013. Vitamin D deficiency in gestational diabetes mellitus and the role of the placenta. *American Journal of Obstetrics and Gynecology*. 209(6):1–8.
- Cho NH, Shaw JE, Karuranga S, Huang Y, da Rocha Fernandes JD, Ohlrogge AW, Malanda B. 2018. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Research and Clinical Practice*. 138:271–281.

- Choi KH, Lee HA, Park MH, Han J. 2017. Cyanidin-3-rutinoside increases glucose uptake by activating the PI3K/Akt pathway in 3T3-L1 adipocytes. *Environmental Toxicology and Pharmacology*. September(54):1-6.
- Chompoo J, Upadhyay A, Kishimoto W, Makise T, Tawata S. 2011. Advanced glycation end products inhibitors from *Alpinia zerumbet* rhizomes. *Food chemistry*. 129(3):709–15.
- Choo CY, Sulong NY, Man F, Wong TW. 2012. Vitexin and isovitexin from the Leaves of *Ficus deltoidea* with in-vivo α -glucosidase inhibition. *Journal of Ethnopharmacology*. 142(2012):776–781.
- Christoffersen CT, Tornqvist H, Vlahos CJ, Bucchini D, Jami J, Meyts P De, Joshi RL. 1998. Insulin and Insulin-like Growth Factor-I Receptor Mediated Differentiation of 3T3-F442A Cells into Adipocytes: Effect of PI 3-Kinase Inhibition. *Biochemical and Biophysical Research Communications*. 430(246):426–430.
- Cokorinos EC, Delmore J, Reyes AR, Albuquerque B, Kjøbsted R, Jørgensen NO, Tran JL, Jatkar A, Cialdea K, Esquejo RM, *et. al.* 2017. Activation of Skeletal Muscle AMPK Promotes Glucose Disposal and Glucose Lowering in Non-human Primates and Mice. *Cell Metabolism*. 25(5):1147-1159.e10.
- Cong L, Chen H, Li Y, Zhou L, McGibbon MA, Taylor SI, Quon MJ. 1997. Physiological Role of Akt in Insulin- Stimulated Translocation of GLUT4 in Transfected Rat Adipose Cells. *Molecular Endocrinology*. 3(13):1881–1890.
- Copps KD, Hancer NJ, Opore-Ado L, Qiu W, Walsh C, White MF. 2010. Irs1 Serine 307 Promotes Insulin Sensitivity in Mice. *Cell Metabolism*. 11(1):84–92.
- Coughlan MT, Yap FYT, Tong DCK, Andrikopoulos S, Gasser A, Thallas-Bonke V, Webster DE, Miyazaki JI, Kay TW, Slattery RM, *et. al.* 2011. Advanced glycation end products are direct modulators of β -cell function. *Diabetes*. 60(10):2523–2532.
- Cousminer DL, Ahlqvist E, Mishra R, Andersen MK, Chesi A, Hawa MI, Davis A, Hodge KM, Bradfield JP, Zhou K, *et. al.* 2018. First genome-wide association study of latent autoimmune diabetes in adults reveals novel insights linking immune and metabolic diabetes. *Diabetes Care*. 41(11):2396–2403.
- Danaei G, Fahimi S, Lu Y, Zhou B, Hajifathalian K, Cesare M Di, Lo WC, Reis-Santos B, Cowan MJ, Shaw JE, *et. al.* 2015. Effects of diabetes definition on global surveillance of diabetes prevalence and diagnosis: A pooled analysis of 96 population-based studies with 331 288 participants. *The Lancet Diabetes and Endocrinology*. 3(8):624–637.
- DeFuria J, Belkina AC, Jagannathan-Bogdan M, Snyder-Cappione J, Carr JD, Nersesova YR, Markham D, Strissel KJ, Watkins AA, Zhu M, *et. al.* 2013. B cells promote inflammation in obesity and type 2 diabetes through regulation of T-cell function and an inflammatory cytokine profile. *Proceedings of the National Academy of Sciences*. 110(13):5133–5138.

- Dekdouk N, Malafronte N, Russo D, Faraone I, De Tommasi N, Ameddah S, Severino L, Milella L. 2015. Phenolic Compounds from *Olea europaea* L. Possess Antioxidant Activity and Inhibit Carbohydrate Metabolizing Enzymes in Vitro. Evidence-based Complementary and Alternative Medicine. 2015:1-9.
- Delong T, Wiles TA, Baker RL, Bradley B, Barbour G, Reisdorph R, Armstrong M, Powell RL, Reisdorph N, Kumar N, *et. al.* 2016. Pathogenic CD4 T cells in type 1 diabetes recognize epitopes formed by peptide fusion. Science. 351(6274):711–714.
- Denizot F, Lang R. 1986. Rapid colorimetric assay for cell growth and survival. Journal of Immunological Methods. 89(2):271–277.
- Detimary P, Gilon P, Nenquin M, Henquin J-C. 1994. Two sites of glucose control of insulin release with distinct dependence on the energy state in pancreatic B-cells. Biochem J. 297:455–461.
- Dhar I, Dhar A, Wu L, Desai KM. 2014. Methylglyoxal, a reactive glucose metabolite, increases renin angiotensin aldosterone and blood pressure in male sprague-dawley rats. American Journal of Hypertension. 27(3):308–316.
- Domeikiene A, Vaivadaite J, Ivanauskiene R, Padaiga Z. 2014. Direct cost of patients with type 2 diabetes mellitus healthcare and its complications in Lithuania. Medicina (Lithuania). 50(1):54–60.
- Dominguez C, Ruiz E, Gussinye M, Carrascosa A. 1998. Oxidative stress at onset and in early stages of type 1 diabetes in children and adolescents. Diabetes Care. 21(10):1736–1742.
- Donald G, Egede LE. 2006. Association Between Complementary and Alternative Medicine Use , Preventive Care Practices , and Use of Conventional Medical. Diabetes Care. 29(1):15–19.
- van Dooren FEP, Nefs G, Schram MT, Verhey FRJ, Denollet J, Pouwer F. 2013. Depression and Risk of Mortality in People with Diabetes Mellitus: A Systematic Review and Meta-Analysis. PLoS ONE. 8(3):e57058.
- Dou H, Ma E, Yin L, Jin Y, Wang H. 2013. The Association between Gene Polymorphism of TCF7L2 and Type 2 Diabetes in Chinese Han Population: A Meta-Analysis. PLoS ONE. 8(3):1–13.
- Douros A, Yin H, Yu OHY, Filion KB, Azoulay L, Suissa S. 2017. Pharmacologic differences of sulfonylureas and the risk of adverse cardiovascular and hypoglycemic events. Diabetes Care. 40(11):1506–1513.
- Duan W, Jiang KCZ, Chen X, Sun L, Li J, Lei J, Xu Q, Ma J, Li X, Han L, *et. al.* 2017. Desmoplasia suppression by metformin-mediated AMPK activation inhibits pancreatic cancer progression. Cancer Letters. 385:225–233.
- Dzolin S, Ahmad R, Zain MM, Ismail MI. 2015. Flavonoid distribution in four varieties of *Ficus deltoidea* (Jack). Journal of Medicinal Plant and Herbal Therapy Research. 2015(3):1–9.

- Eastham LL, Mills CN, Niles RM. 2008. PPAR α/γ expression and activity in mouse and human melanocytes and melanoma cells. *Pharmaceutical Research*. 25(6):1327–1333.
- Elchebly M, Payette P, Michaliszyn E, Cromlish W, Collins S, Loy AL, Normandin D, Cheng A, Himms-hagen J, Chan C, *et. al.* 1999. Increased Insulin Sensitivity and Obesity Resistance in Mice Lacking the Protein Tyrosine. *Science*. 283(March):1544–1548.
- Elham Farsi, Armaghan Shafaei, Sook Yee Hor, Mohamed B. Khadeer Ahamed, Mun Fei Yam, Idress Hamad Attitalla MZA and ZI, Farsi. 2011. Correlation between enzymes inhibitory effects and antioxidant activities of standardized fractions of methanolic extract obtained from *Ficus deltoidea* leaves. *African Journal of Biotechnology*. 10(67):15184–15194.
- Ellman GL. 1959. Tissue sulfhydryl groups. *Archives of Biochemistry and Biophysics*. 82(1):70–77.
- Erkoç Kaya D, Arikoğlu H, Kaya SA, Öztürk O, Gönen MS. 2017. Transcription factor 7-like 2 (TCF7L2) gene polymorphisms are strong predictors of type 2 diabetes among nonobese diabetics in the Turkish population. *Turkish Journal of Medical Sciences*. 47(1):22–28.
- Fan J, Johnson MH, Lila MA, Yousef G, De Mejia EG. 2013. Berry and citrus phenolic compounds inhibit dipeptidyl peptidase IV: Implications in diabetes management. *Evidence-based Complementary and Alternative Medicine*. 2013(2013):479505.
- Fang WL, Huang KH, Lan YT, Lin CH, Chang SC, Chen MH, Chao Y, Lin WC, Lo SS, Li AFY, *et. al.* 2016. Mutations in PI3K/AKT pathway genes and amplifications of PIK3CA are associated with patterns of recurrence in gastric cancers. *Oncotarget*. 7(5):6201–6220.
- Farsi E, Shafaei A, Hor SY, Ahamed MBK, Yam MF, Asmawi MZ, Ismail Z. 2013. Genotoxicity and acute and subchronic toxicity studies of a standardized methanolic extract of *Ficus deltoidea* leaves. *Clinics (São Paulo, Brazil)*. 68(6):865–75.
- Farsi E, Shafaei A, Hor SY, Ahamed MBK, Yam MF, Hamad Attitalla I, Ismail Z, Asmawi MZ. 2011. Correlation between enzymes inhibitory effects and antioxidant activities of standardized fractions of methanolic extract obtained from *Ficus deltoidea* leaves. *African Journal of Biotechnology*. 10(67):15184–15194.
- Fatihah HNN, Mat N, Zaimah ARN, Zuhailah MN, Norhaslinda H, Khairil M, Ghani AY, Ali AM. 2012. Morphological phylogenetic analysis of seven varieties of *Ficus deltoidea* Jack from the Malay Peninsula of Malaysia. *PloS one*. 7(12):e52441.
- Fauzi, Widodo H. 2019. Short Communication: Aphrodisiac plants used by Dayak Ethnic in Central Kalimantan Province, Indonesia. *Biodiversitas Journal of Biological Diversity*. 20(7):1859–1865.

- Feig DS, Shah BR, Lipscombe LL, Wu CF, Ray JG, Lowe J, Hwee J, Booth GL. 2013. Preeclampsia as a Risk Factor for Diabetes: A Population-Based Cohort Study. *PLoS Medicine*. 10(4):1–8.
- Ferdaoussi M, Fu J, Dai X, Fox JEM, Suzuki K, Smith N, Plummer G, Macdonald PE. 2017. SUMOylation and calcium control syntaxin-1A and secretagogin sequestration by tomosyn to regulate insulin exocytosis in human β cells. *Scientific Reports*. 248(7):1–9.
- Fisman EZ, Tenenbaum A, Adler Y, Motro M, Boyko V, Benderly M, Friedensohn A, Kohanovski M, Rotzak R, Schneider H, *et. al.* 2001. Oral antidiabetic treatment in patients with coronary disease: Time-related increased mortality on combined glyburide/metformin therapy over a 7.7-year follow-up. *Clinical Cardiology*. 24(2):151–158.
- Foda AA, Aal IAA. 2012. Metformin as a new therapy for endometriosis, its effects on both clinical picture and cytokines profile. *Middle East Fertility Society Journal*. 17(4):262–267.
- Fort PE, Losiewicz MK, Reiter CEN, Singh RSJ, Nakamura M, Abcouwer SF, Barber AJ, Gardner TW. 2011. Differential roles of hyperglycemia and hypoinsulinemia in diabetes induced retinal cell death: Evidence for retinal insulin resistance. *PLoS ONE*. 6(10):e26498.
- Foukas LC, Berenjano IM, Gray A, Khwaja A, Vanhaesebroeck B. 2010. Activity of any class IA PI3K isoform can sustain cell proliferation and survival. *Proceedings of the National Academy of Sciences of the United States of America*. 107(25):11381–11386.
- Fryer LGD, Parbu-Patel A, Carling D. 2002. The anti-diabetic drugs rosiglitazone and metformin stimulate AMP-activated protein kinase through distinct signaling pathways. *Journal of Biological Chemistry*. 277(28):25226–25232.
- Gall WE, Beebe K, Lawton KA, Adam KP, Mitchell MW, Nakhle PJ, Ryals JA, Milburn M V., Nannipieri M, Camastra S, *et. al.* 2010. α -Hydroxybutyrate Is an Early Biomarker of Insulin Resistance and Glucose Intolerance in a Nondiabetic Population. *PLoS ONE*. 5(5):e10883.
- Ganasegeran K, Renganathan P, Manaf RA, Al-Dubai SAR. 2014. Factors associated with anxiety and depression among type 2 diabetes outpatients in Malaysia: A descriptive cross-sectional single-centre study. *BMJ Open*. 4(4):1–7.
- Gao J, Zhong X, Ding Y, Bai T, Wang H, Wu H, Liu Y, Yang J, Zhang Y. 2016. Inhibition of voltage-gated potassium channels mediates uncarboxylated osteocalcin-regulated insulin secretion in rat pancreatic β cells. *European Journal of Pharmacology*. 777:41–48.
- Garcia-Barrado M-J, Ravier AM, Rolland J-F, Gilon P, Nenquin M, Henquin J. 2001. Inhibition of Protein Synthesis Sequentially Impairs Distinct Steps of Stimulus-secretion Coupling in Pancreatic beta Cells. *Endocrinology*. 142(1):299–307.

- García-Vicente S, Yraola F, Marti L, González-Muñoz E, García-Barrado MJ, Cantó C, Abella A, Bour S, Artuch R, Sierra C, *et. al.* 2007. Oral insulin-mimetic compounds that act independently of insulin. *Diabetes*. 56(2):486–493.
- Gasic S, Tian B, Green A. 1999. Tumor Necrosis Factor α Stimulates Lipolysis in Adipocytes by Decreasing G i Protein Concentrations. *The Journal of biological chemistry*. 274(10):6770–6775.
- German JP, Wisse BE, Thaler JP, Oh-I S, Sarruf DA, Ogimoto K, Kaiyala KJ, Fischer JD, Matsen ME, Taborsky GJ, *et. al.* 2010. Leptin deficiency causes insulin resistance induced by uncontrolled diabetes. *Diabetes*. 59(7):1626–1634.
- Gheni G, Ogura M, Iwasaki M, Yokoi N, Minami K, Nakayama Y, Harada K, Hastoy B, Wu X, Takahashi H, *et. al.* 2014. Glutamate Acts as a Key Signal Linking Glucose Metabolism to Incretin / cAMP Action to Amplify Insulin Article Glutamate Acts as a Key Signal Linking Glucose Metabolism to Incretin / cAMP Action to Amplify Insulin Secretion. *Cell Reports*. 9(2):661–673.
- Ghosh A, Kushwaha HR, Hasan MR, Pareek A, Sopory SK, Singla-Pareek SL. 2016. Presence of unique glyoxalase III proteins in plants indicates the existence of shorter route for methylglyoxal detoxification. *Scientific Reports*. 6(January):1–15.
- Ghosh S, Ahire M, Patil S, Jabgunde A, Bhat Dusane M, Joshi BN, Pardesi K, Jachak S, Dhavale DD, Chopade BA. 2012. Antidiabetic activity of *gnidia glauca* and *dioscorea bulbifera*: Potent amylase and glucosidase inhibitors. *Evidence-based Complementary and Alternative Medicine*. 2012(2012):929051.
- Gomis A, Valdeolmillos M. 1998. Regulation by tolbutamide and diazoxide of the electrical activity in mouse pancreatic β -cells recorded *in vivo*. *British Journal of Pharmacology*. 123(3):443–448.
- Goryakin Y, Monsivais P, Suhrcke M. 2017. Soft drink prices, sales, body mass index and diabetes: Evidence from a panel of low-, middle- and high-income countries. *Food Policy*. 73(September):88–94.
- Graham DJ, Ouellet-Hellstrom R, MaCurdy TE, Ali F, Sholley C, Worrall C, Kelman JA. 2010. Risk of Acute Myocardial Infarction, Stroke, Heart Failure, and Death in Elderly Medicare Patients Treated With Rosiglitazone or Pioglitazone. *Jama*. 304(4):411.
- Guariguata L, Whiting DR, Hambleton I, Beagley J, Linnenkamp U, Shaw JE. 2014. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes Research and Clinical Practice*. 103(2):137–149.
- Gui J, Liu Q, Feng L. 2013. Metformin vs Insulin in the Management of Gestational Diabetes: A Meta-Analysis. *PLoS ONE*. 8(5):e64585.
- Guillausseau P-J, Meas T, Virally M, Laloi-Michelin M, Médeau V, Kevorkian J-P. 2008. Abnormalities in insulin secretion in type 2 diabetes mellitus. *Diabetes & metabolism*. 34 Suppl 2:S43-8.

- Gupte A, Mora S. 2006. Activation of the Cbl insulin signaling pathway in cardiac muscle; Dysregulation in obesity and diabetes. *Biochemical and Biophysical Research Communications*. 342(3):751–757.
- Hajar S, Giribabu N, Mazuin N, Bakar A, Salleh N. 2017. *Marantodes pumilum* (Kacip fatimah) enhances in-vitro glucose uptake in 3T3-L1 adipocyte cells and reduces pancreatic complications in streptozotocin-nicotinamide induced male diabetic rats. *Biomedicine Pharmacotherapy*. 96(July):716–726.
- Hamada Y, Araki N, Koh N, Nakamura J, Horiuchi S, Hotta N. 1996. Rapid Formation of Advanced Glycation End Products by Intermediate Metabolites of Glycolytic Pathway and Polyol Pathway. *Biochemical and Biophysical Research Communications*. 543:539–543.
- Hammes H, Martin S, Frederlin K, Geisen K, Brownlee M. 2006. Aminoguanidine Treatment Inhibits the Development of Experimental Diabetic Retinopathy. *Proceedings of the National Academy of Sciences*. 89(19):9364d – 9364.
- Han MS, Jung DY, Morel C, Lakhani SA, Kim JK, Flavell RA, Davis RJ. 2012. JNK Expression by Macrophages Promotes Obesity-Induced Insulin Resistance and Inflammation. *Science*. 339(6116):218–222.
- Han Y, Zhu G, Han L, Hou F, Huang W, Liu H, Gan J, Jiang T, Li X, Wang W, *et. al.* 2014. Short-term rosuvastatin therapy for prevention of contrast-induced acute kidney injury in patients with diabetes and chronic kidney disease. *Journal of the American College of Cardiology*. 63(1):62–70.
- Hanafi MMM, Afzan A, Yaakob H, Aziz R, Sarmidi MR, Wolfender JL, Prieto JM. 2017. In vitro pro-apoptotic and anti-migratory effects of *Ficus deltoidea* L. plant extracts on the human prostate cancer cell lines PC3. *Frontiers in Pharmacology*. 8(DEC):1–20.
- Hanani Soib H, Ware I, Yaakob H, Mukrish H, Roji Sarmidi M. 2015. Antioxidant and anti-cancer activity of standardized extracts of three varieties of *Ficus deltoidea*'s leaves. *Jurnal Teknologi*. 77(3):19–25.
- Hannan JMA, Marenah L, Ali L, Rokeya B, Flatt PR, Abdel-Wahab YHA. 2006. *Ocimum sanctum* leaf extracts stimulate insulin secretion from perfused pancreas, isolated islets and clonal pancreatic beta-cells. *Journal of Endocrinology*. 189(1):127–136.
- Hanssen NMJ, Beulens JWJ, Van Dieren S, Scheijen JLJM, Van Der A DL, Spijkerman AMW, Van Der Schouw YT, Stehouwer CDA, Schalkwijk CG. 2015. Plasma advanced glycation end products are associated with incident cardiovascular events in individuals with type 2 diabetes: A Case-Cohort study with a median follow-Up of 10 years (EPIC-NL). *Diabetes*. 64(1):257–265.
- Hanssen NMJ, Engelen L, Ferreira I, Scheijen JLJM, Huijberts MS, Van Greevenbroek MMJ, Van Der Kallen CJH, Dekker JM, Nijpels G, Stehouwer CDA, *et. al.* 2013. Plasma levels of advanced glycation endproducts Nε-(carboxymethyl) lysine, Nε-(carboxyethyl)lysine, and pentosidine are not independently associated with cardiovascular disease in individuals with or without type 2 diabetes: The hoorn and CODAM studies. *Journal of Clinical Endocrinology*

and Metabolism. 98(8):1369–1373.

- Hanssen NMJ, Wouters K, Huijberts MS, Gijbels MJ, Sluimer JC, Scheijen JLJM, Heeneman S, Biessen EAL, Daemen MJAP, Brownlee M, *et al.* 2014. Higher levels of advanced glycation endproducts in human carotid atherosclerotic plaques are associated with a rupture-prone phenotype. *European Heart Journal*. 35(17):1137–1146.
- Hardie DG, Carling D, Carlson M. 1998. THE AMP-ACTIVATED/SNF1 PROTEIN KINASE SUBFAMILY: Metabolic Sensors of the Eukaryotic Cell? *Annual Review of Biochemistry*. 67(1):821–855.
- Harinantenaina L, Tanaka M, Takaoka S, Oda M, Mogami O, Uchida M, Asakawa Y. 2006. Momordica charantia constituents and antidiabetic screening of the isolated major compounds. *Chemical and Pharmaceutical Bulletin*. 54(7):1017–1021.
- Harrison AL, Shields N, Taylor NF, Frawley HC. 2016. Exercise improves glycaemic control in women diagnosed with gestational diabetes mellitus: a systematic review. *Journal of Physiotherapy*. 62(4):188–196.
- Hasan SS, Ahmed SI, Bukhari NI, Loon WCW. 2009. Use of complementary and alternative medicine among patients with chronic diseases at outpatient clinics. *Complementary Therapies in Clinical Practice*. 15(3):152–157.
- Hasan SS, Loon WC, Ahmadi K, Ahmed SI, Bukhari NI. 2011. Reasons, perceived efficacy and factors associated with complementary and alternative medicine use among Malaysian patients with diabetes mellitus. *British Journal of Diabetes and Vascular Disease*. 11(2):92–98.
- Haselgrübler R, Stadlbauer V, Stübl F, Schwarzinger B, Rudzionyte I, Himmelsbach M, Iken M, Weghuber J. 2018. Insulin Mimetic Properties of Extracts Prepared from *Bellis perennis*. *Molecules*. 23(10):2605.
- Hasham R, Choi HK, Sarmidi MR, Park CS. 2013. Protective effects of a *Ficus deltoidea* (Mas cotek) extract against UVB-induced photoageing in skin cells. *Biotechnology and Bioprocess Engineering*. 18(1):185–193.
- Hassan RH, De Sousa ACP, Mahfouz R, Hainault I, Blachnio-Zabielska A, Bourron O, Koskas F, Górski J, Ferré P, Fougelle F, *et al.* 2016. Sustained action of ceramide on the insulin signaling pathway in muscle cells: Implication of the double-stranded RNA-activated protein kinase. *Journal of Biological Chemistry*. 291(6):3019–3029.
- Hatlapatka K, Matz M, Schumacher K, Baumann K, Rustenbeck I. 2011. Bidirectional insulin granule turnover in the submembrane space during K⁺ depolarization-induced secretion. *Traffic*. 12(9):1166–1178.
- Hawley SA, Pan DA, Mustard KJ, Ross L, Bain J, Edelman AM, Frenguelli BG, Hardie DG. 2005. Calmodulin-dependent protein kinase kinase- β is an alternative upstream kinase for AMP-activated protein kinase. *Cell Metabolism*. 2(1):9–19.

- Heinonen S, Buzkova J, Muniandy M, Kaksonen R, Ollikainen M, Ismail K, Hakkarainen A, Lundbom J, Lundbom N, Vuolteenaho K, *et. al.* 2015. Impaired mitochondrial biogenesis in adipose tissue in acquired obesity. *Diabetes*. 64(9):3135–3145.
- Hering BJ, Clarke WR, Bridges ND, Eggerman TL, Alejandro R, Bellin MD, Chaloner K, Czarniecki CW, Goldstein JS, Hunsicker LG, *et. al.* 2016. Phase 3 trial of transplantation of human islets in type 1 diabetes complicated by severe hypoglycemia. *Diabetes Care*. 39(7):1230–1240.
- Hirschey MD, Shimazu T, Jing E, Grueter CA, Collins AM, Aouizerat B, Stančáková A, Goetzman E, Lam MM, Schwer B, *et. al.* 2011. SIRT3 deficiency and mitochondrial protein hyperacetylation accelerate the development of the metabolic syndrome. *Molecular Cell*. 44(2):177–190.
- Hoehn KL, Hohnen-Behrens C, Cederberg A, Wu LE, Turner N, Yuasa T, Eбина Y, James DE. 2008. IRS1-Independent Defects Define Major Nodes of Insulin Resistance. *Cell Metabolism*. 7(5):421–433.
- Hoo RLC, Wong JYL, Qiao CF, Xu A, Xu HX, Lam KSL. 2010. The effective fraction isolated from *Radix Astragali* alleviates glucose intolerance, insulin resistance and hypertriglyceridemia in db/db diabetic mice through its anti-inflammatory activity. *Nutrition and Metabolism*. 7:1–12.
- Hotamisligil GS, Arner P, Caro JF, Atkinson RL, Spiegelman BM. 1995. Increased adipose tissue expression of tumor necrosis factor- α in human obesity and insulin resistance. *Journal of Clinical Investigation*. 95(5):2409–2415.
- Hotamisligil GS, David ML, Choy LN, Spiegelman BM. 1994. Tumor necrosis factor α inhibits signaling from the insulin receptor. *Biochemistry*. 91(May):4854–4858.
- Hu J, Nomura Y, Bashir A, Fernandez-Hernandez H, Itzkowitz S, Pei Z, Stone J, Loudon H, Peter I. 2013. Diversified Microbiota of Meconium Is Affected by Maternal Diabetes Status. *PLoS ONE*. 8(11):e78257.
- Huang THW, Peng G, Kota BP, Li GQ, Yamahara J, Roufogalis BD, Li Y. 2005. Anti-diabetic action of *Punica granatum* flower extract: Activation of PPAR- γ and identification of an active component. *Toxicology and Applied Pharmacology*. 207(2):160–169.
- Hurst Von PR, Stonehouse W, Coad J. 2010. Vitamin D supplementation reduces insulin resistance in South Asian women living in New Zealand who are insulin-resistant and vitamin D deficient – a randomised, placebo-controlled trial. *British Journal of Nutrition*. 103(04):549–555.
- Ilyanie Y, Wong TW, Choo CY. 2011. Evaluation of hypoglycemic activity and toxicity profiles of the leaves of *Ficus deltoidea* in rodents. *Journal of Complementary and Integrative Medicine*. 8(1):1–16.
- Irshad Z, Xue M, Ashour A, Larkin JR, Thornalley PJ, Rabbani N. 2019. Activation of the unfolded protein response in high glucose treated endothelial cells is mediated by methylglyoxal. *Scientific Reports*. 9(1):1–14.

- Ito H, Abe M, Mifune M, Oshikiri K, Antoku S, Takeuchi Y, Togane M. 2011. Hyperuricemia Is Independently Associated with Coronary Heart Disease and Renal Dysfunction in Patients with Type 2 Diabetes Mellitus. *PLoS ONE*. 6(11):1–7.
- Jaganjac M, Almuraikhy S, Al-Khelaifi F, Al-Jaber M, Bashah M, Mazloun NA, Zarkovic K, Zarkovic N, Waeg G, Kafienah W, *et. al.* 2017. Combined metformin and insulin treatment reverses metabolically impaired omental adipogenesis and accumulation of 4-hydroxynonenal in obese diabetic patients. *Redox Biology*. 12(February):483–490.
- Jainandusing S, Rita Koole H, van Miert JNI, Rietveld T, Darcos Wattimena JL, Sijbrands EJG, de Rooij FWM. 2018. Transcription factor 7-like 2 gene links increased in vivo insulin synthesis to type 2 diabetes. *EBioMedicine*.30(2017):295-302
- Jaiswal D, Rai PK, Mehta S, Chatterji S, Shukla S, Rai DK, Sharma G, Sharma B, Khair S, Watal G. 2013. Role of *Moringa oleifera* in regulation of diabetes-induced oxidative stress. *Asian Pacific Journal of Tropical Medicine*. 6(6):426–432.
- Jannoo Z, Mamode Khan N. 2019. Medication Adherence and Diabetes Self-Care Activities among Patients with Type 2 Diabetes Mellitus. *Value in Health Regional Issues*. 18:30–35.
- Jannoo Z, Wah YB, Lazim AM, Hassali MA. 2017. Examining diabetes distress, medication adherence, diabetes self-care activities, diabetes-specific quality of life and health-related quality of life among type 2 diabetes mellitus patients. *Journal of Clinical and Translational Endocrinology*. 9:48–54.
- Javanbakht M, Baradaran HR, Mashayekhi A, Haghdoost AA, Khamseh ME, Kharazmi E, Sadeghi A. 2011. Cost-of-illness analysis of type 2 diabetes mellitus in Iran. *PLoS ONE*. 6(10):1–7.
- Jha AK, Huang SCC, Sergushichev A, Lampropoulou V, Ivanova Y, Loginicheva E, Chmielewski K, Stewart KM, Ashall J, Everts B, *et. al.* 2015. Network integration of parallel metabolic and transcriptional data reveals metabolic modules that regulate macrophage polarization. *Immunity*. 42(3):419–430.
- Jheng H-F, Tsai P-J, Guo S-M, Kuo L-H, Chang C-S, Su I-J, Chang C-R, Tsai Y-S. 2012. Mitochondrial Fission Contributes to Mitochondrial Dysfunction and Insulin Resistance in Skeletal Muscle. *Molecular and Cellular Biology*. 32(2):309–319.
- Jin D, Sun J, Huang J, He Y, Yu A, Yu X, Yang Z. 2014. Cytokine TNF- α reduces g0s2 expression and stimulates lipolysis through PPAR-c inhibition in 3T3-L1 adipocytes. *Cytokine*. 69:196–205.
- Jo J, Young Choi M, Koh D. 2009. Beneficial effects of intercellular interactions between pancreatic islet cells in blood glucose regulation. *Journal of Theoretical Biology*. 257(2009):312–319.
- Jones A, Olsen MZ, Perrild HJD, Willaing I. 2016. The psychological impact of living with diabetes: Descriptive findings from the DAWN2 study in Denmark. *Primary Care Diabetes*. 10(1):83–86.

- Jörns A, Wedekind D, Jähne J, Lenzen S. 2020. Pancreas Pathology of Latent Autoimmune Diabetes in Adults (LADA) in Patients and in a LADA Rat Model Compared to Type 1 Diabetes Mellitus. *Diabetes*. 69(April):624–633.
- Juneja A, Dwivedi S, Srivastava DK, Chandra K. 2017. Insulin Resistance in Young Obese Subjects and Its Relation to Smoking (A Pilot Study). *Indian Journal of Clinical Biochemistry*. 32(1):99–102.
- Kahn SE, Haffner SM, Heise MA, Herman WH, Holman RR, Jones NP, Kravitz BG, Lachin JM, O'Neill MC, Zinman B, et al. 2006. Glycemic durability of rosiglitazone, metformin, or glyburide monotherapy. *New England Journal of Medicine*. 355(23):2427–2443.
- Kamal MSA, Ismail NH, Satar NA, Azis NA, Radjeni Z, Mohammad Noor HS, Kasim N, Singh H. 2019. Standardized ethanol-water extract of *Ficus deltoidea Angustifolia* reduces blood pressure in spontaneously hypertensive rats. *Clinical and Experimental Hypertension*. 41(5):444–451.
- Kane S, Sano H, Liu SCH, Asara JM, Lane WS, Garner CC, Lienhard GE. 2002. A method to identify serine kinase substrates. Akt phosphorylates a novel adipocyte protein with a Rab GTPase-activating protein (GAP) domain. *Journal of Biological Chemistry*. 277(25):22115–22118.
- Kaneb HM, Sharp PS, Rahmani-Kondori N, Wells DJ. 2011. Metformin treatment has no beneficial effect in a dose-response survival study in the SOD1 G93A mouse model of ALS and is harmful in female mice. *PLoS ONE*. 6(9):e24189.
- Kaneko K, Ueki K, Takahashi N, Hashimoto S, Okamoto M, Awazawa M, Okazaki Y, Ohsugi M, Inabe K, Umehara T, et al. 2010. Class IA phosphatidylinositol 3-kinase in pancreatic β cells controls insulin secretion by multiple mechanisms. *Cell Metabolism*. 12(6):619–632.
- Kang C, Lee H, Jung ES, Seyedian R, Jo M, Kim J, Kim JS, Kim E. 2012. Saffron (*Crocus sativus L.*) increases glucose uptake and insulin sensitivity in muscle cells via multipathway mechanisms. *Food Chemistry*. 135(4):2350–2358.
- Karges B, Schwandt A, Heidtmann B, Kordonouri O, Binder E, Schierloh U, Boettcher C, Kapellen T, Rosenbauer J, Holl RW. 2017. Association of insulin pump therapy vs insulin injection therapy with severe hypoglycemia, ketoacidosis, and glycemic control among children, adolescents, and young adults with type 1 diabetes. *JAMA - Journal of the American Medical Association*. 318(14):1358–1366.
- Kim M-B, Kim C, Song Y, Hwang J-K. 2014. Antihyperglycemic and Anti-Inflammatory Effects of Standardized *Curcuma xanthorrhiza* Roxb. Extract and Its Active Compound Xanthorrhizol in High-Fat Diet-Induced Obese Mice. *Evidence-Based Complementary and Alternative Medicine*. 2014:1–10.
- Kim TH, Choi SE, Ha ES, Jung JG, Han SJ, Kim HJ, Kim DJ, Kang Y, Lee KW. 2013. IL-6 induction of TLR-4 gene expression via STAT3 has an effect on insulin resistance in human skeletal muscle. *Acta Diabetologica*. 50(2):189–200.

- Kindt ASD, Fuerst RW, Knoop J, Laimighofer M, Telleps T, Hippich M, Woerheide MA, Wahl S, Wilson R, Sedlmeier EM, *et. al.* 2018. Allele-specific methylation of type 1 diabetes susceptibility genes. *Journal of Autoimmunity*. 89:63–74.
- Kirkiz S, Yarali N, Arman Bilir O, Tunc B. 2014. Metformin-induced hemolytic anemia. *Medical Principles and Practice*. 23(2):183–185.
- Ko B-S, Choi SB, Park SK, Jang JS, Kim YE, Park S. 2005. Insulin sensitizing and insulinotropic action of berberine from *Cortidis Rhizoma*. *Biological and Pharmaceutical Bulletin*. 28(8):1431–1437.
- Koffert JP, Mikkola K, Virtanen KA, Andersson A-MD, Faxius L, Hällsten K, Heglind M, Guiducci L, Pham T, Silvola JMU, *et. al.* 2017. Metformin treatment significantly enhances intestinal glucose uptake in patients with type 2 diabetes: Results from a randomized clinical trial. *Diabetes Research and Clinical Practice*. 131:208–216.
- Koivisto UM, Martinez-Valdez H, Bilan PJ, Burdett E, Ramlal T, Klip A. 1991. Differential regulation of the GLUT-1 and GLUT-4 glucose transport systems by glucose and insulin in L6 muscle cells in culture. *Journal of Biological Chemistry*. 266(4):2615–2621.
- Koleva II, Van Beek TA, Linssen JPH, De Groot A, Evstatieva LN. 2002. Screening of plant extracts for antioxidant activity: A comparative study on three testing methods. *Phytochemical Analysis*. 13(1):8–17.
- Komatsu M, Sato Y, Yamada S, Yamauchi K, Hashizume K, Aizawa T. 2002. Triggering of insulin release by a combination of cAMP signal and nutrients: an ATP-sensitive K⁺ channel-independent phenomenon. *Diabetes*. 51(Supplement 1):S29-32.
- König M, Bulik S, Holzhütter HG. 2012. Quantifying the contribution of the liver to glucose homeostasis: A detailed kinetic model of human hepatic glucose metabolism. *PLoS Computational Biology*. 8(6):1–17.
- Krolopp JE, Thornton SM, Abbott MJ. 2016. IL-15 activates the Jak3/STAT3 signaling pathway to mediate glucose uptake in skeletal muscle cells. *Frontiers in Physiology*. 7(DEC):1–10.
- Kubota T, Kubota N, Kumagai H, Yamaguchi S, Kozono H, Takahashi T, Inoue M, Itoh S, Takamoto I, Sasako T, *et. al.* 2009. Impaired Insulin Signaling in Endothelial Cells Reduces Insulin-Induced Glucose Uptake by Skeletal Muscle. *Cell Metabolism*. 13(3):294–307.
- Kumar D, Bajaj S, Mehrotra R. 2006. Knowledge, attitude and practice of complementary and alternative medicines for diabetes. *Public Health*. 120(8):705–711.
- Kuri-harcuch W, Green H. 1978. Adipose conversion of 3T3 cells depends 'on a serum factor. *Proceedings of the National Academy of Sciences*. 75(12):6107–6109.
- Lakshmi BS, Sujatha S, Anand S, Sangeetha KN, Narayanan RB, Katiyar C, Kanaujia A, Duggar R, Singh Y, Srinivas K, *et. al.* 2009. Cinnamic acid, from the bark

- of *Cinnamomum cassia*, regulates glucose transport via activation of GLUT4 on L6 myotubes in a phosphatidylinositol 3-kinase-independent manner. *Journal of Diabetes*. 1(2):99–106.
- Lee-Young RS, Hoffman NJ, Murphy KT, Henstridge DC, Samocha-Bonet D, Siebel AL, Iliades P, Zivanovic B, Hong YH, Colgan TD, *et. al.* 2016. Glucose-6-phosphate dehydrogenase contributes to the regulation of glucose uptake in skeletal muscle. *Molecular Metabolism*. 5(11):1083–1091.
- Lee JO, Lee SK, Kim JH, Kim N, You GY, Moon JW, Kim SJ, Park SH, Kim HS. 2012. Metformin regulates glucose transporter 4 (GLUT4) translocation through AMP-activated protein kinase (AMPK)-mediated Cbl/CAP signaling in 3T3-L1 preadipocyte cells. *Journal of Biological Chemistry*. 287(53):44121–44129.
- Lee JY, Song J, Kwon K, Jang S, Kim C, Baek K, Kim J, Park C. 2012. Human DJ-1 and its homologs are novel glyoxalases. *Human Molecular Genetics*. 21(14):3215–3225.
- Lennon GP, Bettini M, Burton AR, Vincent E, Arnold PY, Santamaria P, Vignali DAA. 2009. T Cell Islet Accumulation in Type 1 Diabetes Is a Tightly Regulated, Cell-Autonomous Event. *Immunity*. 31(4):643–653.
- Levin P, Zhou S, Durden E, Farr AM, Gill J, Wei W. 2016. Clinical and Economic Outcomes Associated with the Timing of Initiation of Basal Insulin in Patients with Type 2 Diabetes Mellitus Previously Treated with Oral Antidiabetes Drugs. *Clinical Therapeutics*. 38(1):110–121.
- Levine RL, Garland D, Oliver CN, Amici A, Climent I, Lenz A-G, Ahn B, Shaltiel S, Stadtman ER. 1990. Determination of Carbonyl Content in Oxidatively Modified Proteins. *Methods in Enzymology*. 186:464–478.
- Li XH, Du LL, Cheng XS, Jiang X, Zhang Y, Lv BL, Liu R, Wang JZ, Zhou XW. 2013. Glycation exacerbates the neuronal toxicity of β -amyloid. *Cell Death and Disease*. 4(6):1–9.
- Li Y, Xu S, Mihaylova MM, Zheng B, Hou X, Jiang B, Park O, Luo Z, Lefai E, Shyy JYJ, *et. al.* 2011. AMPK phosphorylates and inhibits SREBP activity to attenuate hepatic steatosis and atherosclerosis in diet-induced insulin-resistant mice. *Cell Metabolism*. 13(4):376–388.
- Liang Z, Li N, Qi H, Wan L, Cai P, Xu Z, Li B. 2016. Formation of peptide bound pyrrolidine in the maillard model systems with different lys-containing dipeptides and tripeptides. *Molecules*. 21(4):1–21.
- Liggins J, Furth AJ. 1997. Role of protein-bound carbonyl groups in the formation of advanced glycation endproducts. *Biochimica et Biophysica Acta - Molecular Basis of Disease*. 1361(2):123–130.
- Van Der Linden MW, Plat AW, Erkens JA, Emneus M, Herings RMC. 2009. Large impact of antidiabetic drug treatment and hospitalizations on economic burden of diabetes mellitus in the Netherlands during 2000 to 2004. *Value in Health*. 12(6):909–914.

- Linné Y, Barkeling B, Rössner S. 2002. Natural course of gestational diabetes mellitus: Long term follow up of women in the SPAWN study. *BJOG: An International Journal of Obstetrics and Gynaecology*. 109(11):1227–1231.
- Liu E, Meigs JB, Pittas AG, McKeown NM, Economos CD, Booth SL, Jacques PF. 2008. Plasma 25-Hydroxyvitamin D Is Associated with Markers of the Insulin-resistant Phenotype in Nondiabetic Adults. *Journal of Nutrition*. 139(2):329–334.
- Liu L, Zhang X, Rong C, Rui C, Ji H, Qian YJ, Jia R, Sun L. 2014. Distinct DNA methylomes of human placentas between pre-eclampsia and gestational diabetes mellitus. *Cellular Physiology and Biochemistry*. 34(6):1877–1889.
- Liu Q, Liu S, Gao L, Sun S, Huan Y, Li C, Wang Y, Guo N, Shen Z. 2017. Anti-diabetic effects and mechanisms of action of a Chinese herbal medicine preparation JQ-R in vitro and in diabetic KKAY mice. *Acta Pharmaceutica Sinica B*. 7(4):461–469.
- Longnus SL, Ségalen C, Giudicelli J, Sajan MP, Farese R V., Van Obberghen E. 2005. Insulin signalling downstream of protein kinase B is potentiated by 5'AMP-activated protein kinase in rat hearts in vivo. *Diabetologia*. 48(12):2591–2601.
- Van Der Lugt T, Weseler AR, Gebbink WA, Vrolijk MF, Opperhuizen A, Bast A. 2018. Dietary advanced glycation endproducts induce an inflammatory response in human macrophages in vitro. *Nutrients*. 10(12):1–10.
- Luo S, Lin J, Xie Z, Xiang Y, Zheng P, Huang G, Li X, Liao Y, Hagopian WA, Wang CY, *et. al.* 2016. HLA genetic discrepancy between latent autoimmune diabetes in adults and type 1 diabetes: LADA China study No. 6. *Journal of Clinical Endocrinology and Metabolism*. 101(4):1693–1700.
- Luoto R, Kinnunen TI, Aittasalo M, Kolu P, Raitanen J, Ojala K, Mansikkamäki K, Lamberg S, Vasankari T, Komulainen T, *et. al.* 2011. Primary Prevention of Gestational Diabetes Mellitus and Large-for-Gestational-Age Newborns by Lifestyle Counseling: A Cluster-Randomized Vontrolled Trial. *PLoS Medicine*. 8(5):e1001036.
- Lyons TJ, Thorpe SR, Jenkins AJ, Baynes JW, Fu M-X, Requena JR. 2002. The Advanced Glycation End Product, N -(Carboxymethyl)lysine, Is a Product of both Lipid Peroxidation and Glycooxidation Reactions. *Journal of Biological Chemistry*. 271(17):9982–9986.
- Ma RCW, So WY, Yang X, Yu LWL, Kong APS, Ko GTC, Chow CC, Cockram CS, Chan JCN, Tong PCY. 2008. Erectile Dysfunction Predicts Coronary Heart Disease in Type 2 Diabetes. *Journal of the American College of Cardiology*. 51(21):2045–2050.
- Mabilleau G, Mieczkowska A, Edmonds ME. 2010. Thiazolidinediones induce osteocyte apoptosis and increase sclerostin expression. *Diabetic medicine*. 27(8):925–932.

- El Maghraby HA, Nafee T, Guiziry D, Elnashar A. 2015. Randomized controlled trial of the effects of metformin versus combined oral contraceptives in adolescent PCOS women through a 24 month follow up period. *Middle East Fertility Society Journal*. 20(3):131–137.
- Mahmoud MF, Hassan NA, El Bassossy HM, Fahmy A. 2013. Quercetin Protects against Diabetes-Induced Exaggerated Vasoconstriction in Rats: Effect on Low Grade Inflammation. *PLoS ONE*. 8(5):e63784.
- Marques CMS, Nunes EA, Lago L, Pedron CN, Manieri TM, Sato RH, Oliveira VX, Cerchiaro G. 2017. Generation of Advanced Glycation End-Products (AGEs) by glycooxidation mediated by copper and ROS in a human serum albumin (HSA) model peptide: reaction mechanism and damage in motor neuron cells. *Mutation Research - Genetic Toxicology and Environmental Mutagenesis*. 824(October):42–51.
- Martin SS, Haruta T, Morris AJ, Klippel A, Williams LT, Olefsky JM. 1996. Activated Phosphatidylinositol 3-Kinase Is Sufficient to Mediate Actin Rearrangement and GLUT4 Translocation in 3T3-L1 Adipocytes. *The Journal of biological chemistry*. 271(July 26):17605–17609.
- Marupuru S, Senapati P, Pathadka S, Miraj SS, Unnikrishnan MK, Manu MK. 2017. Protective effect of metformin against tuberculosis infections in diabetic patients: an observational study of south Indian tertiary healthcare facility. *The Brazilian Journal of Infectious Diseases*. 21(3):312–316.
- Masharani UB, Maddux BA, Li X, Sakkas GK, Mulligan K, Schambelan M, Goldfine ID, Youngren JF. 2011. Insulin resistance in non-obese subjects is associated with activation of the JNK pathway and impaired insulin signaling in skeletal muscle. *PLoS ONE*. 6(5):1–9.
- Maskarinec G, Jacobs S, Amshoff Y, Setiawan VW, Shvetsov YB, Franke AA, Kolonel LN, Haiman CA, Le Marchand L. 2018. Sleep duration and incidence of type 2 diabetes: the Multiethnic Cohort. *Sleep Health*. 4(1):27–32.
- Masterjohn C, Park Y, Lee J, Noh SK, Koo SI, Bruno RS. 2013. Dietary fructose feeding increases adipose methylglyoxal accumulation in rats in association with low expression and activity of glyoxalase-2. *Nutrients*. 5(8):3311–3328.
- Mat H. NS, Mohd Y. SS, Kadir A. A, Noor M. N, Draman N, Muhammad J. 2019. Depression, anxiety, stress and socio-demographic factors for poor glycaemic control in patients with type II diabetes. *Journal of Taibah University Medical Sciences*. 14(3):268–276.
- Mat N, Rosni NA, Ab Rashid NZ, Haron N, Mohd Nor Z, Hasan Nudin NF, Yunus AG, Ali AM. 2012. Leaf morphological variations and heterophylly in *Ficus deltoidea* Jack (Moraceae). *Sains Malaysiana*. 41(5):527–538.
- Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. 1985. Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 28(7):412–419.

- McVicar CM, Ward M, Colhoun LM, Guduric-Fuchs J, Bierhaus A, Fleming T, Schlotterer A, Kolibabka M, Hammes HP, Chen M, *et. al.* 2015. Role of the receptor for advanced glycation endproducts (RAGE) in retinal vasodegenerative pathology during diabetes in mice. *Diabetologia*. 58(5):1129–1137.
- Mieczkowska A, Baslé MF, Chappard D, Mabileau G. 2012. Thiazolidinediones induce osteocyte apoptosis by a G protein-coupled receptor 40-dependent mechanism. *Journal of Biological Chemistry*. 287(28):23517–23526.
- Misu H, Takamura T, Takayama H, Hayashi H, Matsuzawa-Nagata N, Kurita S, Ishikura K, Ando H, Takeshita Y, Ota T, *et. al.* 2010. A liver-derived secretory protein, selenoprotein P, causes insulin resistance. *Cell Metabolism*. 12(5):483–495.
- Mitchell P, Bandello F, Schmidt-Erfurth U, Lang GE, Massin P, Schlingemann RO, Sutter F, Simader C, Burian G, Gerstner O, *et. al.* 2011. The RESTORE study: Ranibizumab monotherapy or combined with laser versus laser monotherapy for diabetic macular edema. *Ophthalmology*. 118(4):615–625.
- Mitra P, Zheng X, Czech MP. 2004. RNAi-based analysis of CAP, Cbl, and CrkII function in the regulation of GLUT4 by insulin. *Journal of Biological Chemistry*. 279(36):37431–37435.
- Mohamed EAH, Mohamed AJ, Asmawi MZ, Sadikun A, Ebrika OS, Yam MF. 2011. Antihyperglycemic effect of *orthosiphon stamineus benth* leaves extract and its bioassay-guided fractions. *Molecules*. 16(5):3787–3801.
- Mohamed EAH, Siddiqui MJA, Ang LF, Sadikun A, Chan SH, Tan SC, Asmawi MZ, Yam MF. 2012. Potent α -glucosidase and α -amylase inhibitory activities of standardized 50% ethanolic extracts and sinensetin from *Orthosiphon stamineus Benth* as anti-diabetic mechanism. *BMC Complementary and Alternative Medicine*. 2012(12):176.
- Mohammad Noor HS, Ismail NH, Kasim N, Zohdi RM, Ali AM. 2016. Hypoglycemic and glucose tolerance activity of standardized extracts *Ficus deltoidea* varieties in normal rats. *Journal of Medicinal Plants Studies*. 4(5):275–279.
- Monnier VM, Sun W, Gao X, Sell DR, Cleary PA, Lachin JM, Genuth S. 2015. Skin collagen advanced glycation endproducts (AGEs) and the long-term progression of sub-clinical cardiovascular disease in type 1 diabetes. *Cardiovascular Diabetology*. 14(1):1–9.
- Morgenstern J, Fleming T, Schumacher D, Eckstein V, Freichel M, Herzig S, Nawroth P. 2017. Loss of glyoxalase 1 induces compensatory mechanism to achieve dicarbonyl detoxification in mammalian Schwann cells. *Journal of Biological Chemistry*. 292(8):3224–3238.
- Morris AP, Voight BF, Teslovich TM, Ferreira T, Segrè A V., Steinthorsdottir V, Strawbridge RJ, Khan H, Grallert H, Mahajan A, *et. al.* 2012. Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nature Genetics*. 44(9):981–990.

- Mosmann T. 1983. Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*. 65(1–2):55–63.
- Mourão L, Carillo R, Linares SM, Canabarro A, Fischer RG. 2019. Homeopathy and periodontal treatment in type ii diabetic patients: A 1-year randomized clinical trial. *Brazilian Dental Journal*. 30(2):139–145.
- Mu J, Brozinick JT, Valladares O, Bucan M, Birnbaum MJ. 2001. A Role for AMP-Activated Protein Kinase in Contraction- and Hypoxia-Regulated Glucose Transport in Skeletal Muscle. 7:1085–1094.
- Mukhtar HM, Singh A, Kaur H. 2018. Bioassay guided fractionation and in vitro anti-plasmodial activity of *Ficus deltoidea* and *ficus benjamina*. *Pharmacognosy Journal*. 10(2):235–240.
- Naghavi M, Wang H, Lozano R, Davis A, Liang X, Zhou M, Vollset SE, Abbasoglu Ozgoren A, Abdalla S, Abd-Allah F, *et. al.* 2015. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: A systematic analysis for the Global Burden of Disease Study 2013. *The Lancet*. 385(9963):117–171.
- Nasri H, Rafieian-Kopaei M. 2014. Metformin: Current knowledge. *Journal of research in medical sciences : the official journal of Isfahan University of Medical Sciences*. 19(7):658–64.
- Nasser MW, Wani NA, Ahirwar DK, Powell CA, Ravi J, Elbaz M, Zhao H, Padilla L, Zhang X, Shilo K, *et. al.* 2015. RAGE mediates S100A7-induced breast cancer growth and metastasis by modulating the tumor microenvironment. *Cancer Research*. 75(6):974–985.
- Nemet I, Strauch CM, Monnier VM. 2011. Favored and disfavored pathways of protein crosslinking by glucose: Glucose lysine dimer (GLUCOLD) and crossline versus glucosepane. *Amino Acids*. 40(1):167–181.
- Neumann A, Weill A, Ricordeau P, Fagot JP, Alla F, Allemand H. 2012. Pioglitazone and risk of bladder cancer among diabetic patients in France: A population-based cohort study. *Diabetologia*. 55(7):1953–1962.
- Nguyen-Tu MS, Da Silva Xavier G, Leclerc I, Rutter GA. 2018. Transcription factor-7-like 2 (TCF7L2) gene acts downstream of the Lkb1/Stk11 kinase to control mTOR signaling, cell growth, and insulin secretion. *Journal of Biological Chemistry*. 293(36):14178–14189.
- Nishimoto S, Fukuda D, Higashikuni Y, Tanaka K, Hirata Y, Murata C, Kim-Kaneyama JR, Sato F, Bando M, Yagi S, *et. al.* 2016. Obesity-induced DNA released from adipocytes stimulates chronic adipose tissue inflammation and insulin resistance. *Science Advances*. 2(3):1–12.
- Nishimura S, Manabe I, Nagasaki M, Eto K, Yamashita H, Ohsugi M, Otsu M, Hara K, Ueki K, Sugiura S, *et. al.* 2009. CD8+ effector T cells contribute to macrophage recruitment and adipose tissue inflammation in obesity. *Nature Medicine*. 15(8):914–920.

- Nissen SE, Wolski K. 2007. Effect of Rosiglitazone on the Risk of Myocardial Infarction and Death from Cardiovascular Causes. *Journal of Medicine (Cincinnati)*. 356(24):2457–2471.
- Nissen SE, Wolski K. 2007. Effect of Rosiglitazone on the Risk of Myocardial Infarction and Death from Cardiovascular Causes. *The New England Journal of Medicine*. 356(24):2457–2471.
- Niu Y, Li S, Na L, Feng R, Liu L, Li Y, Sun C. 2012. Mangiferin decreases plasma free fatty acids through promoting its catabolism in liver by activation of AMPK. *PLoS ONE*. 7(1):1–8.
- Nivitashekam SN, Asad M, Prasad VS. 2009. Pharmacodynamic interaction of *Momordica charantia* with rosiglitazone in rats. *Chemico-Biological Interactions*. 177(3):247–253.
- Noipha K, Ratanachaiyavong S, Ninla-aesong P. 2010. Enhancement of glucose transport by selected plant foods in muscle cell line L6. *Diabetes Research and Clinical Practice*. 89(2):e22–e26.
- Nur Fatimah HN, Nashriyah M, Nor Zaimah AR, Khairil M, Ali AM. 2014. Leaf morphology and anatomy of 7 varieties of *Ficus deltoidea* (Moraceae). *Turkish Journal of Botany*. 38:677–685.
- O'Neill HM. 2013. AMPK and exercise: Glucose uptake and insulin sensitivity. *Diabetes and Metabolism Journal*. 37(1):1–21.
- Oh MJ, Abdul Hamid M, Ngadiran S, Seo YK, Sarmidi MR, Park CS. 2011. *Ficus deltoidea* (Mas cotek) extract exerted anti-melanogenic activity by preventing tyrosinase activity in vitro and by suppressing tyrosinase gene expression in B16F1 melanoma cells. *Archives of Dermatological Research*. 303(3):161–170.
- Oikarinen M, Tauriainen S, Oikarinen S, Honkanen T, Collin P, Rantala I, Mäki M, Kaukinen K, Hyöty H. 2012. Type 1 diabetes is associated with enterovirus infection in gut mucosa. *Diabetes*. 61(3):687–691.
- Ojo O, Ojo CC. 2015. Insulinotropic actions of *Moringa oleifera* involves the induction of membrane depolarization and enhancement of intracellular calcium concentration. *Journal of Experimental and Integrative Medicine*. 5(1):36–41.
- Omar A, Ganapathy SS, Anuar MFM, Khoo YY, Jeevananthan C, Maria Awaluddin S, Yn JLM, Rao C. 2019. Cause-specific mortality estimates for Malaysia in 2013: Results from a national sample verification study using medical record review and verbal autopsy. *BMC Public Health*. 19(1):1–12.
- Omar MH, Mullen W, Crozier A. 2011. Identification of proanthocyanidin dimers and trimers, flavone C-glycosides, and antioxidants in *Ficus deltoidea*, a Malaysian herbal tea. *Journal of Agricultural and Food Chemistry*. 59(4):1363–1369.
- Ouellet V, Labbe SM, Blondin DP, Phoenix S, Guerin B, Hama F, Turcotte EE, Richard D, Carpentier AC. 2012. Brown adipose tissue oxidative metabolism contributes to energy expenditure during acute cold exposure in humans. *The Journal of Clinical Investigation*. 122(2):545–552.

- Oyaizu M. 1986. Studies on products of browning reaction. Antioxidative activities of products of browning reaction prepared from glucosamine. The Japanese Journal of Nutrition and Dietetics. 44(6):307–315.
- Padiya R, Chowdhury D, Borkar R, Srinivas R, Pal Bhadra M, Banerjee SK. 2014. Garlic attenuates cardiac oxidative stress via activation of PI3K/AKT/Nrf2-Keap1 pathway in fructose-fed diabetic rat. PLoS ONE. 9(5):3–10.
- Park JE, Han JS. 2018. A *Portulaca oleracea* L. extract promotes insulin secretion via a K⁺ATP channel dependent pathway in INS-1 pancreatic β -cells. Nutrition Research and Practice. 12(3):183–190.
- Parsons WJ, Ramkumar V, Stiles GL. 1988. Isobutylmethylxanthine stimulates adenylate cyclase by blocking the inhibitory regulatory protein, G(i). Molecular Pharmacology. 34(1):37–41.
- Patterson ME, Yee JK, Wahjudi P, Mao CS, Lee W-NP. 2018. Acute metabolic responses to high fructose corn syrup ingestion in adolescents with overweight/obesity and diabetes. Journal of Nutrition & Intermediary Metabolism. 14(August):1–7.
- Pedersen BK, Febbraio MA. 2012. Muscles, exercise and obesity: Skeletal muscle as a secretory organ. Nature Reviews Endocrinology. 8(8):457–465.
- Peng W, Chen JQ, Liu C, Malu S, Creasy C, Tetzlaff MT, Xu C, McKenzie JA, Zhang C, Liang X, *et. al.* 2016. Loss of PTEN promotes resistance to T cell-mediated immunotherapy. Cancer Discovery. 6(2):202–216.
- Pérez-Ramírez IF, González-Dávalos ML, Mora O, Gallegos-Corona MA, Reynoso-Camacho R. 2017. Effect of *Ocimum sanctum* and *Crataegus pubescens* aqueous extracts on obesity, inflammation, and glucose metabolism. Journal of Functional Foods. 35:24–31.
- Perez C, Fernandez-Galaz C, Teresa Fernandez-Agullo CA, Andres A, Ros M, Carrascosa JM. 2004. Leptin impairs insulin signalling in isolated rat adipocytes. Diabetologia. 53(19):347–353.
- Petersen KF, Oral EA, Dufour S, Befroy D, Ariyan C, Yu C, Cline GW, Depaoli AM, Taylor SI, Gorden P, *et. al.* 2002. Leptin reverses insulin resistance and hepatic steatosis in patients with severe lipodystrophy. Clinical Investigation. 109(10):1345–1350.
- Phillips SA, Thornalley PJ. 1993. The formation of methylglyoxal from triose phosphates: Investigation using a specific assay for methylglyoxal. European Journal of Biochemistry. 212(1):101–105.
- Phung OJ, Schwartzman E, Allen RW, Engel SS, Rajpathak SN. 2013. Sulphonylureas and risk of cardiovascular disease: Systematic review and meta-analysis. Diabetic Medicine. 30(10):1160–1171.

- Pinent M, Blay M, Bladé MC, Salvadó MJ, Arola L, Ardévol A. 2004. Grape seed-derived procyanidins have an antihyperglycemic effect in streptozotocin-induced diabetic rats and insulinomimetic activity in insulin-sensitive cell lines. *Endocrinology*. 145(11):4985–4990.
- Poudel A, Savari O, Striegel DA, Periwal V, Taxy J, Millis JM, Witkowski P, Atkinson MA, Hara M. 2015. Beta-cell destruction and preservation in childhood and adult onset type 1 diabetes. *Endocrine*. 49(3):693–702.
- Pourahmadi M, Erfanian S, Moradzadeh M, Jahromi AS. 2015. Non-association between rs7903146 and rs12255372 polymorphisms in transcription factor 7-like 2 gene and type 2 diabetes mellitus in Jahrom city, Iran. *Diabetes and Metabolism Journal*. 39(6):512–517.
- Qiu YH, Deng FY, Li MJ, Lei SF. 2014. Identification of novel risk genes associated with type 1 diabetes mellitus using a genome-wide gene-based association analysis. *Journal of Diabetes Investigation*. 5(6):649–656.
- Quaye EA, Amporful EO, Akweongo P, Aikins MK. 2015. Analysis of the Financial Cost of Diabetes Mellitus in Four Cocoa Clinics of Ghana. *Value in Health Regional Issues*. 7:49–53.
- Quesada I, Nadal A, Soria B. 1999. Different effects of tolbutamide and diazoxide in α , and β -cells within intact islets of Langerhans. *Diabetes*. 48(12):2390–2397.
- Rajamani U, Jialal I. 2014. Hyperglycemia Induces Toll-Like Receptor-2 and -4 Expression and Activity in Human Microvascular Retinal Endothelial Cells: Implications for Diabetic Retinopathy. 2014:7–10.
- Rajaraman B, Ramadas N, Krishnasamy S, Ravi V, Pathak A, Devasena CS, Swaminathan K, Ganeshprasad A, Kuppuswamy AA, Vedantham S. 2019. Hyperglycaemia cause vascular inflammation through advanced glycation end products/early growth response-1 axis in gestational diabetes mellitus. *Molecular and Cellular Biochemistry*. 456(1-2):179-190.
- Rakyan VK, Beyan H, Down TA, Hawa MI, Maslau S, Aden D, Daunay A, Busato F, Mein CA, Manfras B, *et. al.* 2011. Identification of type 1 Diabetes-associated DNA methylation variable positions that precede disease diagnosis. *PLoS Genetics*. 7(9):1–9.
- Rao Sirasanagandla S, Ranganath Pai Karkala S, Potu BK, Bhat KMR. 2014. Beneficial effect of *Cissus quadrangularis linn.* on osteopenia associated with streptozotocin-induced type 1 diabetes mellitus in male Wistar rats. *Advances in Pharmacological Sciences*. 2014(2014):483051.
- Raposeiras-Roubín S, Rodiño-Janeiro BK, Paradela-Dobarro B, Grigorian-Shamagian L, García-Acuña JM, Aguiar-Souto P, Jacquet-Hervet M, Reino-Maceiras M V., González-Juanatey JR, Álvarez E. 2013. Fluorescent advanced glycation end products and their soluble receptor: The Birth of New Plasmatic Biomarkers for Risk Stratification of Acute Coronary Syndrome. *PLoS ONE*. 8(9):e74302.
- Ren JM, Marshall BA, Gulve EA, Gao J, Johnson DW, Holloszy JO, Mueckler M. 1993. Evidence from transgenic mice that glucose transport is rate-limiting for

- glycogen deposition and glycolysis in skeletal muscle. *Journal of Biological Chemistry*. 268(22):16113–16115.
- Rezania A, Bruin JE, Arora P, Rubin A, Batushansky I, Asadi A, O'Dwyer S, Quiskamp N, Mojibian M, Albrecht T, *et al.* 2014. Reversal of diabetes with insulin-producing cells derived in vitro from human pluripotent stem cells. *Nature Biotechnology*. 32(11):1121–1133.
- Rhee TG, Westberg SM, Harris IM. 2018. Complementary and Alternative Medicine in U.S. Adults with Diabetes: Reasons for Use and Perceived Benefits. *Journal of Diabetes*. 10(4):310–319.
- da Rocha Fernandes J, Ogurtsova K, Linnenkamp U, Guariguata L, Seuring T, Zhang P, Cavan D, Makaroff LE. 2016. IDF Diabetes Atlas estimates of 2014 global health expenditures on diabetes. *Diabetes Research and Clinical Practice*. 117:48–54.
- Röder P V, Wu B, Liu Y, Weiping Han. 2016. Pancreatic regulation of glucose homeostasis. *Experimental and Molecular Medicine*. 48(3):e219.
- Rosen ED, Spiegelman BM. 2006. Adipocytes as regulators of energy balance and glucose homeostasis. *Nature*. 444(7121):847–853.
- Rossi J, Balthasar N, Olson D, Scott M, Berglund E, Lee CE, Choi MJ, Lauzon D, Lowell BB, Elmquist JK. 2011. Melanocortin-4 receptors expressed by cholinergic neurons regulate energy balance and glucose homeostasis. *Cell Metabolism*. 13(2):195–204.
- Rui L. 2014. Energy metabolism in the liver. *Comprehensive Physiology*. 4(1):177–197.
- Sakai M, Oimomi M, Kasuga M. 2002. Experimental studies on the role of fructose in the development of diabetic complications. *Kobe Journal of Medical Sciences*. 48(5–6):125–136.
- Sakamoto K, Holman GD. 2008. Emerging role for AS160/TBC1D4 and TBC1D1 in the regulation of GLUT4 traffic. *American Journal of Physiology - Endocrinology and Metabolism*. 295(1):29–37.
- Samsulrizal N, Awang Z, MohdNajib MLH, Idzham M, Zarin A. 2011. Effect of *Ficus deltoidea* leaves extracts on sperm quality, LDH-C4 activity and testosterone level in alloxan-induced male diabetic rats. 2011 IEEE Colloquium on Humanities, Science and Engineering, CHUSER 2011.(Chuser):888–891.
- Samsulrizal N, Goh Yong Meng, Ahmad H, Dom SM, Syimal'ain Azmi N, Noor Mohamad Zin NS, Ebrahimi M. 2017. Changes in pancreatic histology, insulin secretion and oxidative status in diabetic rats following treatment with *Ficus deltoidea* and vitexin. *BMC complementary and alternative medicine*. 17(1):290.
- Samsulrizal N, Goh Y. M., Hafandi A, Dom SM, Nur Syimal'ain A, Noor Syaffinaz NM, Ebrahimi M. 2017. Improvement of spatial learning and memory, cortical gyrification patterns and brain oxidative stress markers in diabetic rats treated

- with *Ficus deltoidea* leaf extract and vitexin. *Journal of Traditional and Complementary Medicine*.8(1):190-202.
- Sanders MJ, Grondin PO, Hegarty BD, Snowden MA, Carling D. 2007. Investigating the mechanism for AMP activation of the AMP-activated protein kinase cascade. *Biochemical Journal*. 403(1):139–148.
- Sandholm N, Salem RM, McKnight AJ, Brennan EP, Forsblom C, Isakova T, McKay GJ, Williams WW, Sadlier DM, Mäkinen VP, *et al.* 2012. New Susceptibility Loci Associated with Kidney Disease in Type 1 Diabetes. *PLoS Genetics*. 8(9):1–13.
- Sandouk T, Reda D, Hofmann C. 1993. Antidiabetic agent pioglitazone enhances adipocyte differentiation of 3T3-F442A cells. *American Journal of Physiology-Cell Physiology*. 264(6):C1600–C1608.
- Sano H, Eguez L, Teruel MN, Fukuda M, Chuang TD, Chavez JA, Lienhard GE, McGraw TE. 2007. Rab10, a Target of the AS160 Rab GAP, Is Required for Insulin-Stimulated Translocation of GLUT4 to the Adipocyte Plasma Membrane. *Cell Metabolism*. 5(4):293–303.
- Sano H, Kane S, Sano E, Miinea CP, Asara JM, Lane WS, Garner CW., Lienhard GE. 2003. Insulin-stimulated Phosphorylation of a Rab GTPase-activating Protein Regulates GLUT4 Translocation. *The Journal of Biological Chemistry*. 278(April 25):14599–14602.
- Scheijen LJ, Hanssen NMJ, van Greevenbroek MM, Van der Kallen CJ, Feskens EJM, Stehouwer CDA, Schalkwijk CG. 2018. Dietary intake of advanced glycation endproducts is associated with higher levels of advanced glycation endproducts in plasma and urine: The CODAM study. *Clinical Nutrition*. 37(3):919–925.
- Schmidt S, Jakab M, Jav S, Streif D, Pitschmann A, Zehl M, Purevsuren S, Glasl S, Ritter M. 2013. Extracts from *Leonurus sibiricus* L. increase insulin secretion and proliferation of rat INS-1E insulinoma cells. *Journal of Ethnopharmacology*. 150(1):85–94.
- Schommers P, Thurau A, Bultmann-Mellin I, Guschlbauer M, Klatt AR, Rozman J, Klingenspor M, de Angelis MH, Alber J, Gründemann D, *et al.* 2017. Metformin causes a futile intestinal–hepatic cycle which increases energy expenditure and slows down development of a type 2 diabetes-like state. *Molecular Metabolism*. 6(7):737–747.
- Schwartz A V., Sellmeyer DE, Vittinghoff E, Palermo L, Lecka-Czernik B, Feingold KR, Strotmeyer ES, Resnick HE, Carbone L, Beamer BA, *et al.* 2006. Thiazolidinedione use and bone loss in older diabetic adults. *Journal of Clinical Endocrinology and Metabolism*. 91(9):3349–3354.
- Schwartz J, Foster CM, Satin MS. 1985. Growth hormone and insulin-like growth factors I and II produce distinct alterations in glucose metabolism in 3T3-F442A adipocytes. *Proceedings of the National Academy of Sciences*. 82(December):8724–8728.

- Scirica BM, Bhatt DL, Braunwald E, Steg G, Davidson J, Hirshberg B, Ohman P, Frederick R, Wiviott SD, Hoffman EB, *et. al.* 2013. Saxagliptin and Cardiovascular Outcomes in Patients with Type 2 Diabetes Mellitus. *The New England Journal of Medicine.* 369(14):1–9.
- Sebastián D, Hernández-Alvarez MI, Segalés J, Sorianello E, Muñoz JP, Sala D, Waget A, Liesa M, Paz JC, Gopalacharyulu P, *et. al.* 2012. Mitofusin 2 (Mfn2) links mitochondrial and endoplasmic reticulum function with insulin signaling and is essential for normal glucose homeostasis. *Proceedings of the National Academy of Sciences of the United States of America.* 109(14):5523–5528.
- Segal KR, Landt M, Klein S. 1996. Relationship Between Insulin Sensitivity and Plasma Leptin Concentration in Lean and Obese Men. *Diabetes.* 45(July):988–991.
- Seino S. 2012. Cell signalling in insulin secretion: The molecular targets of ATP, cAMP and sulfonylurea. *Diabetologia.* 55(8):2096–2108.
- Serra V, Markman B, Scaltriti M, Eichhorn PJA, Valero V, Guzman M, Botero ML, Llonch E, Atzori F, Di Cosimo S, *et. al.* 2008. NVP-BEZ235, a dual PI3K/mTOR inhibitor, prevents PI3K signaling and inhibits the growth of cancer cells with activating PI3K mutations. *Cancer Research.* 68(19):8022–8030.
- Seuring T, Goryakin Y, Suhrcke M. 2015. The impact of diabetes on employment in Mexico. *Economics and Human Biology.* 18:85–100.
- Sharif F, Muhajir H, Amin I, Zainah A. 2015. Effects of oil palm (*Elaeis guineensis*) fruit extracts on insulin secretion from BRIN BDII cells. *Malaysian Journal of Nutrition.* 21(1):75-82.
- Shi L, Yu X, Yang H, Wu X. 2013. Advanced Glycation End Products Induce Human Corneal Epithelial Cells Apoptosis through Generation of Reactive Oxygen Species and Activation of JNK and p38 MAPK Pathways. *PLoS ONE.* 8(6):1–10.
- Shimada K, Fujikawa K, Yahara K, Nakamura T. 1992. Antioxidative Properties of Xanthan on the Autoxidation of Soybean Oil in Cyclodextrin Emulsion. *Journal of Agricultural and Food Chemistry.* 40(6):945–948.
- Shimobayashi M, Albert V, Woelnerhanssen B, Frei IC, Weissenberger D, Meyer-Gerspach AC, Clement N, Moes S, Colombi M, Meier JA, *et. al.* 2018. Insulin resistance causes inflammation in adipose tissue. *Journal of Clinical Investigation.* 128(4):1538–1550.
- Shivanagoudra SR, Perera WH, Perez JL, Athrey G, Sun Y, Wu CS, Jayaprakasha GK, Patil BS. 2019. In vitro and in silico elucidation of antidiabetic and anti-inflammatory activities of bioactive compounds from *Momordica charantia L.* *Bioorganic and Medicinal Chemistry.*(March):1–13.
- Shyng SL, Ferrigni T, Nichols CG. 1997. Regulation of K(ATP) channel activity by diazoxide and MgADP: Distinct functions of the two nucleotide binding folds of the sulfonylurea receptor. *Journal of General Physiology.* 110(6):643–654.

- Siew YY, Zareisdehizadeh S, Seetoh WG, Neo SY, Tan CH, Koh HL. 2014. Ethnobotanical survey of usage of fresh medicinal plants in Singapore. *Journal of Ethnopharmacology*. 155(3):1450–1466.
- Singh P, Bano N, Hossain MM, Basit R, Dar MJ. 2019. p110 α and p110 β isoforms of PI3K are involved in protection against H₂O₂ induced oxidative stress in cancer cells. *Breast Cancer*. 26(3):378–385.
- Singh V, Raidoo DM, Harries CS. 2004. The prevalence, patterns of usage and people's attitude towards complementary and alternative medicine (CAM) among the Indian community in Chatsworth, South Africa. *BMC Complementary and Alternative Medicine*. 4:1–7.
- Singh VP, Bali A, Singh N, Jaggi AS. 2014. Advanced Glycation End Products and Diabetic Complications. 18:1–14.
- Singleton VL, Orthofer R, Lamuela-Raventós RM. 1998. Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Methods in Enzymology*. 299(1974):152–178.
- Singleton VL, Rossi JAJ. 1965. Colorimetry to total phenolics with phosphomolybdic acid reagents. *American journal of enology and viniculture*. 16(48):144–58.
- Siqueira CFDQ, Cabral DLV, Peixoto Sobrinho TJDS, De Amorim ELC, De Melo JG, Araújo TADS, De Albuquerque UP. 2012. Levels of tannins and flavonoids in medicinal plants: Evaluating bioprospecting strategies. *Evidence-based Complementary and Alternative Medicine*. 2012:1-7.
- Sivitz WI, Walsh SA, Morgan DA, Thomas MJ, Haynes WG. 1997. Effects of Leptin on Insulin Sensitivity in Normal Rats. *Endocrinology*. 138(8):3395–3401.
- Sohn JW, Harris LE, Berglund ED, Liu T, Vong L, Lowell BB, Balthasar N, Williams KW, Elmquist JK. 2013. Melanocortin 4 receptors reciprocally regulate sympathetic and parasympathetic preganglionic neurons. *Cell*. 152(3):612–619.
- Sokolovska J, Isajevs S, Sugoka O, Sharipova J, Paramonova N, Isajeva D, Rostoka E, Sjakste T, Kalvinsh I, Sjakste N. 2012. Comparison of the effects of glibenclamide on metabolic parameters, GLUT1 expression, and liver injury in rats with severe and mild streptozotocin-induced diabetes mellitus. *Medicina (Lithuania)*. 48(10):532–543.
- Sousa Silva M, Gomes RA, Ferreira AEN, Ponces Freire A, Cordeiro C. 2013. The glyoxalase pathway: The first hundred years... and beyond. *Biochemical Journal*. 453(1):1–15.
- Spauwen PJJ, Van Eupen MGA, Köhler S, Stehouwer CDA, Verhey FRJ, Van Der Kallen CJH, Sep SJS, Koster A, Schaper NC, Dagnelie PC, *et. al.* 2015. Associations of advanced glycation end-products with cognitive functions in individuals with and without type 2 diabetes: The Maastricht study. *Journal of Clinical Endocrinology and Metabolism*. 100(3):951–960.
- Stanford KI, Middelbeek RJW, Townsend KL, An D, Nygaard EB, Hitchcox KM, Markan KR, Nakano K, Hirshman MF, Tseng Y, *et. al.* 2013. Brown adipose

- tissue regulates glucose homeostasis and insulin sensitivity. *The Journal of Clinical Investigation*. 123(1):215–223.
- Stepanenko AA, Dmitrenko V V. 2015. Pitfalls of the MTT assay: Direct and off-target effects of inhibitors can result in over/underestimation of cell viability. *Gene*. 574(2):193–203.
- Stephan D, Winkler M, Kühner P, Russ U, Quast U. 2006. Selectivity of repaglinide and glibenclamide for the pancreatic over the cardiovascular KATP channels. *Diabetologia*. 49(9):2039–2048.
- Stockert JC, Blázquez-Castro A, Cañete M, Horobin RW, Villanueva Á. 2012. MTT assay for cell viability: Intracellular localization of the formazan product is in lipid droplets. *Acta Histochemica*. 114(8):785–796.
- Sulaiman MR, Hussain MK, Zakaria ZA, Somchit MN, Moin S, Mohamad AS, Israf DA. 2008. Evaluation of the antinociceptive activity of *Ficus deltoidea* aqueous extract. *Fitoterapia*. 79(7–8):557–561.
- Szendroedi J, Yoshimura T, Phielix E, Koliaki C, Marcucci M, Zhang D, Jelenik T, Müller J, Herder C, Nowotny P, *et al.* 2014. Role of diacylglycerol activation of PKC θ in lipid-induced muscle insulin resistance in humans. *Proceedings of the National Academy of Sciences of the United States of America*. 111(26):9597–9602.
- Taddeo EP, Laker RC, Breen DS, Akhtar YN, Kenwood BM, Liao JA, Zhang M, Fazakerley DJ, Tomsig JL, Harris TE, *et al.* 2014. Opening of the mitochondrial permeability transition pore links mitochondrial dysfunction to insulin resistance in skeletal muscle. *Molecular Metabolism*. 3(2):124–134.
- Taguchi A, Kislinger T, Pischetsrieder M, Hofmann M, Stern D, Yan SF, Schmidt AM, Huber B, Fu C, Qu W, *et al.* 2002. N ϵ -(Carboxymethyl)Lysine Adducts of Proteins Are Ligands for Receptor for Advanced Glycation End Products That Activate Cell Signaling Pathways and Modulate Gene Expression. *Journal of Biological Chemistry*. 274(44):31740–31749.
- Takamoto I, Kubota N, Nakaya K, Kumagai K, Hashimoto S, Kubota T, Inoue M, Kajiwara E, Katsuyama H, Obata A, *et al.* 2014. TCF7L2 in mouse pancreatic beta cells plays a crucial role in glucose homeostasis by regulating beta cell mass. *Diabetologia*. 57(3):542–553.
- Tan KCB, Shiu SWM, Wong Y, Tam X. 2011. Serum advanced glycation end products (AGEs) are associated with insulin resistance. *Diabetes/Metabolism Research and Reviews*. 27(5):488–492.
- Tan KK, Kim KH. 2013. *Alternanthera sessilis* red ethyl acetate fraction exhibits antidiabetic potential on obese type 2 diabetic rats. *Evidence-based Complementary and Alternative Medicine*. 2013:1-8.
- Tan MJ, Ye JM, Turner N, Hohnen-Behrens C, Ke CQ, Tang CP, Chen T, Weiss HC, Gesing ER, Rowland A, *et al.* 2008. Antidiabetic Activities of Triterpenoids Isolated from Bitter Melon Associated with Activation of the AMPK Pathway. *Chemistry and Biology*. 15(3):263–273.

- Thirone ACP, Carvalheira JBC, Hirata AE, Velloso LA, Saad MJA. 2004. Regulation of Cbl-Associated Protein/Cbl Pathway in Muscle and Adipose Tissues of Two Animal Models of Insulin Resistance. *Endocrinology*. 145(1):281–293.
- Tirosh A, Shai I, Tekes-Manova D, Israeli E, Pereg D, Shochat T, Kochba I, Rudich A. 2005. Normal Fasting Plasma Glucose Levels and Type 2 Diabetes in Young Men. *New England Journal of Medicine*. 353(14):1454–1462.
- Tosi F, Muggeo M, Brun E, Spiazzi G, Perobelli L, Zanolin E, Gori M, Coppini A, Moghetti P. 2003. Combination treatment with metformin and glibenclamide versus single-drug therapies in type 2 diabetes mellitus: A randomized, double-blind, comparative study. *Metabolism*. 52(7):862–867.
- Toumillehto J, Lindström J, Eriksson JG, Valle TT, Uusitupa M. 2001. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *The New England journal of medicine*. 344(18):1343–1350.
- Trebbak JT, Glund S, Deshmukh A, Klein DK, Long YC, Jensen TE, Jørgensen SB, Viollet B, Andersson L, Neumann D, *et al.* 2006. AMPK-mediated AS160 phosphorylation in skeletal muscle is dependent on AMPK catalytic and regulatory subunits. *Diabetes*. 55(7):2051–2058.
- Tucker SJ, Gribble FM, Zhao C, Trapp S, Ashcroft FM. 1997. Truncation of Kir6.2 produces ATP-sensitive K⁺ channels in the absence of the sulphonylurea receptor. *Nature*. 387(6629):179–183.
- Umpierrez G, Issa M, Vlaisnjic A. 2006. Glimepiride versus pioglitazone combination therapy in subjects with type 2 diabetes inadequately controlled on metformin monotherapy: Results of a randomized clinical trial. *Current Medical Research and Opinion*. 22(4):751–759.
- Vanhaesebroeck B, Waterfield MD. 1999. Signaling by distinct classes of phosphoinositide 3-kinases. *Experimental Cell Research*. 253(1):239–254.
- Vendrame F, Pileggi A, Laughlin E, Allende G, Martin-Pagola A, Molano RD, Diamantopoulos S, Standifer N, Geubtner K, Falk BA, *et al.* 2010. Recurrence of type 1 diabetes after simultaneous pancreas-kidney transplantation, despite immunosuppression, is associated with autoantibodies and pathogenic autoreactive CD4 T-cells. *Diabetes*. 59(4):947–957.
- Vianna R, Brault A, Martineau LC, Couture R, Arnason JT, Haddad PS. 2011. In vivo anti-diabetic activity of the ethanolic crude extract of *Sorbus decora* C.K.Schneid. (Rosacea): A medicinal plant used by Canadian James Bay Cree nations to treat symptoms related to diabetes. *Evidence-based Complementary and Alternative Medicine*. 2011(2011):237941.
- Volden S, Wimmelmann CL, Flensburg-Madsen T. 2017. Does vital exhaustion increase the risk of type 2 diabetes? A prospective study. *Journal of Psychosomatic Research*. 99(February):82–88.
- Wan Mustaffa NAA, Hasham R, Sarmidi MR. 2015. An in vitro study of wound healing

- activity of *Ficus deltoidea* leaf extract. Jurnal Teknologi. 77(3):67–72.
- Wang N, Li T, Han P. 2016. The Effect of Tianmai Xiaoke Pian on Insulin Resistance through PI3-K / AKT Signal Pathway. Diabetes Research. 2016(2016):9261259.
- Wang Y, Xiang L, Wang C, Tang C, He X. 2013. Antidiabetic and Antioxidant Effects and Phytochemicals of Mulberry Fruit (*Morus alba* L.) Polyphenol Enhanced Extract. PLoS ONE. 8(7):e71144.
- Weber MA, Bakris GL, Jamerson K, Weir M, Kjeldsen SE, Devereux RB, Velazquez EJ, Dahlöf B, Kelly RY, Hua TA, *et. al.* 2010. Cardiovascular Events During Differing Hypertension Therapies in Patients With Diabetes. Journal of the American College of Cardiology. 56(1):77–85.
- Wensveen FM, Jelenčić V, Valentić S, Šestan M, Wensveen TT, Theurich S, Glasner A, Mendrila D, Štimac D, Wunderlich FT, *et. al.* 2015. NK cells link obesity-induced adipose stress to inflammation and insulin resistance. Nature Immunology. 16(4):376–385.
- Wetzels S, Vanmierlo T, Scheijen JLJM, Van Horssen J, Amor S, Somers V, Schalkwijk CG, Hendriks JJA, Wouters K. 2019. Methylglyoxal-derived advanced glycation endproducts accumulate in multiple sclerosis lesions. Frontiers in Immunology. 10(APR):1–9.
- Winer DA, Winer S, Shen L, Wadia PP, Yantha J, Paltser G, Tsui H, Wu P, Davidson MG, Alonso MN, *et. al.* 2011. B Lymphocytes Promote Insulin Resistance through Modulation of T Lymphocytes and Production of Pathogenic IgG Antibody. Nature Medicine. 17(5):610–617.
- Withers DJ, Gutierrez JS, Towery H, Burks DJ, Ren JM, Previs S, Zhang Y, Bernal D, Pons S, Shulman GI, *et. al.* 1998. Disruption of IRS-2 causes type 2 diabetes in mice. Nature. 391:900–904.
- Woon SM, Seng YW, Ling APK, Chye SM, Koh RY. 2014. Anti-adipogenic effects of extracts of *Ficus deltoidea* var. *deltoidea* and var. *angustifolia* on 3T3-L1 adipocytes. Journal of Zhejiang University. 15(3):295–302.
- World Health Organization. 2016. Global Report on Diabetes.
- Wu CH, Yen GC. 2005. Inhibitory effect of naturally occurring flavonoids on the formation of advanced glycation endproducts. Journal of Agricultural and Food Chemistry. 53(8):3167–3173.
- Wu JW, Hsieh CL, Wang HY, Chen HY. 2009. Inhibitory effects of guava (*Psidium guajava* L.) leaf extracts and its active compounds on the glycation process of protein. Food Chemistry. 113(1):78–84.
- Xia X, Yan J, Shen Y, Tang K, Yin J, Zhang Y, Yang D, Liang H, Ye J, Weng J. 2011. Berberine improves glucose metabolism in diabetic rats by inhibition of hepatic gluconeogenesis. PLoS ONE. 6(2):1–10.
- Xiao B, Sanders MJ, Underwood E, Heath R, Mayer F V., Carmena D, Jing C, Walker

- PA, Eccleston JF, Haire LF, *et. al.* 2011. Structure of mammalian AMPK and its regulation by ADP. *Nature*. 472(7342):230–233.
- Xu M, Xiao Y, Yin J, Hou W, Yu X, Shen L, Liu F, Wei L, Jia W. 2014. Berberine promotes glucose consumption independently of AMP-activated protein kinase activation. *PLoS ONE*. 9(7):1–7.
- Xu XJ, Gauthier MS, Hess DT, Apovian CM, Cacicedo JM, Gokce N, Farb M, Valentine RJ, Ruderman NB. 2012. Insulin sensitive and resistant obesity in humans: AMPK activity, oxidative stress, and depot-specific changes in gene expression in adipose tissue. *Journal of Lipid Research*. 53(4):792–801.
- Yamagishi S, Inagaki Y, Okamoto T, Amano S, Koga K, Takeuchi M, Makita Z. 2002. Advanced Glycation End Product-induced Apoptosis and Overexpression of Vascular Endothelial Growth Factor and Monocyte Chemoattractant Protein-1 in Human-cultured Mesangial Cells. *The Journal of biological chemistry*. 277(23):20309–20315.
- Yan YX, Xiao HB, Wang SS, Zhao J, He Y, Wang W, Dong J. 2016. Investigation of the relationship between chronic stress and insulin resistance in a Chinese population. *Journal of Epidemiology*. 26(7):355–360.
- Yang SJ, Choi JM, Park SE, Rhee EJ, Lee WY, Oh KW, Park SW, Park CY. 2015. Preventive effects of bitter melon (*Momordica charantia*) against insulin resistance and diabetes are associated with the inhibition of NF- κ B and JNK pathways in high-fat-fed OLETF rats. *Journal of Nutritional Biochemistry*. 26(3):234–240.
- Yao D, Brownlee M. 2010. Hyperglycemia-induced reactive oxygen species increase expression of the receptor for advanced glycation end products (RAGE) and RAGE ligands. *Diabetes*. 59(1):249–255.
- Yap A, Nishiumi S, Yoshida KI, Ashida H. 2007. Rat L6 myotubes as an in vitro model system to study GLUT4-dependent glucose uptake stimulated by inositol derivatives. *Cytotechnology*. 55(2–3):103–108.
- Yoshioka Y, Harada E, Ge D, Imai K, Katsuzaki H, Mishima T, Gabazza EC, Ashida H. 2017. Adenosine isolated from *Grifola gargal* promotes glucose uptake via PI3K and AMPK signalling pathways in skeletal muscle cells. *Journal of Functional Foods*. 33:268–277.
- Yu Q, Shuai H, Ahooghalandari P, Gylfe E, Tengholm A. 2019. Glucose controls glucagon secretion by directly modulating cAMP in alpha cells. *Diabetologia*. 62(7):1212–1224.
- Yu Z, Liu J, Zhang R, Huang X, Sun T, Wu Y, Hambly BD, Bao S. 2017. IL-37 and 38 signalling in gestational diabetes. *Journal of Reproductive Immunology*. 124(September):8–14.
- Zahra MASF, Mahmood AA, Hapipah MA, Suzita MN, Salmah I. 2009. Anti-ulcerogenic activity of aqueous extract of *Ficus deltoidea* against ethanol-induced gastric mucosal injury in rats. *Research Journal of Medical Sciences*. 3(2):42–46.

- Zaid SSM, Othman S, Kassim NM. 2018. Protective role of *Ficus deltoidea* against BPA-induced impairments of the follicular development, estrous cycle, gonadotropin and sex steroid hormones level of prepubertal rats. *Journal of Ovarian Research*. 11(1):1–9.
- Zakaria ZA, Hussain MK, Mohamad AS, Abdullah FC, Sulaiman MR. 2012. Anti-inflammatory activity of the aqueous extract of *Ficus deltoidea*. *Biological Research for Nursing*. 14(1):90–97.
- Zargari Samani O, Mahmoodnia L, Izad M, Shirzad H, Jamshidian A, Ghatrehsamani M, Kheiri S, Sadeghian L, Soltani A, Sarmadi A. 2018. Alteration in CD8 + T cell subsets in enterovirus-infected patients: An alarming factor for type 1 diabetes mellitus. *The Kaohsiung Journal of Medical Sciences*.(2017):1–7.
- Zhang C, Qiu C, Hu FB, David RM, van Dam RM, Bralley A, Williams MA. 2008. Maternal plasma 25-hydroxyvitamin D concentrations and the risk for gestational diabetes mellitus. *PLoS ONE*. 3(11):16–19.
- Zhang Y, Liu D. 2011. Flavonol kaempferol improves chronic hyperglycemia-impaired pancreatic beta-cell viability and insulin secretory function. *European Journal of Pharmacology*. 670(1):325–332.
- Zhao C, Dong J, Jiang T, Shi Z, Yu B, Zhu Y, Chen D, Xu J, Huo R, Dai J, *et. al.* 2011. Early second-trimester serum miRNA profiling predicts gestational diabetes mellitus. *PLoS ONE*. 6(8):e23925.
- Zhao S, Mugabo Y, Iglesias J, Xie L, Delghingaro-augusto V, Lussier R, Peyot M, Joly E, Tai B, Davis MA, *et. al.* 2014. a/b -Hydrolase Domain-6-Accessible Monoacylglycerol Controls Glucose-Stimulated Insulin Secretion. 6:993–1007.
- Zhou G, Goodyear LJ, Moller DE, Zhou G, Myers R, Li Y, Chen Y, Shen X, Fenyk-melody J, Wu M, *et. al.* 2001. Role of AMP-activated protein kinase in mechanism of metformin action Find the latest version : Role of AMP-activated protein kinase in mechanism of metformin action. *The Journal of Clinical Investigation*. 108(8):1167–1174.
- Zhou Z, Xiang Y, Ji L, Jia W, Ning G, Huang G, Yang L, Lin J, Liu Z, Hagopian WA, *et. al.* 2013. Frequency, immunogenetics, and clinical characteristics of latent autoimmune diabetes in China (LADA China Study): A nationwide, multicenter, clinic-based cross-sectional study. *Diabetes*. 62(2):543–550.
- Zisman A, Peroni OD, Abel ED, Michael MD, Mauvais-Jarvis F, Lowell BB, Wojtaszewski JFP, Hirsman MF, Virkamaki A, Goodyear LJ, *et. al.* 2000. Targeted disruption of the glucose transporter 4 selectively in muscle causes insulin resistance and glucose intolerance. *Nature Medicine*. 6(8):4–8.

LIST OF PUBLICATIONS

Publications

Nurshieren Yahaya, Nur Sumirah Mohd Dom, Zainah Adam, and Muhajir Hamid, “Insulinotropic Activity of Methanolic Extracts of *Ficus deltoidea* from Seven Varieties”, Evidence-Based Complementary and Alternative Medicine, vol. 2018, Article ID 3769874, 8 pages, 2018.

Nurshieren Yahaya, Nursumirah Mohd Dom, Affrida Abu Hasan, Zainah Adam, and Muhajir Hamid. “Evaluation of Glucose Uptake of Standardized Methanolic *Ficus deltoidea* Extracts from Seven Varieties in L6 Muscle Cells”, Diabetes Research and Clinical Practices, 2020. **(Under review)**.