



UNIVERSITI PUTRA MALAYSIA

**MICROBIOLOGICAL AND PHYSICO-CHEMICAL CHANGES
DURING FERMENTATION OF *THEOBROMA CACAO*, L.:
ISOLATION AND CHARACTERIZATION OF COCOA LIPASE.**

RATNA AGUNG SAMSUMAHARTO

FSAS 2000 27

**MICROBIOLOGICAL AND PHYSICO-CHEMICAL CHANGES
DURING FERMENTATION OF *THEOBROMA CACAO*, L.:
ISOLATION AND CHARACTERIZATION OF COCOA LIPASE.**

By

RATNA AGUNG SAMSUMAHARTO

**Thesis Submitted in Fulfilment of the Requirements for the
Degree of Master of Science in the Faculty of
Science and Environmental Studies
Universiti Putra Malaysia**

February 2000



DEDICATION

“Dedicated especially to my mom, my mom, my mom : *Khasanah*, the soul that I love most; to my father: *H. Soedarto Sastrowardojo*; to my brother; to my sisters and brothers in-law whose sacrifice and support has enabled me to complete this study successfully and to my nieces and nephews for all the love”

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Master of Science

**MICROBIOLOGICAL AND PHYSICO-CHEMICAL CHANGES
DURING FERMENTATION OF *THEOBROMA CACAO*, L.:
ISOLATION AND CHARACTERIZATION OF COCOA LIPASE**

By

RATNA AGUNG SAMSUMAHARTO

February 2000

Chairman : Associate Professor Radzali Muse, Ph.D.

Faculty : Science and Environmental Studies

A study was carried out to examine microbiological and physico-chemical changes during the fermentation of *Theobroma cacao*. Isolation, partial purification, characterization of cocoa lipase were also carried out. Results showed that several microbes were successfully detected during fermentation of cocoa beans. The maximum number of yeast colonies observed in fermented cocoa beans PBC 123 and 159 clones was 4.7×10^8 and 1.5×10^9 CFU/g. fr. wt., respectively. Acetic acid bacteria were found to be dominant, 9.6×10^7 CFU/g. fr. wt. for PBC 123 clone and 1.4×10^8 CFU/g. fr. wt. for PBC 159 clone during the third day of fermentation whilst moulds were present throughout the fermentation period of six days. Spore forming bacteria appeared to be dominant on the fifth day of fermentation period. The maximum number of colonies observed for lactic acid bacteria was 3.5×10^6

CFU/g. fr. wt. for PBC 123 clone and 3.7×10^6 CFU/g. fr. wt. for PBC 159 clone while for lipolytic bacteria of both clones were 2.1×10^4 CFU/g. fr. wt. for PBC 123 clone and 2.7×10^4 CFU/g. fr. wt. for PBC 159 clone. Several major fatty acids were successfully identified in cocoa beans during the six-days of fermentation period. The maximum palmitic acid content for PBC 123 and 159 clones was 22.44 and 28.75% w/w, stearic acid content was 34.58 and 33.52% w/w, and oleic acid content was 32.09 and 36.91% w/w, respectively.

The maximum lipase specific activity from acetone dry powder of cocoa beans was 38.72 $\mu\text{mole}/\text{min}/\text{mg}$ protein (PBC 123 clone) and 98.91 $\mu\text{mole}/\text{min}/\text{mg}$ protein (PBC 159 clone). Lipase from AcDP of cocoa beans was used for partial purification using 40-60 and 60-80% ammonium sulphate precipitation. The resulted indicated 44.73 and 60.51-fold purification with 26.74 and 33.31% recovery lipase activity, respectively. Results from SDS-PAGE analysis showed that the molecular weight of the enzyme was in between 20-45 kDa. The optimum pH for the lipase activity was 7.0–8.0. Substrate specificity determinations were performed for tributyrin, trimyristin and triolein; lipase demonstrated higher affinity for trimyristin, with K_m value of 2.63 mM. Inhibition of lipase occurred in the presence of diisopropyl flourophosphate, *N*-bromosuccinimide and 5,5-dithiobis-(-2-nitrobenzoic acid).

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PERUBAHAN MIKROBIOLOGI DAN FIZIKO-KIMIA SEMASA
PENAPAIAN *THEOBROMA CACAO*, L.: PENGASINGAN DAN
PENCIRIAN LIPASE KOKO**

Oleh

RATNA AGUNG SAMSUMAHARTO

Februari 2000

Pengerusi : Profesor Madya Radzali Muse, Ph.D.

Fakulti : Sains dan Pengajian Alam Sekitar

Satu kajian telah dilakukan untuk meneliti perubahan mikrobiologi dan fiziko-kimia semasa penapaian daripada *Theobroma cacao*. Kemudian pengasingan penulenan separa dan pencirian lipase koko telah dijalankan. Keputusan menunjukkan beberapa mikrob telah berjaya dikesan semasa penapaian biji koko. Bilangan maksimum koloni yis daripada klon PBC 123 dan 159 adalah masing-masing 4.7×10^8 and 1.5×10^9 CFU/g. berat basah. Bilangan koloni yis telah dikesan pada hari pertama penapaian. Bakteria asid asetik di dapati dominan pada hari ketiga penapaian, 9.6×10^7 CFU/g. berat basah bagi klon PBC 123 and 1.4×10^8 CFU/g. berat basah bagi klon PBC 159. Bilangan maksimum koloni bakteria asid laktik adalah 3.5×10^6 CFU/ g. berat basah untuk klon PBC 123 and 3.7×10^6 CFU/g. berat basah untuk klon PBC 159. Manakala kulat hadir di sepanjang tempoh enam hari penapaian.

Bakteria pembentuk spora didapati dominan sehingga hari kelima penapaian. Manakala bagi bakteria lipolitik pula adalah 2.1×10^4 CFU/g. berat basah bagi klon PBC 123 dan 2.7×10^4 CFU/g. berat basah bagi klon PBC 159. Asid lemak utama telah berjaya dikenalpasti dan ditentukan di dalam biji koko semasa enam hari tempoh penapaian. Kandungan maksimumnya bagi setiap klon PBC 123 dan 159, adalah masing-masing 22.44 dan 28.75% w/w bagi asid palmitik, 34.58 dan 33.52% w/w bagi asid stearik serta 32.09 dan 36.91% w/w bagi asid oleik.

Aktiviti lipase spesifik maksimum daripada serbuk kering aseton biji koko adalah 38.72 $\mu\text{mole}/\text{min}/\text{mg}$ protein (klon PBC 123) dan 98.91 $\mu\text{mole}/\text{min}/\text{mg}$ protein (klon PBC 159). Lipase daripada serbuk kering aseton biji koko telah digunakan di dalam penulenan separa melalui 40-60 dan 60-80% pemendapan amonium sulfat telah menghasilkan 44.73 dan 60.51-kali penulenan berperingkat. Perolehan semula bagi penulenan tersebut adalah 26.74 dan 33.31%. Penentuan berat molekul dengan menggunakan SDS-PAGE telah menunjukkan nilai di antara 20–45 kDa. pH optima bagi lipase adalah 7.0–8.0. Kespesifikan substrat telah ditunjukkan oleh tributirin, trimiristin dan triolein, di mana lipase menunjukkan afiniti tertinggi bagi trimiristin dengan nilai K_m adalah 2.63 mM. Perencatan lipase telah berlaku dengan kehadiran suatu julat bahan kimia, termasuklah diisopropil florofosfat, *N*-bromosuksinimida dan 5,5-ditiobis-(-2-asid nitrobenzoik).

ACKNOWLEDGEMENTS

In the name of Allah The Beneficient and The Compassionate.

First of all, I would like to convey thanks and praises to The Almighty of Allah for blessing and guiding me in completing this thesis. I would also like to express my sincere gratitude to my supervisor, Associate Professor Dr. Radzali Muse, for his continuous guidance and encouragement throughout this project. Sincerely thanks to Professor Dr. Jinap Selamat and Associate Professor Dr. Johari Ramli for their supports, invaluable technical advices and critics during this study.

Special thank goes to Associate Professor Dr. Khatijah Mohd. Yusoff, Head Department of the Biochemistry and Microbiology, Universiti Putra Malaysia, Serdang, Selangor Darul Ehsan, Malaysia for allowing me to use all research facilities.

I am also indebted to all departmental staffs especially Mr. Ismail Omar, Mr. Onn Hj. Ahmad, Mr. Ibrahim Yunus, Mr. Hussain Jirangon and Mr. Khalid Ithnin of their help and cooperation.

I wish to thank all postgraduate club members particularly labmates e.g. Mohd. Yunus Shukor. Arief Zaidi Jusoh, Suri Roowi, Jeeven Karruppan,

Shakinaz Desa, Fezah Othman, Asfaliza Ramli and Loo Wai Sum for their generous assistance, encouragement, support and humour.

Last but not least, I would like to express my gratitude to my family for their pray, love, understanding and patience during the study.

APPROVAL SHEETS


I certify that an Examination Committee met on 29th February, 2000 to conduct the final examination of Ratna Agung Samsumaharto on his Master of Science thesis entitled “Microbiological and Physico-chemical Changes during Fermentation of *Theobroma cacao*, L.: Isolation and Characterization of Cocoa Lipase” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

MOHD. ASPOLLAH SUKARI, Ph.D.,
Associate Professor,
Faculty of Science and Environmental Studies,
Universiti Putra Malaysia.
(Chairman)

RADZALI MUSE, Ph.D.,
Associate Professor,
Faculty of Science and Environmental Studies,
Universiti Putra Malaysia.
(Member)

JINAP SELAMAT, Ph.D.,
Professor,
Faculty of Biotechnology and Food Science,
Universiti Putra Malaysia.
(Member)

JOHARI RAMLI, Ph.D.,
Associate Professor,
Faculty of Science and Environmental Studies,
Universiti Putra Malaysia.
(Member)



MOHD. GHAZALI MOHAYIDIN, Ph.D.,
Professor/Deputy Dean of Graduate School,
Universiti Putra Malaysia.

Date: 20 APR 2000

This thesis was submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfilment of the requirements for the degree of Master of Science.




KAMIS AWANG, Ph.D
Associate Professor
Dean of Graduate School,
Universiti Putra Malaysia

Date: 11 MAY 2000

DECLARATION FORM

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



Candidate.

Name : RATNA AGUNG SAMSUMAHARTO

Date : 19 April 2000

TABLE OF CONTENTS

	Page
DEDICATION.....	ii
ABSTRACT.....	iii
ABSTRAK.....	v
ACKNOWLEDGEMENT.....	vii
APPROVAL SHEETS.....	ix
DECLARATION FORM.....	xi
LIST OF TABLES.....	xv
LIST OF FIGURES.....	xvi
LIST OF PLATES.....	xix
LIST OF ABBREVIATIONS.....	xx

CHAPTER

I	GENERAL INTRODUCTION.....	1
	Objectives of The Study.....	5
II	LITERATURE REVIEW.....	6
	Fermentation.....	6
	Factors Affecting on Cocoa Beans Fermentation.	8
	Ripeness of Pods.....	8
	Pulp Preconditioning.....	9
	Pod Diseases.....	12
	Type of Cocoa.....	12
	Quantity of Cocoa	13
	Duration.....	14
	Substrate.....	14
	Turning (Aeration).....	15
	Seasonal Effects.....	19
	Changes Occurring during Fermentation.....	19
	Temperature Changes during Cocoa Beans Fermentation.....	21
	pH Changes during Cocoa Beans Fermentation.....	23
	Microbiological Changes in The Cocoa Pulp.....	25
	Organic Acids Development during Cocoa Beans Fermentation.....	31



Sources of Lipase.....	35
Reactions Catalyzed by Lipases.....	39
Hydrolysis and Esterification.....	40
Transesterification.....	42
Interesterification.....	42
Lipase Activity in Plants.....	43
Purification of Lipase.....	45
Characterization of Lipase.....	47
Effect of Temperature and pH on Lipase Activity.....	47
Effect of Inhibitors on Lipase Activity.....	48
Effect of Substrates on Lipase Activity.....	50
 III	
GENERAL MATERIALS AND METHODS.....	52
Materials.....	52
Methods.....	53
Fermentation and Sampling.....	53
Preparation of Acetone Dry Powder.....	54
Extraction of Crude Enzyme from AcDP..	55
Determination of Soluble Protein Content	55
Substrate Emulsion Preparation.....	56
Assay of Lipase Activity.....	56
 IV	
MICROBIOLOGICAL AND PHYSICO-CHEMICAL CHANGES DURING COCOA BEANS FERMENTATION.....	58
Introduction.....	58
Materials and Methods.....	62
Statistical Analysis.....	70
Results and Discussion.....	70
Fermentation.....	70
Temperature Changes.....	72
pH and Titratable Acidity Changes.....	75
Fermentation Index Changes.....	79
Yeast and Acetic Acid Bacteria, Lactic Acid Bacteria, Spore Forming Bacteria Population	80
Lipolytic Bacteria Population.....	86
Fat Content.....	89
Fatty Acids Content.....	90
Lipase Activity in Cocoa Beans.....	93

V	ISOLATION AND CHARACTERIZATION OF COCOA LIPASE.....	95
	Introduction.....	95
	Materials and Methods.....	97
	Results and Discussion.....	107
	Isolation of Cocoa Lipase.....	107
	Partial Purification of Cocoa Lipase.....	108
	The Purity and Molecular Weight of Cocoa Lipase.....	112
	Characterization of Cocoa Lipase.....	115
VI	GENERAL DISCUSSION AND CONCLUSION	122
	BIBLIOGRAPHY.....	131
	APPENDICES.....	145
	A Additional Methodologies.	146
	B Additional Figures.....	152
	BIODATA OF AUTHOR.....	155

LIST OF TABLES

Table	Page
1 Effect of Ripeness of Pods.....	9
2 Lipase Specific Activity in Crude Enzyme from AcDP and Ammonium Sulphate Precipitated Fractions.....	108
3 Partial Purification Scheme of Lipase from AcDP of Cocoa Beans (40-60% Saturation of Ammonium Sulphate Precipitation).....	113
4 Partial Purification Scheme of Lipase from AcDP of Cocoa Beans (60-80% Saturation of Ammonium Sulphate Precipitation).....	114
5 Effect of Inhibitors on Lipase Specific Activity of Partially Purified Lipase Fractions from AcDP of Cocoa Beans.....	120

LIST OF FIGURES

Figure	Page
1 Stages in The Lipase Catalysed Hydrolysis of An Ester.....	41
2 Temperature Profile of Cocoa Beans Mass during Six-Days of Fermentation.....	74
3 pH Profile of Cocoa Beans Mass during Six-Days of Fermentation.....	77
4 Titratable Acidity Profile of Cocoa Beans Mass during Six-Days of Fermentation.....	78
5 Fermentation Index Profile of Cocoa Beans Mass during Six-Days of Fermentation.....	80
6 Population of Microbial Profile in Cocoa Beans of PBC 159 Clone during Six-Days of Fermentation...	83
7 Population of Microbial Profile in Cocoa Beans of PBC 123 Clone during Six-Days of Fermentation..	84
8 Population of Lipolytic Bacteria Profile in Cocoa Beans of PBC 123 and 159 Clones.during Six-Days of Fermentation.....	87
9 Fat Content Profile in Cocoa Beans of PBC 123 and 159 Clones during Six-Days of Fermentation.	89
10 Fatty Acid Content Profile in Cocoa Beans of PBC 159 Clone during Six-Days of Fermentation.	91
11 Fatty Acids Content Profile in Cocoa Beans of PBC 123 Clone during Six-Days of Fermentation.	92

12 Lipase Specific Activity in Crude Extract and AcDP Extract of Cocoa Beans during Six-Days of Fermentation.....	94
13 Elution Profile of Soluble Lipase from 40-60% Saturation of Ammonium Sulphate Precipitation Obtained by DEAE-cellulose Chromatography. Lipase Enzyme Was Eluted from Column using a 0.0 to 0.5 M NaCl Gradient.....	109
14 Elution Profile of Soluble Lipase from 60-80% Saturation of Ammonium Sulphate Precipitation Obtained by DEAE-cellulose Chromatography. Lipase Enzyme Was Eluted from Column using a 0.0 to 0.5 M NaCl Gradient.....	110
15 Elution Profile of Soluble Lipase from 40-60% Saturation of DEAE-cellulose Elution Obtained Was Obtained by Sephacryl S-200 Gel Filtration Chromatography.....	111
16 Elution Profile of Soluble Lipase from 60-80% Saturation of DEAE-cellulose Elution Obtained Was Obtained by Sephacryl S-200 Gel Filtration Chromatography.....	111
17 Effect of Temperature on Lipase Specific Activity of Cocoa Beans.....	116
18 Effect of pH on Lipase Specific Activity of Cocoa Beans.....	117
19 Lineweaver-Burk Double Reciprocal Plots for The Partially Purified Lipase using Tributyrin, Trimyristin and Triolein as Substrates.....	118
20 Chromatographic of FAMES Standard (A) C14:0= Myristic Acid, (B) C16:0= Palmitic Acid, (C) C18:0= Stearic Acid, (D) C18:1=Oleic Acid..	152
21 Chromatographic of Fatty Acids of Cocoa Beans PBC 159 Clone (A) C16:0= Palmitic Acid, (B) C18:0= Stearic Acid, (C) C18:1= Oleic Acid	153

22	Chromatographic of Fatty Acids of Cocoa Beans	
	PBC 123 Clone (A) C16:0= Palmitic Acid,	
	(B)C18:0= Stearic Acid, (C) C18:1= Oleic	
	Acid).....	154

TABLE OF PLATES

Plate	Page
1 Ripe Fruit of <i>Theobroma cacao</i> , L of PBC 123 and 159 Clones.....	52
2 Non-fermented Beans of <i>Theobroma cacao</i> , L. of PBC 123 and 159 Clones.....	71
3 Fermented Beans of <i>Theobroma cacao</i> , L. of PBC 123 and 159 Clones.....	71
4 Presumptive Test of Lipolytic Bacteria in Tributyrin Agar.....	88
5 Population of Lipolytic Bacteria in Oleic Agar.....	88
6 DEAE-cellulose Ion Exchanger Medium.....	101
7 Sephacryl S-200 Gel Filtration Medium.....	103
8 Sodium Dodecyl Sulphate – Polyacrylamide Gel Electrophoreograms of Cocoa Lipase.....	114

LIST OF ABBREVIATIONS

CFU	- colonies forming unit
g	- gram
g. fr. wt.	- gram fresh weight
(w/w)	- weight/weight
%	- per cent
min	- minute
mg	- milli gram
μmole	- micro mole
kDa	- kilo dalton
mM	- milli molar
μM	- micro molar
ha	- hectare
EC	- enzyme commission
FFA	- free fatty acid
min	- minute
kg	- kilo gram
mL	- meter length
mW	- meter wide
mD	- meter deep
cm	- centi meter
(w/v)	- weight/volume
dwb	- dry weight basis
ml	- milli liter
nm	- nanno meter
AcDP	- acetone dry powder
rpm	- revolution per minute
UV/Vis	- ultra violet/visible
U	- unit
meq	- milli equivalent
BF ₃	- borontriflouride
M	- molar
μl	- micro liter
N	- normality
V/cm	- volt/centi meter
°C	- degree Celcius
Abs	- absorbance
TEMED	- N,N,N',N'-tetramethyl-ethylenediamine
DFP	- diisoprophylflourophosphate

CHAPTER I

GENERAL INTRODUCTION

The fermentation technique is one of the most important factors in determining the quality of cured cocoa beans. According to Lehrian and Patterson (1983), cocoa fermentation is a complex natural process involving a mixture of external microbiological processes occurring in the pulp surrounding the beans and internal structural changes and enzymatic reactions. A succession of microorganism, in particular yeast, lactic acid bacteria, and acetic acid bacteria, grow rapidly in the pulp producing ethanol, lactic acid and acetic acid as major metabolic products (Lehrian and Patterson, 1983). Acetic acid in particular is responsible for the death of the beans, prevents germination, solubilizes polyphenols, aids in diffusing the content of the storage cells into surrounding parenchyma tissue, and prevents the beans from attack by putrefactive bacteria (Ziegleder and Biehl, 1988).

Acids could mask the overall chocolate flavour in cocoa beans (Lopez and McDonald, 1981). However, high concentrations of residual acids may cause the beans to be excessively acidic in flavour. The presence of acetic and lactic acids either alone or combined, have been implicated as

possible causes of high acidic flavour in cocoa beans. Some researchers suggest that only acetic acid is important because it is present in high concentration and tastes more acidic than other acids (Rohan and Stewart, 1964; Biehl, 1965; Lopez, 1983; Jinap and Dimicks, 1990). Holm (1991) reported that oxalic acids could improve cocoa flavour in Malaysian cocoa beans.

The exothermic formation of acetic acid may physiologically cause the temperature of fermenting mass to rise, prevent germination and cause structural changes which remove the compartment action of enzymes and substrates, thereby permitting an increase in enzymes activity (Biehl and Adomako, 1983).

According to Horman and Braco (1986), Staphylakis and Gegiou (1985) the cocoa cotyledon contains common fatty acids such as 25-30% palmitic acid, 32-37% oleic acid and 30-37% stearic acid, 2-4% linolenic acid and 0.7-1% arachidonic acids. It is generally known that fatty acids have an important effect on chocolate flavour. The high content of free fatty acids in cocoa beans cause dull, insipid and flat tastes (Padavatan *et al.*, 1979). A variety of saturated and unsaturated fatty acids is present in triacylglycerols (TAGs) and the way they arrange themselves during crystal formation determines the hardness of the cocoa butter. Among the most abundant fatty acids are myristic (14:0), palmitic (16:0), stearic (18:0) and

oleic (18:1^{Δ9}) acids . Fatty acids content in triacylglycerols (TAGs) are most easily determined by using complete saponification process with NaOH and then followed by esterification of the released fatty acids (Boyer,1986; Ziegleder and Biehl, 1988).

In the cotyledon, the accumulation of acids is responsible for the formation of flavour precursors during fermentation by providing an acidic environment for enzymatic reactions to occur. Lipases (triacylglycerol acylhydrolase, EC 3.1.1.3) are ester hydrolases or esterases since they hydrolyse the ester bonds of triacylglycerol molecules. The lipases are more active with insoluble fatty acid esters and hydrolyse the ester bonds present only at the water-oil interface, whereas carboxylic ester hydrolases that are specific for the soluble esters are simply termed esterases (Schuepp *et al.*, 1997). According to Jensen *et al.* (1983), lipases or acylglycerol hydrolase are enzymes which catalyze the hydrolysis of long chain aliphatic acids from acylglycerol at the oil/water interface. The systematic name is acylglycerol acylhydrolase. The interface is usually provided by emulsion globules or lipoprotein particles, the latter are primarily chylomicrons and very low density lipoprotein. The element providing the interface has been termed the super-substrate.

Bloch (1960) reported that lipase is an enzyme which can be easily found in higher plants, animals, insects and some microorganisms. In plant

this lipolytic enzyme plays an important role in fatty acid metabolism. Highly active lipases are found to catalyze the hydrolysis of reserve triacylglycerols (TAGs). The triacylglycerols (TAGs) are actually localized in subcellular organelles called lipid bodies (Bloch, 1960). Also, lipases have the important physiological role of preparing the fatty acids of water-insoluble triglycerides (TGs) for absorption into and transport through membranes by converting the triglycerides (TGs) to the more polar diglycerides (DGs), monoglycerides (MGs), free fatty acid (FFA) and glycerol (Jensen, 1983). The majority of lipases are extracellular, acidic glycoprotein of molecular weights between 20 and 60 kDa, although some form aggregates in solution (accounting for the high molecular weights reported for some partially purified enzymes). Most purified lipases contain between 2 and 15% carbohydrate, with the major glycoside residue being mannose (Gill and Parish, 1997). The evidence for the presence of lipases in cocoa beans was first reported by Ciferrin in 1931 (as cited by Forsyth and Quesnel, 1963).

The lipolytic enzyme found in the higher plants such as cocoa as important in understanding of their physiological roles as well as their action in agricultural products during storage. The main purpose of this research is to study the changes in microbiological and physico-chemical properties on fermented cocoa beans and isolation and characterization of cocoa lipase.