



Original Article

Syzygium claviflorum fruit extract preadipocyte differentiation inhibition in 3T3-L1 cells



Rozaini M. Zohdi, PhD^{a,b,*}, Muhammad A. Adli, BSc^{a,b}, Shahida M. Mukhtar, BSc^a, Awang H. Awang Junaidi, PhD^c and Md Zuki A. Bakar, PhD^c

^a Atta-ur-Rahman Institute for Natural Product Discovery, Universiti Teknologi MARA, Selangor Branch, Puncak Alam, Selangor, Malaysia

^b Department of Pharmacology and Life Sciences, Faculty of Pharmacy, Universiti Teknologi MARA, Selangor Branch, Puncak Alam, Selangor, Malaysia

^c Department of Veterinary Pre-Clinical Science, Faculty of Veterinary Medicine, Universiti Putra Malaysia, Serdang, Selangor, Malaysia

Received 18 August 2024; revised 1 November 2024; accepted 4 December 2024; Available online 19 December 2024

المخلص

أهداف البحث: دفعت المخاوف بشأن العدد المتزايد من الأفراد الذين يعانون من السمنة المفرطة والمخاطر الصحية المرتبطة بها إلى استكشاف الخيارات العلاجية. وبالمثل، تهدف هذه الدراسة إلى إنشاء سمات مضادة للدهون في مستخلص فاكهة سيجيزيوم كلايفلوروم في خلايا 3T3-L1.

طرق البحث: تم التعرف على المركبات البوليفينولية في فاكهة سيجيزيوم كلايفلوروم باستخدام التحليل اللوني السائل عالي الأداء للمرحلة العكسية. وفي الوقت نفسه، تم استخدام الخلايا الشحمية 3T3-L1 الفارسية، وقياس مستويات الليبتين، وأنواع الأكسجين التفاعلية، ومحتويات الدهون والدهون الثلاثية خلال تقييمات النشاط المضاد للدهون. في الوقت نفسه، تم تحديد تأثيرات فاكهة سيجيزيوم كلايفلوروم على عوامل النسخ الدهنية باستخدام تفاعل البلمرة المتسلسل الكمي في الوقت الحقيقي.

النتائج: أشارت النتائج إلى ثلاثة مركبات بوليفينولية في فاكهة سيجيزيوم كلايفلوروم، بما في ذلك فلافونويد واحد (نارينجين) واثنين من الأحماض الفينولية (سرينجيك وبكوماريك). على الرغم من أن علاجات فاكهة سيجيزيوم كلايفلوروم (٢٥٠-١٠٠٠ ميكروغرام/مل) لم تؤدي إلى سمية الخلايا، إلا أنها خفضت بشكل كبير تراكم الدهون المعتمد على الجرعة، وإنتاج الأكسجين التفاعلي، ومستويات الدهون الثلاثية والليبتين بالنسبة للخلايا الشحمية المتميزة بالتحكم. علاوة على ذلك، قامت فاكهة سيجيزيوم كلايفلوروم بقمع بروتين ربط

العناصر التنظيمية للستيروول-١، العنصر التنظيمي للستيروول، ومستقبلات غاما المنشط بالبيروكسيوسوم، والتعبيرات الجينية لبروتين سي/ايبب ألفا المرتبطة بالمحسن خلال الخلية الريدية الشحمية والتميز إلى الخلايا الشحمية.

الاستنتاجات: كشفت النتائج عن خصائص فاكهة سيجيزيوم كلايفلوروم المضادة للدهون، مما يشير إلى إمكاناته كعلاج طبيعي لإدارة السمنة. ومع ذلك، هناك حاجة إلى مزيد من الدراسات لتوضيح التفاعلات التي تؤدي إلى تأثيرات فاكهة سيجيزيوم كلايفلوروم المضادة للدهون والمكونات النشطة المسؤولة عن الخاصية.

الكلمات المفتاحية: تكوين الشحوم؛ مضاد للدهون؛ في المختبر؛ بدانة؛ الخلايا التحضيرية؛ فاكهة سيجيزيوم كلايفلوروم

Abstract

Objective: Concerns over the increasing number of obese individuals and the associated health risks have prompted therapeutic option explorations. Similarly, this study aimed to establish *Syzygium claviflorum* fruit extract (SCFE) anti-adipogenic attributes in 3T3-L1 cells.

Methods: The polyphenolic compounds in SCFE were identified with Reverse phase-high performance liquid chromatography (RP-HPLC). Meanwhile, murine 3T3-L1 preadipocytes, measuring leptin levels, reactive oxygen species (ROS), and lipid and triglyceride (TG) contents were utilized during anti-adipogenic activity assessments. Concurrently, the effects of SCFE on adipogenic transcription factors were established with quantitative real-time-polymerase chain reaction (qRT-PCR).

* Corresponding address: Atta-ur-Rahman Institute for Natural Product Discovery, Universiti Teknologi MARA, Selangor Branch, Puncak Alam Campus, 42300 Puncak Alam, Selangor, Malaysia.

E-mail: rozainizohdi@uitm.edu.my (R.M. Zohdi)

Peer review under responsibility of Taibah University.



Results: The RP-HPLC results indicated three polyphenolic compounds in SCFE, including one flavonoid (naringin) and two phenolic acids (syringic and p-coumaric). Although SCFE treatments (250–1000 µg/mL) did not result in cell toxicity, they significantly reduced dose-dependent lipid accumulation, ROS production, and TG and leptin levels relative to control-differentiated adipocytes. Moreover, SCFE suppressed sterol regulatory element binding protein-1 (SREBP-1), peroxisome proliferator-activated receptor-gamma (PPAR-γ), and CCAAT/enhancer-binding protein-alpha (C/EBP-α) gene expressions during preadipocyte differentiation into adipocytes.

Conclusion: The findings revealed the anti-adipogenic properties of SCFE, indicating its potential as a natural obesity management remedy. Nevertheless, more studies are necessary to elucidate the reactions resulting in SCFE anti-adipogenic effects and the active constituents responsible for the property.

Keywords: Adipogenesis; Anti-adipogenic; *In vitro*; Obesity; Preadipocytes; *Syzygium claviflorum*

© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Obesity is characterized by excess adipose tissue buildup. The condition is a worldwide health issue due to its complex relationship to various metabolic disturbances, including type 2 diabetes mellitus, hypertension, and non-alcoholic fatty liver disease.^{1,2} Obesity pathophysiology is intricately associated with adipocytes, the primary cellular component of adipose tissue.¹ Adipocytes are required to maintain energy balance by storing excess energy in the form of triglyceride (TG).² Nonetheless, the physiological function is dysregulated in obese individuals, resulting in pathological adipocyte growth and adipose tissue dysfunction.³

Mature adipocyte derivation from preadipocytes is critical for adipose tissue growth and maturation. Although adipocyte differentiation promotes adipocyte development, the process also generates reactive oxygen species (ROS), which exerts oxidative stress in the cellular environment.⁴ A discernible rise in adipokines secretion, including adiponectin and leptin, is also observed during the activity.⁵ Complex reactions between transcription factors, such as CCAAT/enhancer-binding protein-alpha (C/EBP-α), sterol regulatory element-binding protein-1 (SREBP-1), and peroxisome proliferator-activated receptor-gamma (PPAR-γ), manage adipocyte differentiation.⁶ The transcription factors are crucial to coordinating the gene expressions related to adipogenesis and lipid metabolism and regulating critical adipocyte developmental and functional aspects.⁶

Obesity management is a significant challenge for medical interventions.⁷ Moreover, pharmaceutical approaches have often been ineffective and not safe.⁸ The questionable efficacy and possible adverse effects associated with anti-obesity medications have also raised concerns.⁷

Consequently, developing alternative anti-obesity therapies with minimum side effects and long-term efficacy is urgently required. Harnessing the potential of natural compounds for obesity management has gained much attention, specifically polyphenols, which have exhibited obesity and associated health condition mitigation effects.⁹ The compounds reportedly exert advantageous properties through numerous biochemical targets and pathways, collectively providing significant health advantages.⁹

Syzygium claviflorum (Roxb.) Wall. ex Steud. (*S. claviflorum*), or *Eugenia claviflora* Roxb., belongs to the *Myrtaceae* family.¹⁰ In Malaysia, the species is known as 'jambu arang' or 'keriang dot'. The plant possesses well-known distinctive-flavored fruits. Nevertheless, research on the nutritional and medicinal properties of *S. claviflorum* is limited.

Recent studies indicated that *S. claviflorum* fruit pulp and seed extracts are rich in polyphenols offering potent antioxidant properties.¹¹ Nonetheless, the specific health-promoting effects of *S. claviflorum*, particularly for managing obesity, are still unknown. The present study analyzed the phytochemical composition and anti-adipogenic attributes of *S. claviflorum* fruit extract (SCFE) in 3T3-L1 cells. SCFE effects on the expression of critical genes required in adipogenesis and lipid metabolism were also determined.

Materials and Methods

Harvesting the plant specimens

The *S. claviflorum* fruits employed in this study were harvested from residential fruit trees in Alor Setar, Kedah (N 6° 7' 29.2800" E 100° 22' 4.1556") in July 2022. Specimen identification was performed by a botanist from Universiti Kebangsaan Malaysia (UKM), Bangi, Malaysia, and the sample voucher (AuRIns-20220 322) was deposited in the Herbarium of AuRIns, UiTM Puncak Alam.

Dirt was removed from the harvested *S. claviflorum* fruits with multiple washes of distilled water before manually removing the peel and seeds. Finally, the fruit pulp samples were freeze-dried, crushed into powder, and stored at −20 °C before subjecting them to further procedures.

Preparing the SCFE

The extraction methodology outlined by Ahmed et al.¹¹ was applied in the present study with slight adjustments. Firstly, an ethanolic extract was acquired by adding 10 g of the powdered peel to 100 mL of 80:20 (w/v) an ethanolic aqueous solvent system. After being shaken for 24 h at room temperature, the mixture was filtered through Whatman number 1 filter paper [Millipore, Burlington, Massachusetts, United States of America (USA)]. For an optimized extraction, the residue was redissolved in the solvent. The filtrates from the procedure, which was performed in triplicates, were then combined. The resultant mixture was concentrated with a Rotavapor R-215 (Büchi, Flawil, Switzerland) set to 45 °C and 150 rpm for 90 min. Finally, the extract was lyophilized and kept at −20 °C.

The SCFE phytochemical analysis

The present study established the phytochemicals in the SCFE samples with reverse phase-high performance liquid chromatography (RP-HPLC). The methodology reported by Kong et al.¹² was utilized. A C18 column (5 µm particle size; 150 mm × 4.6 mm) [Agilent, California (CA), USA] adjusted to 30 °C was employed. This study also utilized ultrapure water added with 0.3 % formic acid and acetonitrile (ACN) as the mobile phase at an injection volume and flow rate of 10 µL and 1.0 mL/min, respectively.

Caffeic acid phenethyl ester (CAPE), gallic, syringic, cinnamic, and p-coumaric acids, which were phenolic acids, and six flavonoids; chrysin, quercetin, naringin, kaempferol, galangin, and pinocembrin (Sigma–Aldrich, Steinheim, Germany) (see Figure 1) were employed as the standards in this study. P-coumaric and cinnamic acids were determined at 300 and 280 nm, respectively, while the other standards were detected at 260 nm. The gradient sequence applied was 0–3 min, ACN: 10 %, 3–25 min, ACN: 10%–20 %, 25–45 min, ACN: 20%–45 %, 45–55 min, ACN: 45%–60 %, 55–60 min, ACN: 60%–95 %, 60–65 min, ACN: 95 %, 65–70 min, ACN: 95%–10 %, 70–72 min, ACN: 10 %. Chromatograms from triplicate assessments of the standards and the SCFE were compared.

Adipocyte cell cultivation and differentiation

The 3T3-L1 murine preadipocytes procured from the American Type Culture Collection (Manassas, Virginia, USA) employed in this study were grown in a glucose-rich Dulbecco's modified Eagle medium (DMEM) containing 1 % penicillin-streptomycin and 10 % fetal bovine serum (FBS). The guideline reported by Choi et al.¹³ was applied during adipocyte differentiation initiation, as presented in Figure 2.

Upon confluence, a differentiation medium replaced the medium employed. FBS (10 %), 3-isobutyl-1-methylxanthine (0.5 mM), dexamethasone (0.1 µM), and insulin (1 µg/mL) were employed, which was maintained for two days. Subsequently, a regular medium of 1 µg/mL insulin and 10 % FBS was utilized for cell culture for six days. A fresh medium was supplied every two days.

In this study, the influences of SCFE on adipocyte differentiation were evaluated with and without SCFE addition to the differentiation medium two days post-confluent. N-acetyl-L-cysteine (NAC) (5 mM) was employed as the positive control. After eight days, the cells were harvested.

Cell viability assay

The current study established cell viability via the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay. 96-well plates incorporated with 250, 500, and 1000 µg/mL of the SCFE was employed to place the 3T3-L1 preadipocytes for 24 h. Subsequently, cell absorbance at 570 nm was determined post-incubating the cells for 4 h with an MTT solution.

Differentiation media added with the SCFE 250, 500, and 1000 µg/mL of were employed to procure mature adipocytes.

The MTT assay was conducted to establish cell viability after eight days. Viable cell percentages were calculated according to Equation (1).

$$\text{Cell viability (\%)} = \left(\frac{\text{Absorbance of treated cells}}{\text{Absorbance of control cells}} \right) \times 100\% \quad (\text{Eq. 1})$$

Lipid accumulation determination via oil red O (ORO) staining

Minor modifications to the guidelines described by Chung and Hyun¹⁴ was applied in the present study for the ORO staining to assess the SCFE lipid inhibitory properties on differentiated adipocytes. Differentiation of the 3T3-L1 preadipocyte was induced in a 24-well plate containing 250, 500, and 1000 µg/mL of the SCFE. The cells were fixed and rinsed in 10 % formaldehyde for 1 h and phosphate-buffered saline (PBS), respectively, before drying them after eight days.

A 0.5 % ORO dissolved in isopropanol (60 % v/v) at room temperature was employed to stain the dried 3T3-L1 preadipocytes for 30 min. The samples were then washed with water and dried. An Olympus CKX53 microscope (Tokyo, Japan) was utilized for observing and photographing the stained lipid droplets. Quantification of lipid accumulation was performed by solubilizing the stained lipid droplets with isopropanol. The absorbance of the samples was determined at 490 nm with a microplate reader. The lipid accumulation percentage was established with Equation (2).

$$\text{Lipid accumulation (\%)} = \left(\frac{\text{Absorbance of treated cells}}{\text{Absorbance of control cells}} \right) \times 100\% \quad (\text{Eq. 2})$$

This study applied the procedure reported by Choi et al. (2017) during the NBT assay to investigate the effects of the SCFE samples on ROS generation. The 3T3-L1 preadipocytes were induced to differentiate into adipocytes in 96-well plates post-confluence. SCFE of 250, 500, and 1000 µg/mL were then employed to treat the sample cells. The NBT solution was added on the eighth day and left for 90 min. The cells were subjected to dissolution with formazan. Equation (3) was applied to calculate ROS formation percentages upon determining cell absorbance at 570 nm.

$$\text{ROS (\%)} = \left(\frac{\text{Absorbance of treated cells}}{\text{Absorbance of control cells}} \right) \times 100\% \quad (\text{Eq. 3})$$

TG and leptin level evaluations

A commercial colorimetric test kit (Elabsience, Houston, Texas, USA) was utilized to determine the TG levels in cell lysates in this study per the manufacturer's instructions. Post-confluence, the 3T3-L1 preadipocyte cells were treated with 250, 500, and 1000 µg/mL of SCFE in 6-well plates during the eight days of differentiation. The cell lysates were

then scraped and centrifuged at $1000\times g$ for 10 min after being subjected to PBS wash several times.

Isopropanol was added to the cell lysates to resuspend them. The 1-sec pulse sonication technique was then performed. Supernatants were procured by centrifuging the lysates at $10\,000\times g$ for 10 min. The TG deposition results were recorded as TG (mg) per cellular protein (mg). Furthermore, the Lowry technique was employed to determine the total protein content in the lysate cells.

Leptin enzyme-linked immunosorbent (ELISA) assay

The leptin levels in the adipocytes evaluated in the present study were determined with a mouse leptin immunoassay kit (Elabscience, Houston, Texas, USA) following the manufacturer's recommendations. After an eight-day treatment period, the supernatant from the culture was collected. The cells were rinsed with PBS, carefully harvested utilizing trypsinization, and centrifuged at $1000\times g$ for 10 min. Subsequently, cellular pellets were washed thrice with cold PBS before being subjected to three freezing and thawing cycles. The supernatants were analyzed for leptin levels following centrifugation at $1500\times g$ for 10 min. Finally, a standard curve was generated with leptin concentrations between 0 ng/mL and 20 ng/mL to calculate the leptin levels.

RNA extraction and quantitative real-time-polymerase chain reaction (qRT-PCR)

The current study isolated RNA from untreated and treated cells (on day 8) with an SV Total RNA extraction kit [Promega, Wisconsin (WI), USA] to assess the gene expression associated with adipocyte differentiation. Subsequently, the GoScript™ Reverse Transcriptase system (Promega, WI, USA) was employed to synthesize cDNA from the total RNA. The procedures applied followed the guidelines provided by the manufacturers.

The desired genes were amplified with an Applied Biosystems StepOne system (Life Technologies, MA, USA) utilizing 200 ng/reaction of cDNA incorporated with SYBR Green-based qPCR Master Mix (Promega, WI, USA). A total of 40 denaturation cycles at $95\text{ }^{\circ}\text{C}$ for 15 s followed by annealing and extension at $60\text{ }^{\circ}\text{C}$ for 60 s were performed. Relative gene expression was established according to the $2^{-\Delta\Delta C_t}$ technique with β -actin as the internal control. All samples were examined in triplicates and quantified per the relative standard curve approach. The primer sequences employed in this study are listed in Table 1.

Statistical analysis

The statistical analyses conducted in the current study were performed with GraphPad Prism version 7.0. The evaluations included the one-way analysis of variance (ANOVA) followed by Tukey's multiple comparison test to identify significant differences among experimental groups. The data procured were recorded as mean \pm standard deviation (SD), obtained from triplicate runs. Differences demonstrating $p < 0.05$ were considered statistically significant.

Results

RP-HPLC

Figure 3(A) and (B) illustrate the chromatograms of the standards and SCFE, respectively. The SCFE contained phenolic acids; syringic (RT: 10.446 min) and p-coumaric (RT: 15.634 min), and a flavonoid compound, naringin (RT: 24.698 min).

The effects of the SCFE on 3T3-L1 cell viability

The 3T3-L1 preadipocytes and mature adipocytes in this study were exposed to SCFE at varying concentrations (250,




Figure 1: Chemical structures of the five phenolic acids and six flavonoid standards used in the high-performance liquid chromatography (HPLC) analysis.




Figure 2: Schematic timeline of the experiment.

Table 1: Primer sequences used for quantitative reverse transcription PCR (qRT-PCR) analysis of adipogenic gene expression in 3T3-L1 adipocytes.

Gene	Forward (5'-3')	Reverse (5'-3')
PPAR γ	TTTTCAAGGGTGCCAGTTTC	AATCCTTGGCCCTCTGAGAT
C/EBP α	TTACAACAGGCCAGGTTTCC	GGCTGGCGACATACAGTACA
SREBP-1	TGTTGGCATCCTGCTATCTG	AGGGAAAGCTTGGGGTCTA
β -actin	CTGTCCCTGTATGCCTCTG	ATGTCACGCACGATTTC

500, and 1000 $\mu\text{g}/\text{mL}$) to assess the effects of the extract on cell viability. Based on Figure 4(A) and (B), the SCFE treatment did not induce cytotoxicity in either the 3T3-L1 preadipocytes or mature adipocytes. Furthermore, cell viability consistently exceeded 90 % in all concentrations employed. The non-cytotoxic doses were employed in subsequent experiments to identify SCFE anti-adipogenic attributes.

The effects of the SCFE on lipid accumulation and ROS production during adipocyte differentiation

In the mature adipocyte samples evaluated in this study, intracellular lipid accumulation was assessed via ORO staining to determine SCFE anti-adipogenic activities. Microscopic analysis revealed consistent lipid droplet reductions with increasing SCFE concentration [see Figure 5(A)]. The lipid levels diminished significantly ($p < 0.05$) compared to the control group [see Figure 5(B)], recording 78.64 %, 73.96 %, and 55.34 %, when 250, 500, and 1000 $\mu\text{g}/\text{mL}$ of the SCFE were supplied, respectively. At the highest SCFE concentration, 1000 $\mu\text{g}/\text{mL}$, the lipid reduction (55.34 %) was comparable to the cells treated with NAC, the positive control (45.38 %) ($p > 0.05$).

In this study, the NBT assay was conducted to evaluate the effects of the SCFE on ROS production during adipogenic differentiation. Treatment with 250, 500, and 1000 $\mu\text{g}/\text{mL}$ of the SCFE inhibited ROS production considerably ($p < 0.05$), documenting 59.36 %, 46.40 %, and 33.86 %, respectively, relative to the control [see Figure 5(C)]. At 1000 $\mu\text{g}/\text{mL}$, the SCFE reduced ROS formation by 33.86 %, which was comparable to the NAC (25.26 %) ($p < 0.05$). The data indicated that SCFE inhibitory effects on lipid accumulation and ROS production throughout adipogenesis are dose-dependent.

The effects of the SCFE on TG and leptin levels during adipocyte differentiation

The intracellular TG and leptin contents in the adipocytes assessed in this study were quantified eight days following adipogenic differentiation. The cells treated with the SCFE and positive control (NAC) exhibited notably less TG accumulation than the control samples [see Figure 6(A)]. The cells documented TG levels of 1.79, 1.85, and 1.49 mmol/L post-treatment with 250, 500, and 1000 $\mu\text{g}/\text{mL}$ of the SCFE, respectively, while the NAC-treated cells had 1.50 mmol/L. Meanwhile, the control group recorded 2.33 mmol/L of TG. Nevertheless, the 1000 $\mu\text{g}/\text{mL}$ SCFE treatment yielded no notable variation ($p > 0.05$) compared to the NAC-treated cells regarding TG levels.

The leptin levels of the SCFE-treated cell were significantly lower ($p < 0.05$) than that of the control cells in a dose-dependent manner [see Figure 6(B)]. The cells recorded diminished leptin levels (0.47, 0.40, and 0.16 ng/mL) with increasing SCFE concentrations (250, 500, and 1000 $\mu\text{g}/\text{mL}$). The control group exhibited 0.63 ng/mL leptin content. Treatment with 1000 $\mu\text{g}/\text{mL}$ of the SCFE resulted in a more substantial reduction in leptin levels (0.16 ng/mL) relative to the positive control (NAC) (0.23 ng/mL). The observations suggested that the SCFE effectively reduces TG accumulation and leptin levels during adipogenic differentiation, with the highest concentration (1000 $\mu\text{g}/\text{mL}$) demonstrating comparable effects to the positive control.

The influences of the SCFE on adipogenic transcription factor expressions

Gene expression levels of primary adipogenic transcription factors, including C/EBP- α , PPAR- γ , and SREBP-1,




Figure 3: HPLC chromatograms of (A) the standard compounds and (B) SCFE. Peaks correspond to the following polyphenols present in the SCFE: 1, gallic acid; 2, syringic acid; 3, p-coumaric acid; 4, naringin; 5, cinnamic acid; 6, quercetin; 7, kaempferol; 8, chrysin; 9, galangin; 10, CAPE; 11, pinocembrin.




Figure 4: Effect of SCFE on the viability of (A) 3T3-L1 preadipocytes treated for 24h and (B) 3T3-L1 mature adipocytes treated for 8 days, as determined by MTT assay. The control group represents cells treated with 0.5 % dimethyl sulfoxide (DMSO). Results are shown as the mean \pm standard deviation (SD) of three independent experiments.

were assessed utilizing qRT-PCR. The effects of the SCFE on target genes involved in adipogenesis were determined in the evaluation. The results revealed that the SCFE considerably suppressed C/EBP- α , PPAR- γ , and SREBP-1 gene expressions in 3T3-L1 cells in a dose-dependent manner compared to the control group (see Figure 7). Nevertheless, the inhibitory properties of 500 and 1000 $\mu\text{g/mL}$ SCFE on C/EBP- α and PPAR- γ expressions were similar to the NAC ($p > 0.05$). The findings indicated that the SCFE reduced the expression of adipogenic transcription factors essential for adipocyte differentiation.

Discussion

Adipogenesis, the process where precursor cells differentiate into mature adipocytes, is pivotal in the formation of adipose tissue. Dysregulated adipogenesis, commonly observed in obesity, contributes to metabolic complications, including insulin resistance and inflammation.^{15,16} Consequently, targeting adipogenesis exhibits the potential to address obesity-related disorders.

Natural products, particularly polyphenol-rich substances, have gained attention due to their promising anti-




Figure 5: Effects of SCFE on the buildup of lipids and the generation of reactive oxygen species (ROS) production in 3T3-L1 adipocytes. Post-confluence 3T3-L1 cells were subjected to differentiation either without or with SCFE at concentrations of 250, 500, and 1000 $\mu\text{g/mL}$ for 8 days. (A) Microscopic images of adipocytes stained with Oil red O (Magnification: $40\times$, scale bar: 100 μm). (B) Percentage of lipid accumulation after Oil Red O elution. (C) ROS production was determined by the nitroblue tetrazolium assay. The control group represents cells treated with 0.5 % DMSO, and “NAC” represents cells treated with 5 mM N-acetyl-L-cysteine as a positive control. Results are displayed as the mean \pm SD of three independent experiments. Significant differences ($p < 0.05$) are shown by bars with different letters.




Figure 6: Effects of SCFE on (A) triglyceride and (B) leptin concentrations in 3T3-L1 adipocytes. Post-confluent 3T3-L1 cells were subjected to differentiation either without or with SCFE at concentrations of 250, 500, and 1000 µg/mL for 8 days. The control group represents cells treated with 0.5 % DMSO. “NAC” represents cells treated with N-acetyl-L-cysteine at 5 mM and serves as a positive control group. Results are presented as the mean ± SD of three independent experiments. Significant differences ($p < 0.05$) are shown by bars with different letters.

adipogenic effects.^{17,18} The compounds are abundant in fruits, vegetables, and medicinal plants. Various biological activities, such as anti-inflammatory and antioxidant attributes, have also been reported.^{18,19} Ahmed et al.¹¹ performed chromatographic profiling via HPLC of *S. claviflorum* fruit and seed methanolic extracts. The report revealed substantial phenolic compound levels in the extract. The study identified epicatechin, catechin hydrate, and gallic acid as the predominant compounds in the plant parts. Nonetheless, this study detected a distinct set of compounds; syringic and p-coumaric acids and quercetin, in the ethanolic extract of SCFE. The dissimilarities were attributable to variations in the extraction methods and solvents employed. Differences in geographical origins and maturity of the plants utilized might also contribute to the variations.

The SCFE did not exhibit cytotoxic effects on 3T3-L1 preadipocytes and mature adipocytes at concentrations between 250 and 1000 µg/mL. Consequently, a safe concentration range for subsequent experiments was established. The data also indicated that the effects observed during the differentiation were not confounded by cytotoxicity.

Significant lipid buildup and TG contents in 3T3-L1 cell reductions post-exposure to the SCFE compared to the control cells were revealed during subsequent adipogenic differentiation process assessments. The dose-dependent lipid accumulation inhibition suggested that the SCFE might have impeded preadipocyte maturation into adipocytes, potentially involving mechanisms that disrupt lipid synthesis or storage. TG is a primary component of lipid droplets and the principal storage form of lipids in adipocytes, hence lipid accumulation and TG levels are closely




Figure 7: Effect of SCFE on the regulation of adipogenic gene expression in 3T3-L1 adipocytes. Post-confluent 3T3-L1 cells were subjected to differentiation either without or with SCFE at concentrations of 250, 500, and 1000 $\mu\text{g/mL}$ for 8 days. The expression level of each gene was measured by qRT-PCR and normalized using β -actin as an internal control. The control group represents cells treated with 0.5 % DMSO. NAC represents cells treated with N-acetyl-L-cysteine at 5 mM and serves as a positive control group. Results are presented as the mean \pm SD of three separate experiments. A significant difference ($p < 0.05$) is shown by bars with different letters.

linked.²⁰ The TG levels in the cells also increase as adipocytes mature and accumulate more lipids, which enhances lipid droplets and lipid storage capacity.²¹ In terms of adipogenesis and obesity, increased lipid accumulation and TG content in adipocytes are primary adipocyte hypertrophy and hyperplasia characteristics, contributing to adipose tissue mass expansion.²²

Heightened metabolic activity during adipogenesis generates ROS, which activates Nicotinamide adenine dinucleotide phosphate (NADPH) oxidase, endoplasmic reticulum stress, and inflammation-related immune cells.²³ The reactions also contribute to ROS production during preadipocyte transformation into mature adipocytes. The SCFE treatment employed in the current study substantially reduced ROS production, indicating its antioxidative properties. Polyphenol-rich extracts possess potent antioxidative attributes that could mitigate ROS construction and associated oxidative stress in adipogenesis and obesity. The effectiveness of berry extracts in reducing oxidative stress markers and enhancing antioxidant enzymatic activities in obese individuals has been reported, leading to improved metabolic parameters and decreased adiposity.²⁴ Similarly, green tea extracts, containing catechins, such as epigallocatechin-3-gallate (EGCG), exhibited robust antioxidative effects. EGCG, specifically scavenges ROS, inhibits lipid peroxidation and boosts antioxidant enzyme activity in adipocytes and obese animal models.²⁵

Elevated leptin levels are often associated with obesity due to increased adipose tissue mass.²⁶ The SCFE treatment applied in this study led to reduced leptin levels during 3T3-L1 cell adipogenic differentiation, further supporting the anti-adipogenic effects of the extract. Leptin is an adipokine principally secreted by adipocytes. The substance is critical in regulating energy balance and body weight, which signals satiety to the brain and regulates food intake and energy expenditure.⁵ In adipogenesis, preadipocyte differentiation

into mature adipocytes is accompanied by leptin secretion increment, denoting adipose tissue expansion and lipid store accumulation.⁵ Consequently, interventions that suppress adipogenic differentiation and lipid accumulation in adipocytes might also reduce leptin levels.

At the molecular level, the SCFE treatment demonstrated suppressive effects on the expression of primary adipogenic transcription factors, including C/EBP- α , PPAR- γ , and SREBP-1. The gene expression regulation revealed a potential adipocyte differentiation mechanism through which the SCFE inhibits. The PPAR- γ is a principal adipogenesis modulator that promotes preadipocyte differentiation into mature adipocytes by activating genes for lipid uptake and storage.²⁷ Similarly, C/EBP- α aids adipocyte maturation through lipid synthesis and storage-assisted adipogenic gene expression initiations.²⁸ Conversely, lipid metabolism is managed by SREBP-1, which enhances fatty acid and TG synthesis-related gene transcriptions.²⁹ The SCFE might disrupt the adipogenic gene regulatory network by downregulating C/EBP- α , PPAR- γ , and SREBP-1 expressions, hence impairing adipocyte maturation. The interference could lead to reductions in lipid synthesis and storage within adipocytes, inhibiting adipogenesis.

Conclusion

The current study demonstrated the significant anti-adipogenic effects of the SCFE in impeding 3T3-L1 cell preadipocyte differentiation. The findings indicated promising therapeutic attributes of the SCFE as an intervention for managing obesity and its associated metabolic disorders. The results also advocate the employment of natural extracts in addressing global health issues. Nevertheless, identifying the action mechanisms and active components involved requires further examination.

Source of funding

This research work was funded by the Universiti Teknologi MARA (UiTM) internal research grant-Geran Penyelidikan Centre of Excellence UiTM Cawangan Selangor (DUCS-CoE) [Reference number: 600-UITMSEL (PI. 5/4) (036/2022)].

Conflict of interest

The authors declare no conflict of interest.

Ethical approval

This study did not require ethical approval.

Authors contributions

Conceptualization: RMZ. Data curation: MAA, SMM. Formal analysis: MAA, SMM. Funding acquisition: RMZ. Investigation: MAA, SMM. Supervision: RMZ. Validation: RMZ, AHAI, MZAB. Writing – original draft: RMZ, MAA. Writing, reviewing and editing: RMZ, AHAI, MZAB. All authors have critically reviewed and approved the final draft and are responsible for the content and similarity index of the manuscript.

References

- Yang M, Liu S, Zhang C. The related metabolic diseases and treatments of obesity. *Healthcare* **2022**; 10(9): 1616. <https://doi.org/10.3390/healthcare10091616>.
- Chait A, den Hartigh LJ. Adipose tissue distribution, inflammation and its metabolic consequences, including diabetes and cardiovascular disease. *Frontiers in Cardiovascular Medicine* **2020**; 7(22). <https://doi.org/10.3389/fcvm.2020.00022>.
- Palacios-Marin I, Serra D, Jimenez-Chillarón J, Herrero L, Todorčević M. Adipose tissue dynamics: cellular and lipid turnover in health and disease. *Nutrients* **2023**; 15(18): 3968. <https://doi.org/10.3390/nu15183968>.
- Liu GS, Chan EC, Higuchi M, Dusting GJ, Jiang F. Redox mechanisms in regulation of adipocyte differentiation: beyond a general stress response. *Cells* **2012**; 1(4): 976–993. <https://doi.org/10.3390/cells1040976>.
- Picó C, Palou M, Pomar CA, Rodríguez AM, Palou A. Leptin as a key regulator of the adipose organ. *Rev Endocr Metab Disord* **2022**; 23(1): 13–30. <https://doi.org/10.1007/s11154-021-09687-5>.
- Moseti D, Regassa A, Kim WK. Molecular regulation of adipogenesis and potential anti-adipogenic bioactive molecules. *Int J Mol Sci* **2016**; 17(1): 124. <https://doi.org/10.3390/ijms17010124>.
- Lupianez-Merly C, Dilmaghani S, Vosoughi K, Camilleri M. Review article: pharmacologic management of obesity—updates on approved medications, indications, and risks. *Aliment Pharmacol Ther* **2024**; 59(4): 475–491. <https://doi.org/10.1111/apt.17856>.
- Müller TD, Blüher M, Tschöp MH, DiMarchi RD. Anti-obesity drug discovery: advances and challenges. *Nat Rev Drug Discov* **2022**; 21: 201–223. <https://doi.org/10.1038/s41573-021-00337-8>.
- Singh M, Thrimawithana T, Shukla R, Adhikari B. Managing obesity through natural polyphenols: a review. *Future Foods* **2020**; 1–2. <https://doi.org/10.1016/j.fufo.2020.100002>.
- Craven LA, Biffin E. An infrageneric classification of Syzygium (Myrtaceae). *Blumea-Biodiversity. Evolution and Biogeography of Plants* **2010**; 55: 94–99. <https://doi.org/10.3767/000651910X499303>.
- Ahmed S, Jubair A, Hossain MA, Hossain MM, Azam MS, Biswas M. Free radical-scavenging capacity and HPLC-DAD screening of phenolic compounds from pulp and seed of Syzygium claviflorum fruit. *Journal of Agriculture and Food Research* **2021**; 6:100203. <https://doi.org/10.1016/j.jafr.2021.100203>.
- Kong YT, Abdullah V, Mokhtar SU, Kutty RV. Preliminary studies on extraction of propolis using vitamin E d- α -Tocopheryl polyethylene glycol succinate (vitamin E TPGS) and compare their antimicrobial activities. *Malaysian Journal of Microbiology* **2020**; 16(5): 346–352. <https://doi.org/10.21161/mjm.190516>.
- Choi J, Lee SY, Yoo Y, Kim CH. Maturation of adipocytes is suppressed by fluid shear stress. *Cell Biochem Biophys* **2017**; 75: 87–94. <https://doi.org/10.1007/s12013-016-0771-4>.
- Chung YC, Hyun CG. Inhibitory effects of pinostilbene on adipogenesis in 3T3-L1 adipocytes: a study of possible mechanisms. *Int J Mol Sci* **2021**; 22(24):13446. <https://doi.org/10.3390/ijms222413446>.
- Cordeiro A, Costa R, Andrade N, Silva C, Canabrava N, Pena MJ, et al. Does adipose tissue inflammation drive the development of non-alcoholic fatty liver disease in obesity? *Clinics and Research in Hepatology and Gastroenterology* **2020**; 44(4): 394–402. <https://doi.org/10.1016/j.clinre.2019.10.001>.
- Zatterale F, Longo M, Naderi J, Raciti GA, Desiderio A, Miele C, et al. Chronic adipose tissue inflammation linking obesity to insulin resistance and type 2 diabetes. *Front Physiol* **2020**; 10: 1607. <https://doi.org/10.3389/fphys.2019.01607>.
- Jakab J, Miškić B, Mikšić S, Juranić B, Čosić V, Schwarz D, et al. Adipogenesis as a potential anti-obesity target: a review of pharmacological treatment and natural products. *Diabetes, Metabolic Syndrome and Obesity* **2021**; 14: 67–83. <https://doi.org/10.2147/DMSO.S281186>.
- Kowalska K, Olejnik A, Szwajgier D, Olkiewicz M. Inhibitory activity of chokeberry, bilberry, raspberry and cranberry polyphenol-rich extract towards adipogenesis and oxidative stress in differentiated 3T3-L1 adipose cells. *PLoS One* **2017**; 12(11). <https://doi.org/10.1371/journal.pone.0188583>.
- Martino HSD, dos Santos Dias MM, Noratto G, Talcott S, Mertens-Talcott SU. Anti-lipidaemic and anti-inflammatory effect of açai (*Euterpe oleracea* Martius) polyphenols on 3T3-L1 adipocytes. *J Funct Foods* **2016**; 23: 432–443. <https://doi.org/10.1016/j.jff.2016.02.037>.
- Kislev N, Eidelheit S, Perlmutter S, Benayahu D. How to follow lipid droplets dynamics during adipocyte metabolism. *J Cell Physiol* **2022**; 237(11): 4157–4168. <https://doi.org/10.1002/jcp.30857>.
- Li Q, Spalding KL. The regulation of adipocyte growth in white adipose tissue. *Front Cell Dev Biol* **2022**; 10. <https://doi.org/10.3389/fcell.2022.1003219>.
- Lafontan M. Adipose tissue and adipocyte dysregulation. *Diabetes Metabol* **2014**; 40(1): 16–28. <https://doi.org/10.1016/j.diabet.2013.08.002>.
- Masenga SK, Kabwe LS, Chakulya M, Kirabo A. Mechanisms of oxidative stress in metabolic syndrome. *Int J Mol Sci* **2023**; 24(9): 7898. <https://doi.org/10.3390/ijms24097898>.
- Stote KS, Burns G, Mears K, Sweeney M, Blanton C. The effect of berry consumption on oxidative stress biomarkers: a systematic review of randomized controlled trials in humans. *Antioxidants* **2023**; 12(7): 1443. <https://doi.org/10.3390/antiox12071443>.
- Kim HS, Quon MJ, Kim JA. New insights into the mechanisms of polyphenols beyond antioxidant properties; lessons from the green tea polyphenol, epigallocatechin 3-gallate. *Redox Biol* **2014**; 2: 187–195. <https://doi.org/10.1016/j.redox.2013.12.022>.

26. Stern JH, Rutkowski JM, Scherer PE. Adiponectin, leptin, and fatty acids in the maintenance of metabolic homeostasis through adipose tissue crosstalk. *Cell Metabol* **2016**; 23(5): 770–784. <https://doi.org/10.1016/j.cmet.2016.04.011>.
27. Sun C, Mao S, Chen S, Zhang W, Liu C. PPARs-orchestrated metabolic homeostasis in the adipose tissue. *Int J Mol Sci* **2021**; 22(16): 8974. <https://doi.org/10.3390/ijms22168974>.
28. Rosen ED, Hsu CH, Wang X, Sakai S, Freeman MW, Gonzalez FJ, et al. C/EB α induces adipogenesis through PPAR γ : a unified pathway. *Genes Dev* **2002**; 16(1): 22–26. <https://doi.org/10.1101/gad.948702>.
29. Shimano H, Sato R. SREBP-regulated lipid metabolism: convergent physiology-divergent pathophysiology. *Nat Rev Endocrinol* **2017**; 13(12): 710–730. <https://doi.org/10.1038/nrendo.2017.91>.

How to cite this article: Zohdi RM, Adli MA, Mukhtar SM, Awang Junaidi AH, Bakar MZA. *Syzygium claviflorum* fruit extract preadipocyte differentiation inhibition in 3T3-L1 cells. *J Taibah Univ Med Sc* **2024**;19(6):1181–1192.