



UNIVERSITI PUTRA MALAYSIA

**MOLECULAR CHARACTERISATION, PATHOGENICITY AND
IMMUNOLOGICAL STUDIES OF CHICKEN ANAEMIA VIRUS
ISOLATED IN MALAYSIA**

SHAH MD. ZIQRUL HAQ CHOWDHURY

FPV 2001 11

**MOLECULAR CHARACTERISATION, PATHOGENICITY AND
IMMUNOLOGICAL STUDIES OF CHICKEN ANAEMIA VIRUS ISOLATED IN
MALAYSIA**

SHAH MD. ZIQRUL HAQ CHOWDHURY

**DOCTOR OF PHILOSOPHY
UNIVERSITI PUTRA MALAYSIA**

2001



**MOLECULAR CHARACTERISATION, PATHOGENICITY AND
IMMUNOLOGICAL STUDIES OF CHICKEN ANAEMIA VIRUS ISOLATED IN
MALAYSIA**

By

SHAH MD. ZIQRUL HAQ CHOWDHURY

**Thesis Submitted in Fulfilment of the Requirement for the
Degree of Doctor of Philosophy in the
Faculty of Veterinary Medicine
Universiti Putra Malaysia**

October 2001



DEDICATED TO

**My Parents, *SHAH MD. LUTFAR RAHMAN CHOWDHURY*
*AND MRS RABEYA KHATUN CHOWDHURY***

**My Wife, *FAUZIA YASMIN CHOWDHURY* and
My Daughter, *FARZANA YASMIN CHOWDHURY***

My Five Brothers, Three Sisters and One Late Sister



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Doctor of Philosophy

**MOLECULAR CHARACTERISATION, PATHOGENICITY AND
IMMUNOLOGICAL STUDIES OF CHICKEN ANAEMIA VIRUS ISOLATED IN
MALAYSIA**

By

SHAH MD. ZIQRUL HAQ CHOWDHURY

October 2001

Chairman: Professor Dr. Aini Ideris

Faculty: Veterinary Medicine

A comprehensive study was carried out to isolate, identify and characterise chicken anaemia virus (CAV) isolated in Malaysia. The study resulted in the isolation of five CAV isolates from broiler chickens, designated as BL-1, BL-2, BL-3, BL-4 and BL-5. These isolates together with three isolates (SMSC-1, SMSC-2 and 3-1) provided by Veterinary Research Institute (VRI), Malaysia and a reference Cux-1 isolate were analysed by different restriction endonuclease enzymes. The whole genome of each CAV isolate was amplified by PCR into four fragments: Fragments A, B, C and D. Fragment A was digested with *StyI*, fragment B with *StyI*, *HpaII* and *MboI*, fragment C with *HaeIII*, and fragment D with *EcoRI*. The overall analysis revealed that the four isolates, BL-1, BL-2, BL-4 and BL-5, exhibited the same restriction profiles in all enzymatic reactions and they are placed in one group, whereas, the other five



isolates (SMSC-1, SMSC-2, 3-1, BL-3 and Cux-1) were found to be different from each other and also from the group of four isolates mentioned above.

The pathogenicity studies in specific pathogen free (SPF) chickens inoculated with SMSC-1, 3-1 and BL-5 isolates at 1-day old showed that, the isolates produced clinical signs and characteristic lesions suggestive of CAV infection at 14-16 days post inoculation (p.i.). The histopathological lesions in infected chicks showed severe depletion of lymphocytes from thymus, bursa and spleen and aplastic changes in bone marrow. The repeated passages of two VRI isolates, SMSC-1 and 3-1, in MSB1 cell line until passage sixty (P60), and passage 123 (P123), produced attenuated viruses (SMSC-1/P60, 3-1/P60, SMSC-1/P123 and 3-1/P123) which showed significantly reduced level of pathogenicity in SPF chickens compared to the pathogenic parent isolates.

The whole genome of two non-attenuated isolates (SMSC-1 and 3-1) and two attenuated isolates (SMSC-1/P60 and 3-1/P60) were sequenced using the Perkin Elmer's BigDye Terminator Cycle Sequencing Kit. The high G:C regions of the CAV genome were sequenced using the same kit by the development of a modified method. The results showed that the complete genome of all isolates consisted of 2298 nucleotides. Three major ORFs of 1347 bp, 648 bp and 363 bp long were found in the plus DNA strand in all isolates, coding for putative proteins of about 52 kDa (VP1), 24 kDa (VP2) and 13 kDa (VP3), respectively. The alignment and antigenic index of VP1

sequence revealed the appearance of a hypervariable region from amino acid positions 139 to 157. The results showed that 76 nucleotide changes in SMSC-1/P60 and 43 nucleotide changes in 3-1/P60 isolates compared to their parent isolates, were thought to contribute to virus attenuation. Among these nucleotide changes, only one nucleotide difference (T→C) at position 816 resulted in changes of amino acid residues at positions 153 in VP2 from V to A, and 118 in VP3 from C to R. This single nucleotide change is probably important for the change in virus pathogenicity or attenuation. The phylogenetic analysis showed that the SMSC-1 isolate is close to the Australian 704 and Japanese TR20, the 3-1 isolate is close to the German Cux-1 isolate and the attenuated cloned isolate 10 (derived from the Cux-1). The attenuated SMSC-1/P60 and 3-1/P60 isolates were very close to the Japanese isolate A2.

The apoptosis study carried out with electron microscopy and DNA fragmentation analysis, detected apoptosis both in infected thymocytes and infected MSB1 cells. The immunological studies with P1, P60 and P123 isolates of SMSC-1 and 3-1, and also with BL-5 isolate, after inoculation into 1-day-old SPF chickens showed that each of the isolates elicited CAV antibody responses, both at 14-16 days and 30 days p.i. Based on the findings of antibody response and pathogenicity studies, the attenuated isolates of P60 and P123 are potential candidates for live vaccines.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**PENCIRIAN MOLEKUL, KAJIAN KEPATOGENAN DAN IMUNOLOGI VIRUS
ANEMIA AYAM YANG DIPENCILKAN DI MALAYSIA**

Oleh

SHAH MD. ZIQRUL HAQ CHOWDHURY

Oktober 2001

Pengerusi: Profesor Dr. Aini Ideris

Fakulti: Perubatan Veterinar

ABSTRAK

Satu kajian yang komprehensif telah dijalankan untuk memencil, mengenalpasti dan mencirikan virus anemia ayam (CAV) yang di isolat di Malaysia. Kajian ini menghasilkan pemencilan lima isolat CAV daripada ayam pedaging, dinamakan sebagai BL-1, BL-2, BL-3, BL-4 dan BL-5. Isolat-isolat tersebut bersama tiga isolat (SMSC-1, SMSC-2 dan 3-1) diberikan oleh Institut Penyelidikan Veterinar (VRI), Malaysia, dan isolat rujukan Cux-1 telah dianalisis melalui enzim endonukleas pembatas. Keseluruhan genom bagi setiap isolat CAV diampifikasi melalui PCR kepada empat fragmen: Fragmen A, B, C dan D. Fragmen A telah dipotong dengan *StyI*, Fragmen B dengan *StyI*, *HpaII* dan *MboI*, Fragmen C dengan *HaeIII* dan fragmen D dengan *EcoRI*. Analisis keseluruhan menunjukkan bahawa empat isolat, BL-1, BL-2, BL-4 dan BL-5 menghasilkan profil pembatas yang sama di dalam kesemua tindak balas enzim dan isolat berkenaan diletakkan dalam satu kumpulan, manakala lima isolat



(SMSC-1, SMSC-2, 3-1, BL-3 dan Cux-1) didapati berbeza antara satu sama lain dan juga daripada kumpulan empat isolat yang disebut di atas.

Kajian kepatogenan ke atas ayam bebas patogen spesifik (SPF) yang diinokulat dengan SMSC-1, 3-1 dan BL-5 menunjukkan tanda-tanda klinikal dan ciri-ciri lesi CAV. Lesi histopatologi dalam ayam terjangkit menunjukkan pengurangan limfosit daripada timus, bursa dan limpa dan perubahan aplastik dalam sum-sum tulang. Pengulangan inokulasi bagi dua isolat VRI, SMSC-1 dan 3-1, dalam sel MSBI hingga ke inokulasi 60 (P60) dan inokulasi 123 (P123) menghasilkan virus atenuat (SMSC-1/P60, 3-1/P60, SMSC-1/P123 dan 3-1/P123) yang mana menunjukkan pengurangan tahap kepatogenan yang signifikan dalam ayam SPF berbanding dengan isolat patogenik asal.

Keseluruhan genom bagi dua isolat yang tidak diatenuat (SMSC-1 dan 3-1) dan dua isolat yang diatenuat (SMSC-1/P60 dan 3-1/P60) telah diujukkan menggunakan Perkin Elmer's BigDye Terminator Cyclor Sequencing Kit. Kawasan G:C yang tinggi bagi genom CAV diujukkan dengan menggunakan kit sama dengan ubahsuaian. Keputusan menunjukkan bahawa genom yang lengkap bagi kesemua isolat terdiri daripada 2298 nukleotid. Tiga ORF major daripada 1347 bp, 648 bp dan 363 bp telah dijumpai pada bebenang DNA tambahan dalam kesemua isolat, dengan mengekod protein putatif pada anggaran 52kDa (VP1), 24 kDa (VP2) dan 13 kDa (VP3), masing-masing. Jujukan VP1 yang disusun memperlihatkan kemunculan satu kawasan

hiper boleh-ubah daripada asid amino berkedudukan 139 hingga 157. Indeks antigenik VP1 juga menunjukkan kawasan hiper boleh-ubah di antara isolat-isolat, dalam kawasan asid amino 122 hingga 165. Keputusan tersebut menunjukkan bahawa 76 nukleotid terubah dalam SMSC-1/P60 dan 43 nukleotid terubah dalam 3-1/P60 isolat berbanding dengan isolat asal. Perubahan ini dianggapkan menyumbang kepada pengaknuatan virus. Di kalangan perubahan nukleotid ini, hanya perbezaan nukleotid (T→C) pada kedudukan 816, yang menghasilkan perubahan dalam asid amino residu pada kedudukan 153 dalam VP2 daripada V kepada A, dan 118 dalam VP3 daripada C kepada R. Perubahan satu nukleotid adalah penting untuk merubah kepatogenan atau pengaknuatan virus. Analisis filogenetik menunjukkan bahawa isolat SMSC-1 adalah hampir kepada isolat Australia 704 dan Jepun TR20, isolat 3-1 adalah hampir kepada isolat Jerman Cux-1 dan klon isolat 10 yang diatenuat. Isolat SMSC-1/P60 dan 3-1/P60 yang diatenuat adalah sangat hampir kepada isolat Jepun A2.

Kajian apoptosis dengan menggunakan mikroskop elektron dan analisis serpihan DNA, telah mengesan apoptosis dalam timosit terjangkit dan sel MSBI terjangkit. Kajian imunologi ke atas P1, P60 dan P123 bagi isolat SMSC-1 dan 3-1 dan juga dengan isolat BL-5, menunjukkan bahawa setiap isolat menghasilkan tindakbalas antibodi CAV. Berdasarkan penemuan tindakbalas antibodi dan kajian kepatogenan, isolat-isolat yang dilemahkan bagi P60 dan P123 mempunyai potensi sebagai calon vaksin hidup.

ACKNOWLEDGEMENTS

Bismillahir Rahmanir Rahim

All praises and credits are due to Almighty Allah who has given me the ability and strength to complete this challenging task.

I would like to express my deepest sense of appreciation and gratitude to Professor Dr. Aini Ideris, Dean of Graduate School, UPM and Chairman of Supervisory Committee, for her invaluable guidance, advice and suggestions throughout the study. I am also grateful to her for sending me to UKM, to complete two short training courses on molecular works.

My sincere gratitude and appreciation is due to Dr. Abdul Rahman Omar, a member of the supervisory committee for his continuous guidance, suggestions and cooperation throughout the study.

I am expressing my sincere thanks and gratitude to Assoc. Prof. Dr. Mohd. Hair Bejo, a member of the supervisory committee for his guidance, suggestions and encouragement throughout the study.

My sincere appreciation and gratitude is due to Dr. Abdul Aziz Jamaluddin, Director, Veterinary Research Institute (VRI), Ipoh, Perak, Malaysia, and a member of the supervisory committee, for his invaluable guidance, suggestions and cooperation throughout the study. I am really



grateful to him for arranging a training programme for me in VRI, for the isolation and identification of chicken anaemia virus (CAV) from field samples. My cordial thanks are also due to him for providing three VRI CAV isolates for this study.

I would like to express my sincere gratefulness to Dr. Y. Kono, a JICA Specialist at VRI, for his invaluable technical training, suggestions and cooperation during the study. I would like to appreciate the contribution of Dr. Nadzri Salim, Lecturer, Faculty of Veterinary Medicine, UPM, for helping me to analyse my research data statistically.

I would like to express my sincere thank to Assoc. Prof. Dr. Rahmah Mohamed and Dr. Wan Kiew Lian of UKM, Dr. Tan Siang Hee and Dr. Hairkrishna of Genetic Lab., UPM for providing the facilities for sequencing.

My sincere thanks are due to Dr. Badrul Munir Md. Zain of UKM, for his help and cooperation during phylogenetic analysis of the CAV sequences. Thanks are also due to Mr. Mohd. Noor Mat Isa, MTDC, UKM, and Mr. Lee, Genetic Lab., UPM, for their help and cooperation during sequence analysis.

I really appreciate the help and cooperation of Dr. Azizah Darus of VRI, Perak, Malaysia, and Dr. Haas Md Yatim of MTDC, Malaysia, for their help, cooperation and valuable suggestions during collection of CAV samples from

different broiler farms. Thanks are due to Dr. Reuben Sharma of Parasitology Lab., Ms Siti Hasmah Mohtar, Kong Lih Ling and Tan Sheau Wei of Biologics Lab., UPM, for their help and cooperation during the study. I would also like to thank the staffs at Biologics Lab., Histopathology Lab., Haematology Lab., Electron microscopy unit, Genetic Lab. at UPM, and the staffs at VRI, Perak, Malaysia for their help, cooperation and suggestions during the study.

I would like to express my sincere thanks to the World Bank for providing financial support (IDA Credit 2815-BD) for the scholarship under ARMP-BLRI part. I am also grateful to the Government of Bangladesh and Government of Malaysia for providing other financial support to ensure completion of my Ph.D. studies.

My cordial thanks are to the Ministry of Livestock and Fisheries, Bangladesh Secretariate, Dhaka, and Director General, Bangladesh Livestock Research Institute, Savar, Dhaka, Bangladesh, for their continuous help during the study, extension of my scholarship and deputation.

My heartfelt appreciation to my parents, brothers and sisters for their help and encouragement during the study. Finally, I am expressing my utmost gratitude and appreciation to my wife, Fauzia Yasmin Chowdhury and my only daughter, Farzana Yasmin Chowdhury, for their patience, continuous help, cooperation and encouragement throughout the study period.



TABLE OF CONTENTS

	Page
DEDICATION -----	ii
ABSTRACT -----	iii
ABSTRAK -----	vi
ACKNOWLEDGEMENTS -----	ix
APPROVAL -----	xii
APPROVAL -----	xiii
DECLARATION -----	xiv
LIST OF TABLES -----	xxi
LIST OF FIGURES -----	xxiv
LIST OF PLATES -----	xxvi
LIST OF ABBREVIATIONS -----	xxx
CHAPTER	
I GENERAL INTRODUCTION -----	1
II REVIEW OF LITERATURE -----	8
Chicken Anaemia Virus -----	8
Classification -----	10
History -----	10
Virus Properties -----	11
Molecular Biology of CAV -----	13
Viral Proteins -----	19
Restriction Endonuclease Enzyme Analysis -----	21
Incidence and Distribution -----	23
Pathogenesis, Pathogenicity and Antigenicity -----	25
Disease Signs -----	30
Naturally Occurring Disease -----	30
Experimental Disease -----	32
Morbidity and Mortality -----	34
Gross Lesions -----	35
Histopathology -----	36
Bonemarrow -----	37
Thymus -----	40
Bursa of Fabricius -----	42
Spleen -----	42
Liver -----	43
Other Organs -----	43
Heamatology -----	44



Apoptosis -----	46
Virus Attenuation and Molecular Basis for Attenuation -----	51
Immunity against CAV -----	56
Active Immunity -----	56
Passive Immunity -----	58
Immunosuppression -----	59
Economic Effect -----	61
Diagnosis -----	62
Isolation and Identification of the Virus -----	62
Differential Diagnosis -----	68
CAV Vaccine, Prevention and Control -----	69
III ISOLATION AND IDENTIFICATION OF MALAYSIAN ISOLATES OF CHICKEN ANAEMIA VIRUS -----	75
Introduction -----	75
Materials and Methods -----	77
Viruses -----	77
Cells and Cell Culture -----	78
Chickens -----	78
Sample Collection -----	78
Preparation of Virus Inoculum -----	79
Isolation of Virus in MDCC-MSB1 Cells -----	79
DNA Extraction from Samples -----	80
DNA Extraction from Infected Culture Cells -----	80
DNA Quantification and Purity -----	81
Polymerase Chain Reaction (PCR) -----	81
Agarose Gel Electrophoresis -----	83
Purification of DNA -----	83
Preparation of Stock from Newly Isolated Virus -----	83
Indirect Immunofluorescence Assay (IIFA) -----	84
Treatment with Restriction Endonuclease Enzyme -----	85
Treatment with Chloroform and Heat -----	85
Chicken Inoculation -----	86
Results -----	86
Virus Isolation -----	86
Detection of CAV DNA in Samples and in Infected MSB1 Cells by PCR -----	87
Detection of CAV Antigens in MSB1 Cells by IIFA -----	87
Restriction Enzyme Analysis -----	88
Treatment with Chloroform and Heat -----	88
Chicken Inoculation -----	88
Discussion -----	92
IV ANALYSIS OF THE GENOMES OF DIFFERENT ISOLATES OF CHICKEN ANAEMIA VIRUS BY RESTRICTION ENDONUCLEASE ENZYMES -----	95

Introduction -----	95
Materials and Methods -----	97
Viruses -----	97
Cells and Cell Culture -----	97
Virus Stock Preparation -----	97
Virus Inoculation into MSB1 Cells -----	98
DNA Extraction and Precipitation -----	98
Amplification of DNA Fragments by Polymerase Chain Reaction (PCR) -----	98
PCR Reaction Mixture and Thermal Incubations -----	100
Agarose Gel Electrophoresis and Purification of DNA Fragments -----	100
Analysis of Amplified DNA Fragments by Restriction Endonucleases -----	102
Agarose Gel Electrophoresis -----	102
Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis (SDS-PAGE) -----	104
Results -----	106
PCR Amplification of Different Fragments -----	106
Restriction Endonuclease Analysis -----	107
Discussion -----	118

V	MOLECULAR CLONING OF CHICKEN ANAEMIA VIRUS GENOME AMPLIFIED BY POLYMERASE CHAIN REACTION ----	124
	Introduction -----	124
	Materials and Methods -----	125
	Viruses -----	125
	Cells and Cell Culture -----	126
	Virus Inoculation into MSB1 Cells -----	126
	DNA Extraction and Precipitation -----	126
	Determination of DNA Concentration and Purity -----	126
	Amplification of CAV DNA by Polymerase Chain Reaction (PCR) -----	126
	PCR Reaction Mixture and Thermal Conditions -----	127
	Agarose Gel Electrophoresis -----	127
	Purification of PCR Products from Agarose Gel -----	127
	Cloning -----	127
	Transformation Efficiency -----	132
	Analysis of Positive Clones by PCR Screening -----	133
	Preparation of Glycerol Stock -----	133
	Extraction of Plasmid DNA by Conventional Method -----	134
	Extraction of Plasmid DNA by Commercial Kit -----	135
	Determination of Concentration and Purity of Plasmid DNA ----	136
	Analysis of Positive Clones by Restriction Endonuclease (RE) Analysis -----	137
	Results -----	137



	Amplification of CAV DNA by PCR -----	137
	Cloning -----	139
	Transformation Efficiency -----	139
	Plasmid DNA Concentration and Purity -----	142
	Analysis of Positive Clones -----	142
	Discussion -----	151
VI	DNA SEQUENCING AND PHYLOGENETIC ANALYSIS OF THE GENOME OF NON-ATTENUATED AND ATTENUATED ISOLATES OF CHICKEN ANAEMIA VIRUS -----	154
	Introduction -----	154
	Materials and Methods -----	156
	Viruses -----	156
	Construction of CAV Recombinant Plasmid -----	156
	Extraction of Plasmid -----	156
	DNA Sequencing -----	157
	The Sequencing Reaction Mixture -----	157
	Purification of Extension Products -----	158
	Electrophoresis on the ABI Prism [®] 377 DNA Sequencer -----	169
	Sequencing High G:C Regions in CAV Genome -----	162
	Sequencing the Isolate SMSC-1/P 123 -----	164
	Sequence Analysis -----	165
	GenBank Accession Numbers for Nucleotide Sequences -----	166
	CAV Sequences Collected from GenBank -----	166
	Hydrophilicity and Antigenicity of VP1 -----	167
	Phylogenetic Analysis -----	167
	Results -----	171
	Development of a Modified Method for Sequencing High G:C Rich Regions -----	171
	Sequence Analysis -----	173
	Nucleotide Sequence Motifs and Regulation of Transcription -	191
	Sequence Alignment -----	198
	Sequence of CAV Isolate SMSC-1/P 123 -----	199
	Hydrophilicity and Antigenicity of the VP1 Protein -----	199
	Phylogenetic Analysis -----	203
	Discussion -----	211
VII	PATHOGENICITY OF NON-ATTENUATED ISOLATES OF CHICKEN ANAEMIA VIRUS IN SPECIFIC PATHOGEN FREE CHICKENS -----	222
	Introduction -----	222
	Materials and Methods -----	224
	Viruses -----	224
	Cells and Cell Culture -----	224
	Chickens -----	224
	Virus Stock Preparation -----	225



	Virus Titration with MDCC-MSB1 Cells -----	225
	Experimental Design -----	226
	Determination of Haematocrit (Packed Cell Volume- PCV) ----	228
	Postmortem Examination -----	228
	Organ Weight Calculation -----	229
	Statistical Analysis -----	229
	Histopathology -----	230
	Results -----	230
	Virus Titration -----	230
	Clinical Signs -----	231
	Body Weight and Organ Weight -----	232
	Haematocrit Values and Anaemia -----	234
	Lesion Score -----	236
	Histopathology -----	237
	Discussion -----	253
VIII	ATTENUATION OF CHICKEN ANAEMIA VIRUS BY REPEATED PASSAGE IN MDCC-MSB1 CELL LINE FOR DEVELOPMENT OF LIVE ATTENUATED VACCINE -----	257
	Introduction -----	257
	Materials and Methods -----	260
	Viruses -----	260
	Cells and Cell Culture -----	260
	Chickens -----	260
	Counting of MSB1 Cells -----	261
	Virus Attenuation by Repeated Passage in MSB1 Cells -----	261
	CPE Scoring -----	262
	Virus Stock Preparation -----	262
	Virus Titration -----	262
	Indirect Immunofluorescence Assay (IIFA) -----	263
	Chicken Inoculation and Experimental Design -----	263
	Determination of Haematocrit Values -----	264
	Postmortem Examination -----	265
	Organ Weight Calculation -----	265
	Statistical Analysis -----	265
	Results -----	265
	Virus Attenuation by Repeated Passage in MSB1 Cells -----	265
	Titration of CAV Isolates -----	268
	Indirect Immunofluorescence Assay (IIFA) -----	269
	Chicken Inoculation -----	271
	Discussion -----	276
IX	APOPTOSIS AND IMMUNOLOGICAL STUDIES ON MALAYSIAN ISOLATES OF CHICKEN ANAEMIA VIRUS -----	282
	Introduction -----	282
	Materials and Methods -----	284



Viruses -----	284
Chickens -----	284
Cells and Cell Culture -----	285
Apoptosis Study -----	285
Transmission Electron Microscopy (TEM) -----	286
DNA Fragmentation Analysis-----	289
Immunological Study -----	290
Enzyme-linked Immunosorbent Assay (ELISA) -----	291
Statistical Analysis -----	297
RESULTS -----	297
Electron Microscopy (TEM) of Thymuses after <i>in vivo</i> Infection -----	297
Electron Microscopy (TEM) of MDCC-MSB1 Cell line after <i>in</i> <i>vitro</i> Infection -----	298
DNA Fragmentation Analysis of Thymuses and MDCC-MSB1 cell line -----	298
Antibody Development by CAV SMSC-1/P1, SMSC-1/P60 and SMSC-1/P123 -----	305
Antibody Development by CAV 3-1/P1, 3-1/P60 and 3-1/P123	305
Antibody Development by CAV BL-5 isolate	306
DISCUSSION -----	309
X GENERAL DISCUSSION AND CONCLUSION -----	315
Future Prospects and Suggestions -----	324
BIBLIOGRAPHY -----	327
APPENDICES -----	351
BIODATA OF AUTHOR -----	393



LIST OF TABLES

Table		Page
Table 2.1:	The economic impact of CAV on commercial broiler production -----	62
Table 4.1:	Restriction endonuclease analysis of different PCR-amplified DNAs specified by different CAV isolates -----	108
Table 4.2:	Number of DNA fragments produced in different PCR-amplified genomic fragments of different Isolates after digestion with various restriction endonucleases -----	109
Table 5.1:	Formation of colonies after transformation of the recombinant plasmids into Top10 <i>E. coli</i> cells -----	141
Table 5.2:	Transformation efficiency of the recombinant plasmid pCR [®] 2.1 with insert fragment A or fragment B of different CAV isolates into chemically competent Top10 cells -----	141
Table 6.1:	List of primers for sequencing fragments A and B of the CAV genomes from different isolates -----	168
Table 6.2:	Sequence of the forward primers used for sequencing the CAV genome from different isolates -----	169
Table 6.3:	Sequence of the reverse primers used for sequencing the CAV genome from different isolates -----	170
Table 6.4:	Molecular weights of the putative proteins encoded by the three major ORFs of 1347 bp, 648 bp and 363 bp in the plus DNA strand of different isolates -----	176
Table 6.5:	Percentage homologies, differences of nucleotide and amino acid sequences between SMSC-1 isolate and other CAV isolates -----	179
Table 6.6:	Percentage homologies, differences of nucleotide and protein sequences between 3-1 isolate and other CAV isolates -----	184
Table 6.7:	Percentage homologies, differences of nucleotide and protein sequences between SMSC-1/P60 isolate and other	



	CAV isolates -----	186
Table 6.8:	Percentage homologies, differences of nucleotide and protein sequences between 3-1/P60 isolate and other CAV isolates -----	190
Table 6.9:	Nucleotide sequence motifs in SMSC-1 isolate -----	194
Table 6.10:	Nucleotide sequence motifs in 3-1 isolate -----	195
Table 6.11:	Nucleotide sequence motifs in SMSC-1/P60 isolate -----	196
Table 6.12:	Nucleotide sequence motifs in 3-1P60 isolate -----	197
Table 6.13:	Summary of variations along the sequences across CAV isolates -----	204
Table 6.14:	Tajima and Nei Distance (x100) Matrix derived from different CAV sequences -----	210
Table 7.1:	Body weights and organ weights of SPF chickens 14 days following infection with SMSC-1 isolate at 1 day of age ----	233
Table 7.2:	Body weights and organ weights of SPF chickens 16 days following infection with 3-1 and BL-5 isolates at 1 day of age -----	233
Table 7.3:	Pathogenicity evaluation in SPF chickens 14 days after inoculation with SMSC-1 isolate at 1 day of age -----	235
Table 7.4:	Pathogenicity evaluation of SPF chickens 16 days after inoculation with 3-1 and BL-5 isolates at 1 day of age -----	236
Table 8.1:	Body weights and organ weights of SPF chickens 16 days following infection with SMSC-1/P60 and 3-1/P60 isolates at 1-day of age -----	274
Table 8.2:	Pathogenicity evaluation of SMSC-1/P60 and 3-1/P60 isolates 16 days following infection in 1 day old SPF chickens -----	275
Table 8.3:	Body weights and organ weights of SPF chickens 16 days following infection with SMSC-1/P123 and 3-1/P123 isolates at 1-day of age -----	275
Table 8.4:	Pathogenicity evaluation of SMSC-1/P123 and 3-1/P123	



	isolates 16 days following infection in 1 day old SPF chickens -----	276
Table 9.1:	Serum antibody titres of chickens following infection with SMSC-1/P1, SMSC-1/P60 and SMSC-1/P123 isolates at 1-day of age -----	307
Table 9.2:	Serum antibody titres of chickens following infection with 3-1/P1, 3-1/P60 and 3-1/P123 isolates at 1-day of age -----	308
Table 9.3:	Serum antibody titres of chickens following infection with BL-5 isolate at 1-day of age -----	308

