



UNIVERSITI PUTRA MALAYSIA

**PHENOTYPIC AND MOLECULAR CHARACTERIZATION OF
PASTEURELLA MULTOCIDA OBTAINED FROM POULTRY IN IRAN**

AHMAD REZA JABBARI

FPV 2001 10

**PHENOTYPIC AND MOLECULAR CHARACTERIZATION OF
PASTEURELLA MULTOCIDA OBTAINED FROM POULTRY IN IRAN**

By

AHMAD REZA JABBARI ,

**Thesis Submitted in Fulfilment of the Requirement for the Degree of
Doctor of Philosophy in the Faculty of Veterinary Medicine
Universiti Putra Malaysia**

September 2001



**DEDICATED TO MY WIFE, MAHTAB MOZAFFARI,
OUR TWO SONS, MOHAMMAD VAHID
AND MOHAMMAD MOEIN**



Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

PHENOTYPIC AND MOLECULAR CHARACTERIZATION OF PASTEURELLA MULTOCIDA OBTAINED FROM POULTRY IN IRAN

By

AHMAD REZA JABBARI

September 2001

Chairman: Associate Professor Dr. Abdul Aziz Saharee

Faculty: Veterinary Medicine

A collection of twenty five *Pasteurella multocida* isolates obtained from avian pasteurellosis in northern part (endemic area) of Iran were studied for some of their phenotypic and molecular characteristics. This research is the first study on conducting of serotyping and molecular characterization of avian *P. multocida* in Iran.

Based on the biochemical characteristics, all *P. multocida* isolates tested belonged to subspecies (biotype) *multocida*. Antimicrobial sensitivity test showed that all the isolates examined were resistant to at least three of the thirteen antimicrobials tested. Among the antimicrobial agents, chloramphenicol, combination of sulfametoxazin and trimetoprim and nitrofurantoin were found to be the most effective (100% sensitivity)



followed by tetracycline (96% sensitivity), penicillin (88% sensitivity) and gentamycin (76% sensitivity). The highest percentage of resistance was found against lincomycin, bacitracin and cloxacillin (100% resistant) followed by furazolidone and colistin (84% and 68 % respectively).

Agar gel diffusion precipitation (AGDP) test was used to determine somatic serotypes of the isolates. According to the results of the AGDP test, Serotype 1 was dominant among avian isolates from endemic area. Serotypes 3, 3 × 4 and 4, found for the first time in the country were also identified among the isolates.

Electrophoresis protein patterns of the isolates were studied by using sodium dodecyl sulphate polyacrylamide gel (SDS-PAGE). All strains were similar in the majority of protein bands. The main difference between protein patterns of the isolates was revealed in the position of one of the major band (H Protein) presented in the 34- 38 KDa region. According to H protein position, three distinguishable groups were identified. In protein type I, the molecular mass of H protein was about 38 KDa but in protein types II and III it was 36.5 and 34 KDa respectively.

The minimum lethal dose (MLD) of the strains with protein types I, II and III as a virulence determinant was identified in mice. It was revealed that the strain with protein type III had the least virulence and the strain with protein

type I had the greatest virulence in mice. Immunization of mice with strain PMI030 (protein type I) induced a good protection against homologous protein type challenge.

Restriction enzyme analysis (REA) of chromosomal DNA and repetitive extragenic palindromic elements PCR (REP-PCR) were used for determination of genetic diversity among the isolates. DNA fingerprinting by HpaII digestion divided the twenty-five isolates into 7 REA groups, 2 of which contained a single isolate.

DNA fingerprinting with REP-PCR revealed a great genetic diversity among the isolates. According to amplified DNA patterns, a total of 9 REP-PCR groups were determined. REP-PCR produced amplified bands ranging in size from approximately 700 bp to 3.6 Kb with two species-specific bands of 0.8 Kb and 2.3 Kb. REP-PCR was able to differentiate *P.multocida* isolates from different source and geographical areas. Results of this study showed that the use of REP element amplification by polymerase chain reaction is highly reproducible and can be suggested as a suitable epidemiologic tool especially for investigation on the origin of outbreaks and similarity between different avian isolates of *P.multocida*.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

CIRI-CIRI FENOTIPIK DAN MOLEKULAR *PASTEURELA MULTOCIDA* YANG DIPEROLEHI DARIPADA TERNAKAN AYAM ITIK DI IRAN

Oleh

AHMAD REZA JABBARI

September 2001

Pengerusi: Profesor Madya Dr. Abdul Aziz Saharee

Fakulti Perubatan Veterinar

Satu kajian tentang sekumpulan 25 *Pasteurella multocida* yang diperolehi daripada pasteurelosis avian di bahagian utara (kawasan endemi) Iran telah dijalankan untuk meneliti ciri-ciri fenotipik dan molekularnya. Kajian ini merupakan yang pertama mengkaji tentang kelakuan serotip dan ciri-ciri molekul *P.multocida* avian di Iran.

Berasaskan ciri-ciri biokimia, kesemua pemencilan *P.multocida* yang diuji tergolong sebagai subspecies (biotaip) multocida. Ujian sensitiviti antimikrob menunjukkan bahawa kesemua pemencilan yang diteliti menolak sekurang-kurangnya 3 daripada 13 antimikrob yang diuji. Di kalangan agen-agen antimikrob, kloramfenikol, kombinasi sulfametoksazin dan trimetoprim serta nitrofurantin didapati paling efektif (sensitiviti 100%) diikuti oleh tetrasiklin

(sensitiviti 98%), penisilin (sensitiviti 88%) dan gentamisin (sensitiviti 76%). Peratus tertinggi resistan didapati lebih kepada linkomisin, basitrasin dan kloksasilin (resistan 100%), diikuti oleh furazolidon dan kolistin (masing-masing 84% dan 68%).

Ujian Agar Gel Diffusion Precipitation (AGDP) telah digunakan untuk menentukan pemencilan somatik serotip. Berdasarkan keputusan ujian AGDP, Serotip 1 didapati dominan berbanding avian yang dipencilkan daripada kawasan endemik. Serotip 3, 3 x 4 dan 4 yang pertama kali ditemui di negara Iran juga dikenal pasti di kalangan pemencilan-pemencilan lain.

Corak protein elektroforisis bagi pemencilan ini telah dikaji dengan menggunakan gel sodium dodisil sulfat poliklamid (SDS-PAGE). Kesemua strain adalah sama dalam kebanyakan jalur-jalur protein. Perbezaan utama antara corak-corak protein pemencilan ini begitu ketara pada kedudukan salah satu daripada jalur utama (H Protein) yang dibentangkan dalam kawasan 34-38 Kda. Merujuk kepada kedudukan protein H, tiga kumpulan yang berbeza telah dikenal pasti. Pada protein jenis I, jisim molekul protein H adalah lebih kurang 38 Kda tetapi di dalam protein jenis II dan jenis III, masing-masing 36.5 dan 34 Kda.

Dos lethal minimum (MLD) strain berprotein jenis I, II dan III sebagai penentu virulen telah dikenal pasti pada tikus. Ia menunjukkan bahawa strain berprotein jenis III mempunyai virulen paling sedikit dan strain berprotein jenis I mempunyai virulen paling banyak terdapat pada tikus.

Pembatasan analisis enzim (REA) kromosomal DNA dan REP-PCR digunakan untuk menentukan kepelbagaian genetik di kalangan pemencilan. Cap jari DNA oleh pencernaan HpaII membahagikan 25 pemencilan ini kepada 7 kumpulan REA, yang mana dua daripadanya mengandungi pemencilan tunggal.

Cap jari DNA beserta REP-PCR menunjukkan kepelbagaian genetik yang amat ketara di kalangan pemencilan. Berdasarkan corak DNA yang luas, sejumlah 9 REP-PCR telah ditentukan. REP-PCR menghasilkan jalur-jalur yang kuat, selari mengikut saiz daripada kira-kira 700 bp hingga 3.6 Kb beserta dua jalur spesifik-spesies 0.8 Kb dan 2.3 Kb. REP-PCR berupaya untuk membezakan pemencilan *P.multocida* di kalangan sumber-sumber dan kawasan-kawasan geografi yang berbeza. Kesimpulan daripada kajian ini menunjukkan bahawa penggunaan amplifikasi elemen-elemen REP oleh tindak balas berantai polimeras sangat produktif dan boleh dicadangkan sebagai alat epidemiologi yang sesuai untuk penyelidikan terhadap kerebakan asal dan keserupaan antara pemencilan avian *P.multocida* yang berlainan.

ACKNOWLEDGEMENTS

In the Name of Allah, the Most Beneficent and Most Merciful

First and foremost, my heartfelt thanks to the Almighty for giving me the strength and willpower to complete this challenging task.

I would like to express my appreciation and sincere gratitude to my supervisor, Associate Prof. Dr Abdul Aziz Saharee, for his invaluable guidance, suggestions and constant encouragement throughout the course of study and in the preparation of this dissertation. I am very much indebted to him for providing all the help, patiently guiding me until the completion of this thesis.

My deep appreciation and gratitude also to Prof. Dr Abdul Rani Bahaman, Dr Abdul Rahim Mutalib and Prof Aini Ideris, members of my supervisory committee in UPM for their invaluable guidance, suggestions and encouragement.

I wish to thank Dr Farhad Esmaily, Department of Biotechnology, Razi Vaccine and Serum Research Institute (RVSRI), Iran for his help and constructive suggestions. I would like to thank the Director of the RVSRI, Dr M. A. Akhavizadegan, his deputy in research affairs, Dr M. Hashemzadeh

and also the Head of the Department of Biotechnology, RVSRI, Dr M. Ashtiani, for their kind support and providing the research facilities.

I am grateful to the Ministry of Jihad Agriculture, for providing me the scholarship to enable me to complete my studies.

I am grateful to Dr M.V. Marandi, Dr S. A. Poorbakhsh, Dr A. Sotoodehnia and Dr J. V. Yousefi for supplying *P.multocida* cultures and laboratory support. I would like to thank the staff of the Department of Biotechnology, RVSRI, especially Mrs N. Mozafarian, Mr M. Esmailzadeh and Dr Z. Fayazi for their technical support. Also Thanks to Dr Siti Khairani Bejo, Institute of Bioscience, UPM for her laboratory support.

Last but not least, I wish to express my deepest appreciation to my wife, Mrs Mahtab Mozaffari, who has not only given me encouragement, moral and loving support throughout the course of my studies, but also has been very patient in taking care of our two sons whenever I needed to be alone or involved in the laboratory. Also thanks to my sons Vahid and Moein who were patient during my work as I didn't have enough time to spend with them. My gratitude and thanks are also for my mother, brothers and sisters for their encouragement. Finally, I also would like to thank all my friends who gave me encouragement and support during the period of my study.

TABLE OF CONTENTS

		Page
DEDICATION		2
ABSTRACT		3
ABSTRAK		6
ACKNOWLEDGEMENTS		9
APPROVAL SHEETS		11
DECLARATION FORM		13
LIST OF TABLES		17
LIST OF FIGURES		20
LIST OF ABBREVIATIONS		23
CHAPTER		
I	INTRODUCTION	25
II	LITERATURE REVIEW	30
	<i>Pasteurella multocida</i> , General Characteristics	30
	Taxonomy of <i>Pasteurella</i> spp	30
	Identification of <i>Pasteurella multocida</i>	31
	Isolation and Maintenance	34
	Antimicrobial Sensitivity Pattern	34
	Serodiagnosis of Pasteurellosis	35
	Cell Components of <i>P. multocida</i>	36
	Capsule	36
	Lipopolysaccharide	38
	Membrane Proteins	40
	Toxins	43
	Enzymes	45
	Fimbriae	47
	Heat Shock Proteins	48
	Plasmids	49
	Avian Pasteurellosis	49
	Classification of <i>Pasteurella multocida</i>	54
	Biotyping	54
	Serum Protection Typing	56
	Capsular Typing	57
	Somatic Serotyping	60
	Bacteriophage Typing	62
	Protein Fingerprinting	63
	Restriction Enzyme Analysis of DNA	67
	Ribotyping	71
	PCR-Based Methods	73
	Immunity to <i>P. multocida</i> Infection	81
	<i>Pasteurella multocida</i> Vaccines	82
	Whole Cell Bacterins	82

	Live Vaccines	83
	Subunit Vaccines	84
	Recombinant Vaccines	85
III	ISOLATION, BIOCHEMICAL CHARACTERIZATION AND ANTIBACTERIAL SENSITIVITY DETERMINATION	86
	Introduction	86
	Materials and Methods	87
	Isolation	87
	Maintenance of the Isolates	88
	Identification Tests	89
	Biochemical Tests	89
	Antimicrobial Sensitivity Determination	91
	Results	92
	Isolates	92
	Biochemical Characteristics	92
	Antimicrobial Sensitivity Test	93
	Discussion	102
IV	SEROTYPING OF <i>PASTEURELLA MULTOCIDA</i> ISOLATES	107
	Introduction	107
	Materials and Methods	109
	Cultures	109
	Bacterins	109
	Chickens	110
	Immunization	110
	Type Specific Antisera	111
	Preparation of Antigen	111
	Agar Gel Diffusion Precipitin Test	112
	Results	113
	Discussion	120
V	ELECTROPHORETIC PROTEIN PROFILES OF <i>PASTEURELLA MULTOCIDA</i> ISOLATES	125
	Introduction	125
	Materials and Methods	128
	Cultures	128
	Protein Preparation	128
	Polyacrylamide Gel Electrophoresis	129
	Determination of Molecular Weight	131
	Determination of Minimum Lethal Dose	132
	Bacterial Dilutions and Colony Count	133
	Inoculation	134
	Cross Protection Study	134

	Statistic Analysis	135
	Results	136
	Electrophoretic Protein Patterns of Isolates	136
	Minimum Lethal Dose	138
	Cross Protection	139
	Discussion	148
VI	CHARACTERIZATION OF <i>PASTEURELLA MULTOCIDA</i> ISOLATES BY RESTRICTION ENDONUCLEASE ANALYSIS OF DNA AND AMPLIFICATION OF REPETITIVE EXTRAGENIC ELEMENTS	155
	Introduction	155
	Materials and Methods	158
	Cultures	158
	Preparation of Whole Cell DNA	158
	Restriction Enzyme Digestion of the Genomic DNA	160
	Polymerase Chain Reaction	161
	Agarose Gel Electrophoresis	163
	Results	165
	Restriction Enzyme Analysis	165
	Amplification of Repetitive palindromic (REP) Elements by PCR	174
	Discussion	185
	Restriction Enzyme Analysis	185
	Amplification of REP Elements by PCR	190
VII	GENERAL DISCUSSION AND CONCLUSION	194
	BIBLIOGRAPHY	206
	APPENDICES	232
	BIODATA OF AUTHOR	252

LIST OF THE TABLES

Table		Page
2-1	Differential properties of <i>P.multocida</i> compared to other avian Pasteurella species.	33
2-2	Sulphonamides and antibiotics beneficially used to treat fowl cholera.	53
2-3	Differential properties of <i>P.multocida</i> subspecies.	55
2-4	Differential properties of <i>P.multocida</i> biovars.	56
2-5	Serologic and nonserologic methods introduced for capsular typing of <i>P.multocida</i>	59
2-6	The recognition sequences for some of the most frequently used restriction endonucleases	69
2-7	Sequences of PCR primers that have been used in the identification and characterization of <i>P.multocida</i> .	75
3-1	Origins of twenty-five <i>P.multocida</i> isolates obtained from poultry in northern part of Iran.	95
3-2	Distribution of <i>P.multocida</i> isolated from poultry in endemic area of Fowl cholera.	96
3-3	Biochemical characteristics of 25 isolates of <i>P.multocida</i> obtained from poultry in Iran.	97
3-4	The results of antimicrobial sensitivity test in 25 <i>P.multocida</i> isolates obtained from poultry in Iran.	98
3-5	Number and percent of sensitivity, intermediate and	

	resistant isolates of avian <i>P.multocida</i> against 13 therapeutic agents.	101
3-6	Distribution of <i>P.multocida</i> isolates according to their multiple antibacterial resistance.	102
4-1	Serotypes of 25 isolates of <i>P.multocida</i> obtained from poultry in Iran.	118
4-2	Distribution of <i>P.multocida</i> serotypes according to sources in the endemic provinces.	119
4-3	Frequency of identified serotypes among chickens, ducks and geese.	120
5-1	Results of protein pattern of 25 <i>P.multocida</i> isolates obtained from poultry in Iran.	144
5-2	Protein types of 25 <i>P.multocida</i> isolates obtained from poultry in Iran.	145
5-3	Distribution of somatic serotypes among three protein typing of avian <i>P.multocida</i> isolates	146
5-4	Distribution of protein types of <i>P.multocida</i> among different sources in endemic area of Iran.	146
5-5	Results of challenge with dilutions of <i>P.multocida</i> for determination of MLD in mice.	147
5-6	Results of challenge of immunized mice by strain PMI030 against homologous and heterologous protein types.	147
6-1	Results of restriction enzyme (HpaII) analysis of chromosomal DNA of 25 <i>P.multocida</i> strains isolated from poultry.	171
6-2	REA types of 25 <i>P.multocida</i> obtained from poultry generated by restriction enzyme HpaII.	173
6-3	Distribution of <i>P.multocida</i> isolates according to their HpaII REA types and somatic serotypes,	173
6-4	Distribution of <i>P.multocida</i> isolates according to their HpaII REA types and protein fingerprinting patterns.	174

6-5	Results of REP-PCR DNA fingerprinting of 25 <i>P.multocida</i> isolates obtained from poultry in Iran.	181
6-6	REP types of 25 <i>P.multocida</i> isolates obtained from poultry in Iran.	182
6-7	Distribution of REP types among different sources of <i>P.multocida</i> in the endemic area of FC in Iran.	183
6-8	Distribution of REP types of <i>P.multocida</i> among different district in the FC endemic area.	183
6-9	Presence (+) or absence (-) of bands amplified by REP-PCR In the DNA fingerprinting of 25 <i>P.multocida</i> isolated from poultry in Iran.	184

LIST OF FIGURES

Figure		Page
4-1	Gel precipitin pattern showing the reaction between <i>P.multocida</i> PMI033 heat stable antigen and serotype 1 antisera. The five reference antisera (1-5) are in the outer wells and heat stable antigen in the centre well.	114
4-2	Gel precipitin pattern of <i>P.multocida</i> PMI030 heat stable antigen and serotype 1 antisera. The five reference antisera (1-5) are in the outer wells and heat stable antigen in the centre well.	115
4-3	Immunodiffusion gel showing the reaction between <i>P.multocida</i> PMI035 heat stable antigen and serotype 3 antisera. The antigen was loaded in centre well and typing antisera were loaded in the outer wells. Wells 1-5 contained typing antisera against serotypes 1-5 respectively.	116
4-4	Immunodiffusion pattern of <i>P.multocida</i> PMI032 heat stable antigen (centre well) and serotype 3 and 4 reference antisera). Typing antisera were: serotypes 1 (well 1); serotype 2 (well 2); serotype 3 (well 3); serotype 4 (well 4); Serotype 5 (well 5).	117
5-1	SDS-PAGE protein profile of <i>P.multocida</i> isolates. Lanes:1, The protein marker; 2, Strain PMI030; 3, Strain PMI033; 4, Strain PMI034; 5, Strain PMI036; 6, Strain PMI037; 7, Strain PMI039; 8, Strain PMI038; 9, Strain PMI040; 10, Strain PMI041.	140
5-2	Protein fingerprinting patterns of <i>P.multocida</i> isolates. Lanes: 1, Strain PMI022; 2, Strain PMI047; 3, Strain PMI026; 4, Strain PMI024; 5, Strain PMI023; 6, Strain PMI 044; 7, Strain PMI043; 8, Strain PMI042; 9, Strain PMI030 and 10 The protein molecular weight marker.	141
5-3	SDS-PAGE protein fingerprinting of <i>P.multocida</i> isolates. Lanes 1: The protein Marker; 2, Strain PMI030;	

- 3, Strain PMI028; 4, Strain PMI047; 5, Strain PMI046; 6, Strain PMI020; 7, Strain PMI031; 8, Strain PMI025; 9, PMI032; and 10 protein molecular weight marker. 142
- 5-4 Protein fingerprinting pattern of reference strains compare to local vaccine strain. Lanes: 1, strain P-1702 (Serotype 5); 2, strain P-1662 (Serotype 4); 3, strain P-1056 (Serotype 3); 4, strain M-1404 (Serotype 2); 5, strain X-73 (Serotype 1) and 6, strain PMI030 (Serotype 1, vaccine strain). 143
- 6-1 Agarose gel electrophoresis of whole cell DNA obtained from *P.multocida* isolated from poultry with fowl cholera after digested with HpaII. Lane contain DNAs from isolates PMI031 (lane 1), PMI047 (lane 2), PMI035 (lane 3), PMI032 (lane 4), PMI037 (lane 5), PMI024 (lane 6), PMI023 (lane 7), PMI043 (lane 8) and the DNA marker (SPP1 DNA cleaved with EcoR1), with numbers on the right representing the number of kilo base pairs of the respective fragments. 167
- 6-2 Restriction fragment patterns of chromosomal DNA of *P.multocida* strains obtained from poultry in Iran. Samples were digested with HpaII and electrophoresed in agarose as described in materials and methods. Lanes: 1, Strain PMI041; 2, Strain PMI042; 3, Strain PMI040; 4, Strain PMI039; 5, strain PMI038; 6, Strain PMI036; 7, Strain PMI033; 8, Strain PMI030 and lane 9, SPP1 DNA cleaved with EcoR1 as marker. 168
- 6-3 Restriction fragment pattern of chromosomal DNA of *P.multocida* isolates. Lanes: 1, Strain PMI026; 2, strain PMI046; 3, Strain PMI045; Strain PMI044; 5, Strain PMI034; and 6, EcoR1 digested SPP1 DNA marker. 169
- 6-4 Restriction fragment patterns of 5 *P.multocida* isolates. Samples were digested with HpaII and electrophoresed in agarose. Lanes: 1, Strain PMI020; 2, Strain PMI047; 3, Strain PMI022; 4, Strain PMI028; 5, Strain PMI025, and 6 the DNA marker. 170

- 6-5 Schematic diagram of REA groups (I-VII) of *P.multocida* isolates according to the pattern of the heavy DNA fragments that were produced following digestion with restriction enzyme HpaII. 171
- 6-6 REP-PCR products generated by amplification of *Pasteurella multocida* genomic DNA with REP primers. PCR conditions included an annealing temperature of 42°C and a total of 34 cycles. Lanes: 2, Strain PMI030; 3, Strain PMI033, 4, Strain PMI034; 5, Strain PMI036; 6, Strain PMI039; 7, Strain PMI041; 8, Strain PMI038; 9, Strain PMI045 and 10 strain PMI046. Lane 1 is SPP1 DNA cleaved with EcoR1 as marker. Numbers representing the number of kilobase pairs (kbp) of the respective fragments. 177
- 6-7 Amplification of *P.multocida* genomic DNA by REP primers. REP-PCR products were electrophoresed in 2 percent agarose as explained in materials and methods. Lanes: 1, Strain PMI044; 2, Strain PMI037; 3, Strain PMI023; 4, Strain PMI035; 5, Strain PMI047; 6, PMI032; 7, Strain PMI043; 8, Strain PMI042; 9, Strain PMI030 and 10 DNA Marker. 178
- 6-8 REP-PCR products generated by amplification of *P.multocida* chromosomal DNA with REP primers. PCR conditions included an annealing temperature of 42C and a total of 34 cycles. Lanes: 1, PMI031; 2, PMI028; 3, PMI025; 4, PMI047; 5, PMI040; 6, PMI026; 7, PMI024; 8, PMI020; 9, PMI030 and 10 DNA marker. 179
- 6-9 Schematic representation of the nine REP types (I-IX) of the *P.multocida* isolates following genomic DNA amplification by REP-PCR. 180

LIST OF ABBREVIATIONS

AGDP	Agar Gel Diffusion Precipitin Test
AP-PCR	Arbitrary Primed PCR
APS	Ammonium Persulfate
BHI	Brain Heart Infusion
°C	Degree Celcius
CCA	Crude Capsule Antigens
CCF	Cell-Free Culture Filtrate
CFU	Colony Forming Unit
CP	Capsular Polysaccharide
CPFs	Cross-Protection Factors
CU strain	Clemson University Strain
DATP	Deoxyadenosine Triphosphate
DCTP	Deoxycytidine Triphosphate
DGTP	Deoxyguanosine Triphosphate
DTTP	Deoxythymidine Triphosphate
DNTP	Deoxynucleotide Triphosphate
DIE	Descriptive Identification Epithet
EDTA	Ethylendiamine Tetra Acetate
ELISA	Enzyme Linked Immunosorbent Assay
FC	Fowl Cholera
HS	Haemorrhagic Septicaemia
HSA	Heat Stable Antigen
HSPs	Heat Shock Proteins
IHA	Indirect Haemagglutination Test
IVO	Iranian Veterinary Organization
KDa	Kilo Daltons
KSCN	Potassium Thiocyanate Extracts
LD 100	Lethal Dose of 100 Percent
LPS	Lipopolysaccharide
MAb	Monoclonal Antibody
Mda	Mega Daltons
MHA	Muller-Hinton Agar
MLD	Minimum Lethal Dose
MOMPs	Major Outer Membrane Proteins
OMPs	Outer Memberane Proteins
<i>P.multocida</i>	<i>Pasteurella multocida</i>
PAGE	Polyacrylamide Gel Electrophoresis
PCR	Polymerase Chain Reaction
PMSF	Phenylmethyl Sulfonyl Fluoride
PMT	<i>Pasteurella multocida</i> Toxin

RAPD	Random Amplified Polymorphic DNA
RD Complex	Respiratory Disease Complex
REA	Restriction Endonuclease Analysis
REP	Repetitive Extragenic Palindrome
rRNA	Ribosomal RNA
RVSRI	Razi Vaccine and Serum Research Institute
<i>Taq</i>	<i>Termophilus aquaticus</i>
SDS	Sodium Dodecyl sulphate
SPF	Specific Pathogen Free
Subsp	Subspecies
TAE	Tris Acetate EDTA
TEMED	Tetrametylendiamine
Tox	Toxin
TSI	Three Sugar Iron Agar
U	Unit
VRI	Veterinary Research Institute
WC	Whole Cell