

August 2024

First serological evidence of *Coxiella burnetii* and *Mycobacterium avium* subspecies *paratuberculosis* (MAP) among deer in an institutional farm in Malaysia

Bura Thlama Paul
Universiti Putra Malaysia, burapaul@upm.edu.my

Faez Firdaus Abdullah Jesse
Universiti Putra Malaysia, jesse@upm.edu.my

Shivnraj Sundar Universiti Putra Malaysia
Universiti Putra Malaysia, 201215@student.upm.edu.my

Nur Hanis Nadhirah Yosryhan
Universiti Putra Malaysia, hanisnadhira10@gmail.com

Yonis Ahmed Ahmed Jimale Dr
Universiti Putra Malaysia, yonisahmed2018@gmail.com



Part of the [Large or Food Animal and Equine Medicine Commons](#), [Life Sciences Commons](#), and the [Veterinary Preventive Medicine, Epidemiology, and Public Health Commons](#)
See next page for additional authors

Recommended Citation

Paul, Bura Thlama; Jesse, Faez Firdaus Abdullah; Sundar, Shivnraj Universiti Putra Malaysia; Yosryhan, Nur Hanis Nadhirah; Jimale, Yonis Ahmed Ahmed Dr; Chung, Eric Lim Teik Dr; and Mohd Lila, Mohd Azmi Prof (2024) "First serological evidence of *Coxiella burnetii* and *Mycobacterium avium* subspecies *paratuberculosis* (MAP) among deer in an institutional farm in Malaysia," *The Thai Journal of Veterinary Medicine*: Vol. 54: Iss. 1, Article 10.

DOI: <https://doi.org/10.56808/2985-1130.3665>

Available at: <https://digital.car.chula.ac.th/tjvm/vol54/iss1/10>

This Short Communication is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

First serological evidence of *Coxiella burnetii* and *Mycobacterium avium* subspecies *paratuberculosis* (MAP) among deer in an institutional farm in Malaysia

Authors

Bura Thlama Paul, Faez Firdaus Abdullah Jesse, Shivrroj Sundar Universiti Putra Malaysia, Nur Hanis Nadhirah Yosryhan, Yonis Ahmed Ahmed Jimale Dr, Eric Lim Teik Chung Dr, and Mohd Azmi Mohd Lila Prof

**First serological evidence of *Coxiella burnetii* and
Mycobacterium avium subspecies *paratuberculosis* (MAP)
among deer in an institutional farm in Malaysia**

Bura Thlama Paul^{1,2} Faez Firdaus Abdullah Jesse^{3*} Shivnraj Sundar³

Nur Hanis Nadhirah Yosryhan³ Nur Amira Azhar³ Yonis Ahmed Jimale^{3,4}

Mohd Jefri Norsidin³ Eric Lim Teik Chung⁵ Mohd Azmi Mohd Lila⁶

Abstract

Q fever is a zoonotic disease caused by *Coxiella burnetii*, which is associated with reproductive disorders such as abortion and reproductive problems in ruminants, while paratuberculosis is an underdiagnosed disease caused by *Mycobacterium avium paratuberculosis* (MAP) which causes chronic wasting, low fertility, and mortality in ruminants. Although both *C. burnetii* and MAP infections have been reported in domestic ruminants worldwide, there is no data on their seroprevalence status among deer in Selangor, Malaysia. This study investigated the seroprevalence status of *C. burnetii* and MAP among deer in Malaysia. Ninety-two blood sera collected from an institutional deer farm were screened using commercial Multi-species ID Screen indirect ELISA tests to detect specific anti-*C. burnetii* (FQS-MS/1117) and anti-MAP (PARAS Ver 0516) antibodies. The ELISA results showed a 14.1% (95% CI = 8.5-22.7) apparent and 14.8% (95% CI = 8.4-24.4) true prevalence for *C. burnetii* and a 2.2% (95% CI = 0.6-7.6) apparent and 1.3% (95% CI = -0.45-7.40) true prevalence for MAP among deer. There were statistically significant associations between seropositivity to *C. burnetii* and the age groups ($\chi^2 = 7.325$, $P = 0.011$) or herds of deer ($\chi^2 = 9.732$, $P = 0.001$). To the best of our knowledge, this study provides the first preliminary serological evidence of *C. burnetii* and MAP in *Cervus timorensis*, suggesting potential threats to animal and public health in Malaysia.

Keywords: Deer, *Coxiella burnetii*, *Mycobacterium avium paratuberculosis*, seroprevalence, Malaysia

¹Department of Animal Science and Fisheries, Faculty of Agricultural and Forestry Science, Universiti Putra Malaysia Campus Bintulu Sarawak, Malaysia

²Veterinary Teaching Hospital, Faculty of Veterinary Medicine, University of Maiduguri, 600230 Maiduguri, Borno State, Nigeria

³Department of Veterinary Clinical Studies, Faculty of Veterinary Medicine, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

⁴Faculty of Veterinary Medicine and Animal Husbandry, Somali National University, Mogadishu P.O. Box 15, Somalia

⁵Department of Animal Science, Faculty of Agriculture, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia

⁶Department of Veterinary Microbiology and Pathology, Faculty of Veterinary Medicine, 43400 UPM Serdang, Selangor Malaysia

*Correspondence: jesse@upm.edu.my (F.F.A. Jesse)

Received February 13, 2024

Accepted May 26, 2024

Introduction

Coxiella burnetii is a zoonotic Gram-negative pathogenic bacterium that causes coxiellosis in animals and Q fever in humans (Celina and Cerný, 2022). The organism is present in body secretions or excretions (Agerholm, 2013). Transmission in animals may be airborne by inhalation or vector-borne by ticks, while human infection may be due to direct anthroponotic transmission from animals through body secretions/excretions, tick-borne transmission, or indirect occupational exposure (Celina and Cerný, 2022). Coxiellosis is mostly asymptomatic, but metritis, infertility, abortion, premature delivery, stillbirth, and the birth of weak neonates may be observed (Maurin and Raoult, 1999; Arricau-Bouvery and Rodolakis, 2005; Eldin et al., 2017). Q fever causes mild flu-like symptoms or severe fatal complications in humans (Angelakis and Raoult, 2010). It is a disease of economic and public health significance (Keshavamurthy et al., 2019; Jesse et al., 2020), and wildlife species, including deer, serve as reservoir hosts (Celina and Cerný, 2022). Although *C. burnetii* has been detected and reported in humans (Rai et al., 2011; Khor et al., 2018) and livestock (Nurkunasegran et al., 2017; Khor et al., 2018; Jesse et al., 2020) in Malaysia, there is still a lack of published information on its seroprevalence status and risk factors among wild ruminants and farmed deer.

Mycobacterium avium subspecies *paratuberculosis* (MAP) is a zoonotic bacterium that causes paratuberculosis or Johne's disease (JD) in ruminants (Ssekitoleko et al., 2021; Yu et al., 2022) and Crohn's disease in humans (Hermon-Taylor et al., 2000). The MAP organism is shed in the feces, milk, and colostrum of infected animals, and transmission occurs via the fecal-oral route (Khol et al., 2017). Johne's disease has a global distribution and is observed in domestic (Barrero-Domínguez et al., 2019; Pourmahdi Borujeni et al., 2021) and wild (Budhe et al., 2014; Matos et al., 2017) ruminants. Although JD is a notifiable disease, it is grossly underreported due to its subclinical course and lack of routine monitoring mechanisms in many countries (Manning and Collins, 2001). Although there is a recent report on prevalence of MAP among small ruminant livestock in Malaysia (Jimale et al., 2024), there is paucity of information on its seroprevalence among deer in the country.

This study was necessary due to the global economic and zoonotic impacts of Q fever and paratuberculosis and the recognition of wildlife as a reservoir and source of infection for domestic ruminants and man, coupled with the lack of epidemiological data on the seroprevalence status of these important zoonoses in Malaysia. In this study, we investigated the seroprevalence status of Q fever and paratuberculosis in farmed deer livestock to shed light on the epidemiology of *C. burnetii* and MAP and furnish preliminary data for implementing further studies and designing comprehensive control programs in Malaysia.

Materials and Methods

Study design and sampling: The study received ethical clearance from the Institutional Animal Care and Use Committee (IACUC) of Universiti Putra Malaysia

(UPM/IACUC/AUP-U042/2023). The deer stock (*Cervus timorensis*) belonged to an institutional farm located within latitude 2.982962 and longitude 101.729190 in Selangor, Malaysia. The deer herds were semi-intensively managed, openly grazed during the day with supplemented pelletized concentrate feeding, and were allowed to breed naturally. A total of 92 blood sera comprising 36 fresh samples collected between January and August 2023 (new herd) and 56 archived serum samples collected in 2017 (old herd) were included in this study. Sample collection was done conveniently based on availability due to the difficulty in restraining the deer. Whenever possible, individuals were manually restrained before withdrawing 5 mL of whole blood by jugular venipuncture for serum extraction. Epidemiological variables of the deer, such as age, gender, and herd, were also recorded. The deer were further categorized according to their age group as young (≤ 2 years) and adults (> 2 years).

Detection of serum antibody by ELISA: Commercial Multi-species ID Screen (ID.VET, France) indirect ELISA screening tests were used to detect serum antibodies directed against *C. burnetii* (FQS-MS/1117) and MAP (PARAS Ver 0516) according to manufacturers instructions. The optical densities of ELISA microplates were measured at 450 nm wavelength using an ELISA microplate absorbance reader (Infinite® 200 PRO). The samples with (S/P% > 70%) were interpreted as positive for MAP, while samples with S/P% > 50% were interpreted as positive for *C. burnetii*.

Statistical analysis: The Epitools statistical calculator was used to calculate the apparent and true prevalence of *C. burnetii* or MAP with their respective 95% confidence intervals (CI) (Rogan and Gladen, 1978; Sergeant, 2018). The Chi-square test using the Fisher's Exact method was computed using the Statistical Package for Social Sciences (SPSS) Software version 29.0 macOS to determine the associations between the seroprevalence of *C. burnetii* or MAP infection as the dependent variables and the age, gender, or herd of deer as the independent variables.

Results and Discussion

The ELISA assays revealed 1.32% (95% CI = -0.5-7.4) and 14.8% (95% CI = 8.4-24.4) true prevalence for MAP and *C. burnetii* (Table 1) with significant associations between *C. burnetii* infection and the age groups ($\chi^2 = 7.325$, $P = 0.011$) and herds ($\chi^2 = 9.732$, $P = 0.001$) of deer. This study is the first report on the prevalence of *C. burnetii* and MAP infection among deer in Malaysia. The result of this study agrees with previous reports of 12.3% for *C. burnetii* in China (Cong et al., 2015), 15.4% in Spain (Ruiz-Fons et al., 2008), and 14.5% in the United States (Kirchgesner et al., 2013). However, our result is higher than previous reports of 1.4% (Yangho et al., 2011), 9.18% (Shin et al., 2014) in Korea, and 3.4% in eastern Australia (Voss et al., 2023). The high seroprevalence of *C. burnetii* infection observed among deer livestock in this study is due to optimal environmental conditions

(Mori and Roest, 2018). Grazing of deer in semi-intensive management, as seen in this study, increases their risk of exposure to the natural tick vectors of *C. burnetii* (Nusinovici et al., 2015). *Hyalomma marginatum marginatum* and *Hyalomma lusitanicum* increased the risk of coxiellosis in Spain (Ruiz-Fons et al., 2008). The higher seropositivity in adult deer may be due to repeated exposure with increasing age (Rizzo et al., 2016). Moreover, their proximity to cattle and goat livestock may lead to cross-infection (Celina and Cerný, 2022). Although a high seroprevalence of *C. burnetii* infection is accompanied by reproductive issues in animals and humans (Ullah et al., 2022), no clinical cases of abortion, stillbirths, or infertility were previously reported on the farm. Thus, the absence of clinical disease or history of reproductive disorders in the studied deer farm has led us to suggest that the observed high seropositivity could be attributed to past exposure of deer to *C. burnetii* because serological testing used in this study only indicates exposure. Without further molecular diagnostics, it is not likely to suggest that the presence of antibodies indicates an active infection.

The current observation of 1.3% seropositivity to MAP among captive deer livestock agrees with previous reports of 1.9%, 3.8, 3.4% in various types of deer in Portugal (Tryland et al., 2004), 1.7% in Mauritius (Jori et al., 2014), and 0.5% (Sleeman et al., 2009) and 0.3% (Raizman et al., 2005) in the United States. However, our result is lower than 16% in Mexico (Lozano-Cavazos et al., 2021), 17.6% in China (Meng et al., 2015), and 30.2% in Spain (Reyes-García et

al., 2008). The MAP organism is known to be very hardy and survives for up to 48 weeks outside the host in water bodies and sediments (Whittington et al., 2019). The result represents a significant risk for domestic animals and other wildlife due to their potential to harbor subclinical disease (Sleeman et al., 2009). MAP infection may potentially spill over from deer to domestic animals and wildlife in areas that practice semi-intensive or extensive management systems (Geraghty et al., 2014).

This study provides preliminary serological data showing a high level of exposure to *C. burnetii* among deer livestock, suggesting potential threats to animal and public health in Malaysia. Furthermore, this study is, to the best of our knowledge, the first documented evidence of serum antibodies towards *C. burnetii* and MAP in *Cervus timorensis* in Malaysia. Although no vaccination was practiced against Q fever and paratuberculosis in the herds to suggest vaccine antibodies, further studies using more sensitive molecular diagnostics are needed to confirm and characterize MAP and *C. burnetii* circulating among deer and elucidate their potential role in the epidemiology of coxiellosis and paratuberculosis in Malaysia. A one-health approach towards controlling paratuberculosis and Q fever is recommended among livestock and people with occupational exposure to MAP and *C. burnetii*, such as veterinarians, animal health workers, abattoir workers, animal traders, environmental workers, and public health care workers.

Table 1 The apparent and true Seroprevalence of *Coxiella. burnetii* and *M. avium paratuberculosis* in deer livestock in Selangor, Malaysia

Variables	Tested	<i>Coxiella. burnetii</i>				<i>M. avium paratuberculosis</i>			
		Apparent	95 % CI	True	95 % CI	Apparent	95 % CI	True	95 % CI
Age									
Young	59	4 (6.8)	2.7-16.2	6.5	1.9-17.1	0 (0.0)	0.0- 6.1	-1.12	-0.1-5.7
Adult	33	9 (27.3)	15.1-44.2	29.5	15.8-48.6	2 (6.1)	0.2-19.6	5.69	0.8-20.9
Gender									
Male	22	1 (4.5)	0.8-21.8	4.0	-0.9-23.4	0 (0.0)	0.0-14.9	-0.01	-0.0-15.6
Female	70	12 (17.1)	10.1-27.6	18.1	10.2-29.9	2 (2.9)	0.8-9.8	2.09	-0.2-9.9
Herd									
Old	56	13 (23.2)	14.1-39.1	24.9	14.7-39.1	2 (3.6)	0.9-12.1	2.89	0.0-12.5
New	36	0 (0.0)	0.0-9.6	-1.1	-1.1-9.7	0 (0.0)	0.0-9.6	-1.12	-1.1-9.7
Overall	92	13 (14.1)	8.5-22.7	14.8	8.4-24.4	2 (2.2)	0.6-7.6	1.3	-0.5-7.4

CI: confidence interval.

Table 2 Univariable analysis for the associations between seropositivity and epidemiological variables

Variables	Tested	<i>Coxiella. burnetii</i>				<i>M. avium paratuberculosis</i>			
		positive	negative	χ2	p-value	positive	negative	χ2	p-value
Age									
Young	59	4 (6.8)	55 (93.2)	7.325	0.011*	0 (0.00)	59 (100)	3.655	0.126
Adult	33	9 (27.3)	24 (72.7)			2 (6.1)	31 (93.9)		
Gender									
Male	22	1 (4.5)	21 (95.5)	2.189	0.179	0 (0.0)	22 (100)	0.643	1.000
Female	70	12 (17.1)	58 (82.9)			2 (2.9)	68 (97.1)		
Herd									
Old	56	13 (23.2)	43 (76.8)	9.732	0.001*	2 (3.6)	54 (96.4)	1.314	0.518
New	36	0 (0.0)	36 (100)			0 (0.0)	36 (100)		

χ²: Chi-square, P-values (Fisher's Exact test) with an asterisk (*) are statistically significant.

Acknowledgments

The authors are grateful to Mr Jefri Mohammed Norsidin in the Clinical Research Laboratory and Puan Wan Nur Ayuni Wan Noor in the Virology Laboratory, Faculty of Veterinary Medicine, Universiti Putra Malaysia, for supporting the field and laboratory aspects of this research.

References

- Agerholm J 2013. *Coxiella burnetii* associated reproductive disorders in domestic animals-a critical review. *Acta Vet Scand.* 55:13-24.
- Angelakis E and Raoult D 2010. Q fever. *Vet Microbiol.* 140(3-4):297-309.
- Arricau-Bouvery N, Rodolakis A 2005. Is Q fever an emerging or re-emerging zoonosis? *Vet Res.* 36:327-49.
- Barrero-Domínguez B, Luque I, Huerta B, Gomez-Laguna J, Galán-Relaño Á, Gómez-Gascón L, Sánchez, M and Astorga R.J 2019. Paratuberculosis in dairy goat flocks from southern Spain: Risk factors associated with seroprevalence. *Vet Rec.* 185:600.
- Bauman CA, Jones-Bitton A, Menzies P, Toft N, Jansen J, Kelton D 2016. Prevalence of paratuberculosis in the dairy goat and dairy sheep industries in Ontario, Canada. *Can Vet J.* 57(2):169.
- Browne AS, Fèvre EM, Kinnaird M, Muloi DM, Wang CA, Larsen PS, O'Brien T and Deem SL 2017. Serosurvey of *Coxiella burnetii* (Q fever) in Dromedary Camels (*Camelus dromedarius*) in Laikipia County, Kenya. *Zoonoses Public Health.* 64(7):543-9.
- Budhe SD, Rode AM and Pawade MM 2014. Prevalence of tuberculosis and paratuberculosis in some captive deer of Maharashtra. *Indian J F Vet.* 10(2):43-4.
- Carbonero A, Guzmán LT, Montaña K, Torralbo A, Arenas-Montes A and Saa LR 2015. *Coxiella burnetii* seroprevalence and associated risk factors in dairy and mixed cattle farms from Ecuador. *Prev Vet Med.* 118(4):427-35.
- Cecchi F, Fratini F, Cerri D, Bandecchi P, Cantile C and Mazzei M 2019. Small ruminant lentivirus and *Mycobacterium avium* subsp. *Paratuberculosis*: Co-infection prevalence and preliminary investigation on genetic resistance to both infections in a garfagnina goat flock. *Large Anim Rev.* 25(3):89-92.
- Celina SS and Cerný J 2022. *Coxiella burnetii* in ticks, livestock, pets and wildlife: A mini-review. *Front Vet Sci.* 9:1068129.
- Cong W, Meng QF, Shan XF, Sun WW, Kang YH, Chen L, Wang WL and Qian AD 2015. *Coxiella burnetii* (Q fever) Infection in Farmed Ruminants in Three Northeastern Provinces and Inner Mongolia Autonomous Region, China. *Vector-Borne Zoonotic Dis.* 15(8):512-4.
- de Souza EAR, de Castro EMS, de Oliveira GMB, Azevedo SS, Peixoto R de M, Labruna MB and Horta MC 2018. Serological diagnosis and risk factors for *Coxiella burnetii* in goats and sheep in a semi-arid region of northeastern Brazil. *Rev Bras Parasitol Vet.* 27(4):514-20.
- Paul B.T. et al. / *Thai J Vet Med.* 2024. 54(1): 77-82.
- Eldin C, Mélenotte C, Mediannikov O, Ghigo E, Million M, Edouard S, Mege JL, Maurin M and Raoult D 2017. From Q fever to *Coxiella burnetii* infection: a paradigm change. *Clin Microbiol Rev.* 30(1):115 LP - 190.
- Elmagzoub WA, Adam NM, Idris SM, Mukhtar ME, Abdelaziz SA, Okuni JB, Ojok L, Wahed AA, El Eltayeb E, Gameel AA and Eltom KH 2020. Seroprevalence of *Mycobacterium avium* subsp. *Paratuberculosis* in dairy cattle in Khartoum state, Sudan. *Vet Sci.* 7(4):1-8.
- Geraghty T, Graham DA, Mullowney P and More SJ 2014. A review of bovine Johne's disease control activities in 6 endemically infected countries. *Prev Vet Med.* 116(1-2):1-11.
- Hermon-Taylor J, Bull TJ, Sheridan JM, Cheng J, Stellakis ML and Sumar N 2000. Causation of Crohn's disease by *Mycobacterium avium subspecies paratuberculosis*. *Can J Gastroenterol.* 14(6):521-39.
- Jesse FFA, Paul BT, Hashi HA, Chung ELT, Abdurrahman NA, Lila MAM 2020. Seroprevalence and risk factors of Q fever in small ruminant flocks in selected States of Peninsular Malaysia. *Thai J Vet Med.* 50(4):511-7.
- Jimale YA, Jesse FF, Paul BT, Chung EL, Zakaria A, Azhar NA and Mohd Lila MA 2024. Seroprevalence and contributing factors of transboundary animal diseases in sheep and goats: a study in Peninsular Malaysia. *Trop Anim Hlth Prod.* 56(6):212.
- Jolianne MR, Roes HIJ, Tulden PW van, Kik MJL, IJzer J and Gröne A 2011. *Coxiella burnetii* Infection in Roe Deer during Q Fever Epidemic, the Netherlands. *Emerg Infect Dis.* 17(12):2369-71.
- Jori F, Godfroid J, Michel AL, Potts AD, Jaumally MR, Sauzier J and Roger M 2014. An assessment of zoonotic and production limiting pathogens in rusa deer (*Cervus timorensis* rusa) from Mauritius. *Transbound Emerg Dis.* 61:31-42.
- Keshavamurthy R, Singh BB, Kalambhe DG, Aulakh RS and Dhand NK 2019. Prevalence of *Coxiella burnetii* in cattle and buffalo populations in Punjab, India. *Prev Vet Med* 166:16-20.
- Khol JL, Braun AL, Slana I, Kralik P and Wittek T 2017. Testing of milk replacers for *Mycobacterium avium* subsp. *paratuberculosis* by PCR and bacterial culture as a possible source for Johne's disease (*paratuberculosis*) in calves. *Prev Vet Med.* 144:53-6.
- Khor CS, Mohd-Rahim NF, Hassan H, Chandren JR, Nore SS and Johari J, Loong SK, Abd-Jamil J, Khoo JJ, Lee HY, Pike BL, Wong LP, Lim YAL., Zandi K and Abubakar S 2018. Seroprevalence of Q Fever among the Indigenous People (Orang Asli) of Peninsular Malaysia. *Vector-Borne Zoonotic Dis.* 18(3):131-7.
- Kirchgessner MS, Dubovi EJ and Whipps CM 2013. Disease Risk Surface for *Coxiella burnetii* Seroprevalence in White-Tailed Deer. *Zoonoses Public Health.* 60(7):457-60.
- Kopečna M, Lamka J, Parmova I, Trcka I, Svastova P, Bartos M and Pavlik I 2006. Paratuberculosis in wild ruminants in the Czech Republic from 1997 to 2004. *Vet Arh.* 76:19-26.
- Kováčová E, Kazár J and Spanělová D 1998. Suitability of various *Coxiella burnetii* antigen preparations for

- detection of serum antibodies by various tests. *Acta Virol.* 42(6):365–8.
- Lozano-Cavazos EA, Velázquez-Morales JV, Santillán-Flores MA, Tarango-Arámbula LA, Olmos-Oropeza G, Martínez-Montoya JF and Cortez-Romero C 2021. Seroprevalence of antibodies against *Mycobacterium avium* subsp. *paratuberculosis* and its relationship to age and sex of Texas white-tailed deer (*Odocoileus virginianus*) in Coahuila, Mexico. *J Wildl Dis.* 57(2):321–6.
- Manning EJB and Collins MT 2001. *Mycobacterium avium* subsp. *paratuberculosis*: Pathogen, pathogenesis and diagnosis. *OIE Rev Sci Tech.* 20(1):133–50.
- Matos AC, Figueira L, Martins MH, Matos M, Álvares S, Mendes A, Pinto ML and Coelho AC 2017. Detection of mycobacterium *avium* subsp. *Paratuberculosis* in kidney samples of red deer (*Cervus elaphus*) in Portugal: Evaluation of different methods. *J Vet Med Sci.* 79(3):692–8.
- Maurin M and Raoult M 1999. Q fever. *Clin Microbiol Rev.* 12(4):518–53.
- McCaughey C, Murray LJ, McKenna JP, Menzies FD, McCullough SJ, O’neill HJ, Wyatt DE, Cardwell CR and Coyle PV 2010. *Coxiella burnetii* (Q fever) seroprevalence in cattle. *Epidemiol Infect.* 138(1):21–7.
- Meng QF, Li Y, Yang F, Yao GZ, Qian AD and Wang WL and Cong W 2015. Seroprevalence and risk factors of *Mycobacterium avium* subspecies *paratuberculosis* infection in domestic sika deer in China. *Trop Anim Health Prod.* 47(5):999–1003.
- Morales-Pablos MI, Mejía-Sánchez P, Díaz-Aparicio E, Palomares-Resendiz EG and Gutiérrez-Hernández JL, Reyna-Granados JR, Luna-Nevárez P, Munguía-Xóchihua JA, Segura-Correa JC and Leyva-Corona JC 2020. Risk factors associated with the seroprevalence of *paratuberculosis* in sheep flocks in the hot-arid region of Sonora, México. *Trop Anim Health Prod.* 52(3):1357–63.
- Mori M and Roest HJ 2018. Farming, Q fever and public health: Agricultural practices and beyond. *Arch Public Heal.* 76(1):1–9.
- Mpenda FN and Buza J 2014. Seroprevalence of *paratuberculosis* in goats and sheep in Arusha, Northern Tanzania. *Int J Sci Res.* 3(9):2319–7064.
- Nurkunasegran M, Kho K, Koh F, Tan P, Nizam Q, Ong B, Panchadcharam C, Mat Amin M, Abdul Majid N, Ramli R and Tay S 2017. Molecular Detection of *Coxiella burnetii* from Farm Animals and Ticks in Malaysia. *Trop Biomed.* 34(3):675–80.
- Nusinovici S, Frössling J, Widgren S, Beaudeau F and Lindberg A 2015. Q fever infection in dairy cattle herds: Increased risk with high wind speed and low precipitation. *Epidemiol Infect.* 143(15):3316–3326.
- Pourmahdi Borujeni M, Haji Hajikolaei MR, Ghorbanpoor M, Elhaei Sahar H, Bagheri S and Roveyshedzadeh S 2021. Comparison of *Mycobacterium avium* subsp. *paratuberculosis* infection in cattle, sheep and goats in the Khuzestan Province of Iran: Results of a preliminary survey. *Vet Med Sci* 7(5):1970–9.
- Rai SB, Kamaludin F, Ts C and Ck Y 2011. First Documented Zoonotic Case of Q Fever in Penang, Malaysia. *Outbreak Surveill Investig Rep.* 4(1):1–5.
- Raizman EA, Wells SJ, Jordan PA, DelGiudice GD and Bey RR 2005. *Mycobacterium avium* subsp. *paratuberculosis* from free-ranging deer and rabbits surrounding Minnesota dairy herds. *Can J Vet Res.* 69(1):32–8.
- Reyes-García R, Pérez-de-la-Lastra JM, Vicente J, Ruiz-Fons F, Garrido JM and Gortázar C 2008. Large-scale ELISA testing of Spanish red deer for *paratuberculosis*. *Vet Immunol Immunopathol.* 124(1–2):75–81.
- Rizzo F, Vitale N, Ballardini M, Borromeo V, Luzzago C, Chiavacci L and Mandola ML 2016. Q fever seroprevalence and risk factors in sheep and goats in northwest Italy. *Prev Vet Med.* 130:10–7.
- Rogan WJ and Gladen B 1978. Estimating Prevalence from the Results of a Screening Test. *Am J Epidemiol* 107(1):71–6.
- Ruiz-Fons F, Rodríguez Ó, Torina A, Naranjo V, Gortázar C and de la Fuente J 2008. Prevalence of *Coxiella burnetii* infection in wild and farmed ungulates. *Vet Microbiol.* 126(1–3):282–6.
- Sergeant ES 2018. Epitools Epidemiological Calculators. <https://epitools.ausvet.com.au/cited> 2023 Sep 15.
- Sharma S, Gautam A, Singh SV, Chaubey KK, Mehta R, Gupta S, Sharma M, Rose MK and Jain VK 2020. Prevalence of *Mycobacterium avium* subspecies *paratuberculosis* (MAP) infection in suspected diarrhoeic buffaloes and cattle reporting at Veterinary University in India. *Comp Immunol Microbiol Infect Dis.* 73(January):101533.
- Shin GW, Kim EJ, Lee HB and Cho HS 2014. The prevalence of *Coxiella burnetii* infection in wild Korean water deer, Korea. *J Vet Med Sci.* 76(7):1069–71.
- Sleeman JM, Manning EJB, Rohm JH, Sims JP, Sanchez S, Gerhold RW and Keel MK 2009. Johne’s Disease in a Free-Ranging White-tailed Deer from Virginia and Subsequent Surveillance for *Mycobacterium avium* subspecies *paratuberculosis*. *J Wildl Dis.* 45(1):201–6.
- Ssekitoleko J, Ojok L, Abd A, Wahed E, Erume J, Amanzada A, Eltayeb E, Eltom KH and Okuni JB 2021. *Mycobacterium avium* subsp. *paratuberculosis* Virulence: A Review. *Microorganisms.* 9:1–16.
- Stau A, Seelig B, Walter D, Schroeder C and Ganter M 2012. Seroprevalence of *Mycobacterium avium* subsp. *paratuberculosis* in small ruminants in Germany. *Small Rumin Res.* 105(1–3):361–5.
- Tryland M, Olsen I, Vikøren T, Handeland K, Arnemo JM, Tharaldsen J, Djønne B, Josefsen TD and Reitan LJ 2004. Serologic survey for antibodies against *Mycobacterium avium* subsp. *paratuberculosis* in free-ranging cervids from Norway. *J Wildl Dis.* 40(1):32–41.
- Ullah Q, Jamil T, Saqib M, Iqbal M and Neubauer H 2022. Q Fever-A Neglected Zoonosis. *Microorganisms.* 10(8):1–17.
- Voss L, Huaman J, Pacioni C, Tolpinrud A, Helbig K, Carvalho TG and Firestone SM 2023. Seroprevalence of *Coxiella burnetii* antibodies in wild deer populations in eastern Australia. *Aust Vet J.* 101(3):106–14.
- Whittington R, Donat K, Weber MF, Kelton D, Nielsen SS, Eisenberg S, et al. 2019. Control of

- paratuberculosis: Who, why and how. A review of 48 countries. BMC Vet Res. 15(1):1–29.
- Yangho J, Kim H, Heo I, Park Y, Kim S, Lee M and Choe N 2011. Seroprevalence of *Coxiella burnetii* in cattle and farm-raised deer in Korea. African J Microbiol Res. 5(24):4234–6.
- Yu Y, Zhang S, Xu G, Xu D, Zheng H, Li B, Shen K and Fu L 2022. Identification of *Mycobacterium avium* subspecies *paratuberculosis* in sheep farms in Bayannaoer, Inner Mongolia, China (short communication). BMC Vet Res. 18(1):1