



**EVALUATION OF DIFFERENT ANTHROPOMETRIC INDICES AND THEIR
ASSOCIATIONS WITH ELEVATED BLOOD PRESSURE AMONG
ADOLESCENTS IN SELANGOR, MALAYSIA**

By

KARTHIKAH RADAKRISHNAN

**Dissertation Submitted to Universiti Putra Malaysia, in Fulfilment of the
Requirements for the Degree of Master in Nutritional Sciences**

August 2022

FPSK (m) 2022 58

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



EVALUATION OF DIFFERENT ANTHROPOMETRIC INDICES AND THEIR ASSOCIATIONS WITH ELEVATED BLOOD PRESSURE AMONG ADOLESCENTS IN SELANGOR, MALAYSIA

By

KARTHIKAH RADAKRISHNAN

AUGUST 2022

Chair : Associate Professor Dr Gan Wan Ying, PhD

A trend of increasing prevalence of hypertension among children and adolescents was observed globally, including Malaysia. Early detection and prompt treatment of hypertension in children and adolescents helps in reducing associated risks and long term effect. Anthropometry indices have been used in several studies to predict hypertension. This cross-sectional study aims to compare the predictive ability of the 7 anthropometric indices: BMI-for-age z-score (BAZ), Waist circumference (WC), Waist-to-height ratio (WHtR), Conicity Index (C-Index), A Body Shape Index (ABSI), Body Roundness Index (BRI), and Triponderal Mass Index (TMI) in identifying elevated blood pressure in adolescents aged 13 to 17 years old in Selangor. Two schools in Selangor state was randomly selected using probability proportionate to size where a total of 934 respondents were recruited. Self-administered questionnaires was used to get data of sociodemographic background, physical activity and sleep quality. Weight, height, waist circumference and blood pressure were measured. Using the receiver operating characteristic curve and established procedures, the prediction power of anthropometric indicators was according to sex. A total 934 children (387 boys and 547 girls) were examined. The prevalence of elevated BP was 19.6%, hypertension stage 1 was 7.7% and hypertension stage 2 was 4.8%. Multiple logistic regression analysis upon controlling sex, age, ethnicity and physical activity level showed WC (AOR=1.274, 95% CI = 0.746, 2.176) and TMI (AOR=1.310, 95% CI = 1.219, 1.407) were significantly associated with elevated BP. Overall, AUCs of BMI-for-age (AUC= 0.752, CI 95%= 0.717, 0.787), WC (AUC=0.767, CI 95%= 0.735, 0.800), WHtR (AUC= 0.701, CI 95%= 0.664, 0.738), TMI (AUC= 0.731, CI 95%= 0.695, 0.767) and BRI (AUC= 0.701, CI 95%= 0.664, 0.738) indicated good predictive power in assessing elevated BP. Considering the results of AUC, sensitivity, specificity and Youden index, WC was considered as the best anthropometric indices in predicting high BP among adolescents in this study cut off values of 71.12 cm in males and 74.27 cm in

females. These results suggested that WC may be a helpful signal for assessing the risk of high blood pressure in routine screening for adolescents. Future studies are suggested to include dietary factors and stress factors to investigate the association with BP



ACKNOWLEDGEMENTS

First of all, I would like to thank my supervisor, Assoc. Prof. Dr. Gan Wan Ying for the patient guidance, encouragement, and advice she has provided throughout this period. I choose this moment to acknowledge her contribution gratefully.

Secondly, my sincere thanks go to Dr. Lim Poh Ying, for her assistance, expertise and suggestions that guided me to improve this research. Apart from that, I would also like to express my gratitude to all the lecturers of Faculty of Medicine & Health Sciences, who guided me in many areas and made my education journey memorable. Apart from that, special gratitude to all the respondents that participated in the questionnaire collected by Joyce Ying Hui Tee, that has contributed a major part in my dissertation.

Besides that, I would like to express my sincere gratitude to my parents who have been supporting me throughout my master program by offering moral support and endless prayer. Last but not least, I would like to thank those who contributed in my dissertation directly and indirectly. I would like to acknowledge their opinions and recommendation, which was crucial for the successful completion of this study.

Declaration by the Graduate Student

I hereby confirm that:

- This dissertation is my original work;
- quotations, illustrations and citations have been duly referenced;
- this dissertation has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the dissertation and the copyright of the dissertation are fully-owned by Universiti Putra Malaysia, as stipulated in the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from the supervisor and the office of the Deputy Vice-Chancellor (Research and innovation) before the dissertation is published in any written, printed or electronic form (including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials) as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the dissertation, and scholarly integrity is upheld in accordance with the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2015-2016) and the Universiti Putra Malaysia (Research) Rules 2012. The dissertation has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No.: _____

Declaration by Members of the Supervisory Committee

This is to confirm that:

- the research and the writing of this dissertation were done under our supervision;
- supervisory responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2015-2016) are adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: _____

Signature: _____
Name of Member of
Supervisory
Committee: _____

Signature: _____
Name of Member of
Supervisory
Committee: _____

Signature: _____
Name of Member of
Supervisory
Committee: _____

TABLE OF CONTENTS

	Page
ABSTRACT	i
ACKNOWLEDGEMENTS	iii
DECLARATION	iv
TABLE OF CONTENT	vi
LIST OF TABLES	viii
LIST OF FIGURES	ix
LIST OF ABBREVIATIONS	x
CHAPTER	
1 INTRODUCTION	1
1.1 Background	1
1.2 Problem Statement	2
1.3 Research Questions	3
1.4 Significance of Study	3
1.5 Research Objectives	4
1.5.1 General Objectives	4
1.5.2 Specific Objectives	4
1.6 Research Hypotheses	4
1.7 Conceptual Framework	5
2 LITERATURE REVIEW	6
2.1 Hypertension	6
2.1.1 Hypertension among adolescents	6
2.2 Risk factors of hypertension among adolescents	12
2.2.1 Sociodemographic factors	12
2.2.2 Obesity	13
2.2.3 Physical Activity	14
2.2.4 Sleep Quality	15
2.3 Anthropometric Indices	16
2.3.1 BMI-for-age z-score (BAZ)	16
2.3.2 Waist circumference (WC)	17
2.3.3 Waist-to-height ratio (WHtR)	17
2.3.4 Conicity Index (C-Index)	18
2.3.5 A Body Shape Index (ABSI)	18
2.3.6 Body Roundness Index (BRI)	19
2.3.7 Triponderal Mass Index (TMI)	20
3 METHODOLOGY	21
3.1 Study Design	21
3.2 Study Location	21
3.3 Study Respondents	21
3.4 Sample Size Determination	22
3.5 Sampling Design	23
3.6 Ethics Approval	23
3.7 Measures	23
3.7.1 Sociodemographic Background	23
3.7.2 Physical Activity Level	23

3.7.3	Sleep Quality	24
3.7.4	Blood Pressure	25
3.7.5	Anthropometric Measurements	25
3.8	Statistical Analysis	28
4	RESULTS	29
4.1	Sociodemographic background	29
4.2	Physical Activity Level	31
4.3	Sleep Quality	37
4.4	Anthropometry Measurements	41
4.5	Blood Pressure	43
4.6	Univariate Logistic Regression	44
4.7	Multiple logistic regression	46
4.8	Receiver Operating Curve and Cut Off Points	46
5	DISCUSSION	49
5.1	Prevalence of Elevated BP	49
5.2	Factors Associated with Elevated Blood Pressure Among Adolescents	49
5.3	Anthropometric Index and Elevated Blood Pressure	50
5.4	Strength and limitations	54
6	CONCLUSION & RECOMMENDATIONS	55
6.1	Conclusion	55
6.2	Recommendations	55
	REFERENCES/BIBLIOGRAPHY	57
	APPENDICES	75

LIST OF TABLES

Table		Page
2.1	Classification of clinic blood pressure levels in adults	8
2.2	Classification of children and adolescent blood pressure level	10
2.3	Prevalence of hypertension in adolescents globally	12
2.4	Prevalence of hypertension in adolescents in Malaysia	14
3.1	Inclusion and exclusion criteria	28
3.2	Physical activity level categories and PAQ-C scores	31
3.3	PSQI score and sleep quality indication	32
3.4	Classification of blood pressure for adolescents in this study	32
3.5	Classification of BMI-for-age z-score for adolescents	33
3.6	Classifications of WC-for-age for Malaysian adolescents	34
4.1	Sociodemographic background of the subjects of study	39
4.2	Distribution of respondents by items in the Physical Activity Questionnaire for Older Children (PAQ-C)	41
4.3	Physical Activity Questionnaire for Older Children (PAQ-C) score and physical activity (PA) level categorisation of respondents	44
4.4	Distribution of score by components in the Pittsburgh Sleep Quality Index (PSQI)	46
4.5	Global Pittsburgh Sleep Quality Index (PSQI) score categorisation of respondents	48
4.6	Anthropometry data of respondents	50
4.7	Blood pressure classification of respondents	51
4.8	Univariate logistic regression of factors associated with elevated blood pressure among adolescents	53
4.9	Multiple logistic regression results	54
4.10	Area under ROC curve (AUC), optimal cut-off values, sensitivities, specificities and Youden index of anthropometric indices in predicting high blood pressure according to sex	56

LIST OF FIGURES

Figure		Page
1.1	Conceptual framework of this study	7
3.1	G*Power sample calculation result	29



LIST OF ABBREVIATIONS

BP	Blood Pressure
WC	Waist Circumference
WHtR	Waist to height ratio
BAZ	BMI-for-age z-score
C-Index	Conicity Index
ABSI	A Body Shape Index
BRI	Body Roundness Index
TMI	Triponderal Mass Index
WHO	World Health Organization
BMI	Body mass index
PA	Physical activity
OR	Odds ratio
AOR	Adjusted odds ratio
NHMS	National Health Morbidity Survey
MOH	Ministry of Health Malaysia
NCD	Non-communicable diseases
SD	Standard deviation
CPG	Clinical Practice Guidelines
SPSS	Statistical Package for the Social Sciences
CI	Confidence Level
SBP	Systolic blood pressure
BDP	Diastolic blood pressure

CHAPTER 1

INTRODUCTION

1.1 Background

Hypertension is a rising concern globally (Song et al., 2019). The biggest cause of morbidity and mortality in the world today is hypertension, which accounts for 10.4 million deaths per year (Stanaway et al., 2018). A systematic review and meta-analysis reported a trend of increasing prevalence of hypertension among children and adolescents was observed globally during the past two decades, with a relative increment of 75% to 79% from 2000 to 2015 (Song et al., 2019). Hypertension that starts in childhood would have high chances to continue into adulthood, but childhood hypertension is largely undiagnosed (AHA, 2003).

Obesity has been shown as a contributing factor and to increase prevalence of several metabolic diseases, including hypertension (Jiang et al., 2016). Obesity is one of the main public health problems worldwide (Powell-Wiley et al., 2021). This is impacting children and adolescents as well where globally, the number of obese children and adolescents has risen tenfold in the past forty years (NCD-RisC, 2017). Additionally, WHO has predicted that by 2022, a greater number of children and adolescents will be affected by obesity than by underweight and this is alarming (NCD-RisC, 2017).

The relationship between hypertension and obesity has been extensively studied in both adults and children across both sexes (Shariq & Mckenzie, 2020). There is a strong link that higher body mass index (BMI) causes higher blood pressure, in which a previous study showed that each 10 BMI unit increase was related with a 10 mmHg rise in systolic blood pressure and a 3 mmHg increase in diastolic blood pressure (Kelly et al., 2015). Similarly, this is supported by another study in China where it was found that among children and adolescents, being overweight or obese significantly raised the risk of developing hypertension (Dong et al., 2019).

South Asian children and adolescents have been shown to have a higher body fat percentage at similar BMI compared with Caucasians placing them at greater risk of metabolic diseases, including hypertension (Eyre et al., 2017; Heo et al., 2020). This shows that Asian are more at risk compared to Caucasian in developing hypertension. Among adolescents, the prevalence of hypertension is quite alarming, especially in Asian countries such as India (7.6%), China (13.45%), and Indonesia (8%) (Daniel et al., 2020; Kurnianto et al., 2020a; Yang et al., 2021a). In Malaysia, Liew et al. (2019) found that the prevalence of hypertension among adolescents was 24.5% with the highest being recorded among Malays (28.7%). This was comparatively higher than a previous study conducted in 2011, where the prevalence was found to be 11.7% (Rampal et al., 2011).

Reducing related risks and long-term effects of hypertension in children and adolescents requires early detection and prompt treatment. Among the early detection methods, anthropometry indices have been used in several studies to predict hypertension (Nkwana et al., 2021; Ononamadu et al., 2017; Tee et al., 2020). For many years, the primary metrics used to evaluate a child's growth have been their height and weight (Erdal et al., 2020). This is referring to the conventional BMI that is still promulgated as the main epidemiological measure of obesity (Nuttall, 2015). In adolescents, BMI-for-age z-score (BAZ) is more suitable to be applied (Freedman et al., 2009). However, the disadvantage of BMI or BAZ is that it cannot reflect fat distribution and adiposity level in an individual (Nevill et al., 2021).

Alternatively, abdominal obesity indices, which includes waist circumference (WC), waist-to-height ratio (WHtR) and conicity index (C-Index) have been proposed, because they alter the BMI's limiting factor (Lee et al., 2015; Lo et al., 2016). Recent review showed that WHtR has the strongest association with adiposity, in which it is a better predictor of hypertension than BMI and WC (Hwaung et al., 2020). Conicity index was also positively associated with hypertension in the adult population (Nkwana et al., 2021).

Some new anthropometric indices have been proposed for screening hypertension, including body roundness index (BRI), a body shape index (ABSI) and Triponderal mass index (TMI). Body Roundness Index (BRI) and A Body Shape Index (ABSI) were proposed for that they can predict the distributions of body fat and were associated with hypertension (Chang et al., 2016; Nkwana et al., 2021). A recent study has shown that TMI can be used to characterize adolescents with overweight and hypertensive (Malavazos et al., 2021). However, it is still challenging to identify hypertensive risk factors and to determine which anthropometric measurements are appropriate at various stages of development in childhood and adolescence due to the mixed findings.

1.2 Problem Statement

Many nations and ethnic groups have explored the connection between anthropometric indicators of obesity and hypertension (Eyre et al., 2017; Ononamadu et al., 2017; Sardinha et al., 2016; Tee et al., 2020). A study conducted among Indian children found that BMI, WC, WHtR, triceps skin fold thickness and wrist measurements has equal sensitivity and specificity in detection of hypertension (Kajale et al., 2014). However, a study among Nigerian secondary school children found that WC and waist to hip ratio were significant predictors of hypertension compared to BMI (Uwaezuoke et al., 2016). Comparatively, Rodea-Montero et al. (2014) found that WHtR was the best comparing to BMI and WC in a study among Mexican adolescents. The findings of several studies indicate that the appropriate anthropometric index for predicting hypertension is still disputed and inconclusive.

In Malaysia, there are few and conflicting results from research that evaluated how well anthropometric indices predict the likelihood that adolescents may develop hypertension. Tee et al. (2020) evaluated the anthropometric

indices for the screening of hypertension among adolescents; however, only four indices (BMI, WC, WHtR, ABSI) were assessed, in which results showed that WHtR was the best index to predict hypertension. Nevertheless, another study done by Cheah et al. (2018) showed that BMI and WC were the best index in predicting hypertension. Therefore, the findings are contradicting and the best predictive anthropometric index is still inconclusive.

Cut off values are important in the predicting power of the indices. According to studies from many nations and ethnic groups, certain anthropometric indices and the associated cut-off values for identifying high blood pressure are superior than others (Nkwana et al., 2021; Ononamadu et al., 2017; Tee et al., 2020). Multiple cut-off values and the use of different anthropometric measures to predict high blood pressure may be necessary due to ethnic heterogeneity among populations from different locations. It is important to generate a cut off value that is suitable for Malaysian population since our body composition and fat distribution may differ from the existing optimal cut off values that has been established in other studies from other countries (Eyre et al., 2017; Heo et al., 2020) .

Overall, most of the studies on anthropometric indices and hypertension were conducted in the adult age group (Andrade et al., 2016; Bertoli et al., 2017; Calderón-García et al., 2021). The association among adolescents is not fully understood. Therefore, this study aimed to compare the predictive ability of the 7 anthropometric indices: BAZ, WC, WHtR, C-index, TMI, BRI, and ABSI in identifying elevated blood pressure among adolescents from age 13 to 17 years old.

1.3 Research Questions

This study addressed a few important research questions as follows:

1. What are the associations between BAZ, WC, WHtR, C-index, TMI, BRI, and ABSI with high blood pressure in adolescents aged 13 to 17 years old?
2. Which of these anthropometric indices (BAZ, WC, WHtR, C-index, TMI, BRI, and ABSI) are the best predictors of high blood pressure in adolescents aged 13 to 17 years old?
3. What is the optimal cut off value for each anthropometric indices in the prediction of high BP among adolescents aged 13 to 17 years old?

1.4 Significance of the Study

As hypertension is increasing globally and in Malaysia, early detection of hypertension is important. Anthropometry indices are simple methods which can be obtained using basic measurements that can be easily measured even in the school settings.

By identifying the best anthropometric measure and the cut-off values, further action can be taken to implement the anthropometric measure in schools and hospitals as a predictive tool. The anthropometric index can be used as a primary screening to identify the high risk individuals in developing hypertension. Anthropometric index is a cheaper and easier way to determine the risk group as it does not need any biochemical values.

Findings of this study can be used by healthcare professionals to assess hypertension risk easily, subsequently reducing the risk in adolescents by targeting the anthropometric indices. This research will advance knowledge in the area and provide data for subsequent intervention studies on adolescents. The information obtained from this study can be used by policy makers to plan preventive measures by implementing the anthropometric index in predicting hypertension among adolescents.

1.5 Research Objectives

1.5.1 General Objective

To compare the predictive ability of the 7 anthropometric indices: BAZ, WC, WHtR, C-index, TMI, BRI, and ABSI in identifying elevated blood pressure in adolescents aged 13 to 17 years old in Selangor.

1.5.2 Specific objectives

1. To assess BAZ, WC, WHtR, C-index, TMI, BRI, ABSI and blood pressure levels in adolescents.
2. To determine the associations between the 7 anthropometric indices (BAZ, WC, WHtR, C-index, TMI, BRI, and ABSI) with blood pressure among adolescents after adjustment of covariates.
3. To compare the performance of different anthropometric indices in predicting elevated blood pressure in adolescents.
4. To identify the optimal sex specific cut off values for anthropometric indices in predicting elevated blood pressure in adolescents.

1.6 Research Hypotheses

1. Obesity indices are associated with elevated blood pressure among adolescents after adjustment for covariates.
2. There are differences in the performance of different anthropometric indices in predicting elevated BP in adolescents.

1.7 Conceptual Framework

Figure 1.1 explains the framework of this study. Anthropometry indices (BAZ, WC, WHtR, C-index, TMI, BRI and ABSI) are the independent variables and blood pressure is the dependant variable. The confounding variables in this study would be the lifestyle factors including physical activity level (PAL), sleep quality, fast food consumption frequency and sociodemographic factors, which includes age, sex, ethnicity, education, and socioeconomic status. The anthropometry indices are associated with blood pressure and are used as a predictive tool in predicting hypertension.




Figure 1.1: Conceptual framework of this study

REFERENCES

- Abbaszadeh, F., Sarafranz, N., Atrian, M. K., Sadat, Z., Bagheri, A., & Moravveji, A. (2017). Anthropometric indices in the prediction of hypertension in female adolescents. *Iranian Red Crescent Medical Journal*, 19(11). <https://doi.org/10.5812/ircmj.14591>
- Abdullah, N. F., Teo, P. S., & Foo, L. H. (2016). Ethnic differences in the food intake patterns and its associated factors of adolescents in Kelantan, Malaysia. *Nutrients*, 8(9), 551. <https://doi.org/10.3390/nu8090551>
- Almahmoud, O. H., Arabiat, D. H., & Saleh, M. Y. (2022). Systematic review and meta-analysis: Prevalence of hypertension among adolescents in the Arab countries. *Journal of Pediatric Nursing*, 65, (e72-e79) <https://doi.org/10.1016/j.pedn.2022.03.002>
- Al-Sadat, N., Majid, H. A., Sim, P. Y., Su, T. T., Dahlui, M., Bakar, M. F. A., Dzaki, N., Norbaya, S., Murray, L., Cantwell, M. M., & Jalaludin, M. Y. (2016). Vitamin D deficiency in Malaysian adolescents aged 13 years: Findings from the Malaysian Health and Adolescents Longitudinal Research Team study (MyHeARTs). *BMJ Open*, 6(8), e010689. <https://doi.org/10.1136/bmjopen-2015-010689>
- Andrade, M. D., Freitas, M. C. P. D., Sakumoto, A. M., Pappiani, C., Andrade, S. C. D., Vieira, V. L., & Damasceno, N. R. T. (2016). Association of the conicity index with diabetes and hypertension in Brazilian women. *Archives of Endocrinology and Metabolism*, 60, 436-442. <https://doi.org/10.1590/2359-3997000000187>
- Anujuo, K. O., Vrijkotte, T. G., Stronks, K., Jean-Louis, G., & Agyemang, C. O. (2016). Ethnic differences in sleep duration at 5 years, and its relationship with overweight and blood pressure. *The European Journal of Public Health*, 26(6), 1001-1006. <https://doi.org/10.1093/eurpub/ckw084>
- Anyaegbu, E. I., & Dharnidharka, V. R. (2014). Hypertension in the teenager. *Pediatric Clinics of North America*, 61(1), 131-151. <https://doi.org/10.1016/j.pcl.2013.09.011>
- Ashley-Martin, J., Ensenuer, R., Maguire, B., & Kuhle, S. (2019). Predicting cardiometabolic markers in children using tri-ponderal mass index: A cross-sectional study. *Archives of Disease in Childhood*, 104(6), 577-582. <https://doi.org/10.1136/archdischild-2018-316028>
- Azemati, B., Kelishadi, R., Ahadi, Z., Shafiee, G., Taheri, M. Z., Ziaodini, H., Qorbani, M., & Heshmat, R. (2020). Association between junk food consumption and cardiometabolic risk factors in a national sample of Iranian children and adolescents population: the CASPIAN-V study. *Eating and Weight Disorders*, 25(2), 329-335. <https://doi.org/10.1007/s40519-018-0591-1>
- Bahadoran, Z., Mirmiran, P., & Azizi, F. (2015). Fast food pattern and cardiometabolic disorders: a review of current studies. *Health Promotion Perspectives*, 5(4), 231.

- Batsky, D. (2012). Blood pressure variability, prehypertension, and hypertension in adolescents. *Adolescent Health, Medicine and Therapeutics*, 43. <https://doi.org/10.2147/ahmt.s15942>
- Beck, C. C., Lopes, A. da S., & Pitanga, F. J. G. (2011). Anthropometric indicators as predictors of high blood pressure in adolescents. *Arquivos Brasileiros de Cardiologia*, 96(2), 126–133.
- Bentham, J., di Cesare, M., Bilano, V., Bixby, H., Zhou, B., Stevens, G. A., Riley, L. M., Taddei, C., Hajifathalian, K., Lu, Y., Savin, S., Cowan, M. J., Paciorek, C. J., Chirita-Emandi, A., Hayes, A. J., Katz, J., Kelishadi, R., Kengne, A. P., Khang, Y. H., ... Cisneros, J. Z. (2017). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *The Lancet*, 390(10113), 2627–2642. [https://doi.org/10.1016/S0140-6736\(17\)32129-3](https://doi.org/10.1016/S0140-6736(17)32129-3)
- Bertoli, S., Leone, A., Krakauer, N.Y., Bedogni, G., Vanzulli, A., Redaelli, V.I., De Amicis, R., Vignati, L., Krakauer, J.C. and Battezzati, A. (2017). Association of Body Shape Index (ABSI) with cardio-metabolic risk factors: A cross-sectional study of 6081 Caucasian adults. *PloS One*, 12(9), e0185013. <https://doi.org/10.1371/journal.pone.0185013>
- Bijlsma, M. W., Blufpand, H. N., Kaspers, G. J., & Bökenkamp, A. (2014). Why pediatricians fail to diagnose hypertension: A multicenter survey. *The Journal of Pediatrics*, 164(1), 173-177. <https://doi.org/10.1016/j.jpeds.2013.08.066>
- Bloch, K. V., Klein, C. H., Szklo, M., Kuschner, M. C. C., de Azevedo Abreu, G., Barufaldi, L. A., da Veiga, G. V., Schaan, B., da Silva, T. L. N., de Vasconcellos, M. T. L., Moraes, A. J. P., Borges, A. L., de Oliveira, A. M. A., Tavares, B. M., de Oliveira, C. L., de Freitas Cunha, C., Giannini, D. T., Belfort, D. R., Santos, E. L., ... Goldberg, T. B. L. (2016). ERICA: Prevalences of hypertension and obesity in Brazilian adolescents. *Revista de Saude Publica*, 50, 1s–12s. <https://doi.org/10.1590/S01518-8787.2016050006685>
- Brown, I. J., Stamler, J., van Horn, L., Robertson, C. E., Chan, Q., Dyer, A. R., Huang, C. C., Rodriguez, B. L., Zhao, L., Daviglius, M. L., Ueshima, H., & Elliott, P. (2011). Sugar-sweetened beverage, sugar intake of individuals, and their blood pressure: International study of macro/micronutrients and blood pressure. *Hypertension*, 57(4), 695–701. <https://doi.org/10.1161/HYPERTENSIONAHA.110.165456>
- Browning, L. M., Hsieh, S. D., & Ashwell, M. (2010). A systematic review of waist-to-height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 05 could be a suitable global boundary value. *Nutrition Research Reviews*, 23(2), 247–269. <https://doi.org/10.1017/S0954422410000144>
- Butte, N. F., Watson, K. B., Ridley, K., Zakeri, I. F., McMurray, R. G., Pfeiffer, K. A., Crouter, S. E., Herrmann, S. D., Bassett, D. R., Long, A., Berhane, Z., Trost, S. G., Ainsworth, B. E., Berrigan, D., & Fulton, J. E. (2018). A youth compendium

of physical activities: Activity codes and metabolic intensities. *Medicine and Science in Sports and Exercise*, 50(2), 246–256. <https://doi.org/10.1249/MSS.0000000000001430>

- Cai, S., Dong, J., Cheng, B., Zhang, A., Sun, J., Li, M., Su, Y., Bao, Q., Zhu, P., & Wang, S. (2022). Relationship of a new anthropometric index with left ventricular hypertrophy in hypertensive patients among the Han Chinese. *BMC Cardiovascular Disorders*, 22(1), 16. <https://doi.org/10.1186/s12872-022-02463-6>
- Calderón-García, J. F., Roncero-Martín, R., Rico-Martín, S., de Nicolás-Jiménez, J. M., López-Espuela, F., Santano-Mogena, E., Alfageme-García, P., & Muñoz-Torrero, J. F. S. (2021). Effectiveness of body roundness index (BRI) and a body shape index (ABSI) in predicting hypertension: A systematic review and meta-analysis of observational studies. *International Journal of Environmental Research and Public Health*, 18(21), 11607. <https://doi.org/10.3390/ijerph182111607>
- Çam, H. H., & Ustuner Top, F. (2021). Prevalence of Hypertension and Its Association with Body Mass Index and Waist Circumference Among Adolescents in Turkey: A Cross-Sectional Study. *Journal of Pediatric Nursing*, 57, e29–e33. <https://doi.org/10.1016/j.pedn.2020.09.017>
- Capers, P. L., Fobian, A. D., Kaiser, K. A., Borah, R., & Allison, D. B. (2015). A systematic review and meta-analysis of randomized controlled trials of the impact of sleep duration on adiposity and components of energy balance. *Obesity Reviews*, 16(9), 771–782. <https://doi.org/10.1111/obr.12296>
- Carpenter, C. L., Yan, E., Chen, S., Hong, K., Arechiga, A., Kim, W. S., Deng, M., Li, Z., & Heber, D. (2013). Body fat and body-mass index among a multiethnic sample of college-age men and women. *Journal of Obesity*, 2013. <https://doi.org/10.1155/2013/790654>
- Carson, V., Rinaldi, R.L., Torrance, B., Maximova, K., Ball, G.D.C., Majumdar, S.R., Plotnikoff, R.C., Veugelers, P., Boule, N.G., Wozny, P. and McCargar, L. (2014). Vigorous physical activity and longitudinal associations with cardiometabolic risk factors in youth. *International Journal of Obesity*, 38(1), 16-21. <https://doi.org/10.1038/ijo.2013.135>
- Chang, Y., Guo, X., Guo, L., Li, Z., Li, Y., & Sun, Y. (2016). The feasibility of two new anthropometric indices to identify hypertension in rural China A cross-sectional study. *Medicine (United States)*, 95(44), e5301. <https://doi.org/10.1097/MD.0000000000005301>
- Chang, Y., Guo, X., Li, T., Li, S., Guo, J., & Sun, Y. (2015). A body shape index and body roundness index: two new body indices to identify left ventricular hypertrophy among rural populations in northeast China. *BMC Public Health*, 15, 794. <https://doi.org/10.1186/s12889-015-2150-2>

- Cheah, W. L., Chang, C. T., Hazmi, H., & Kho, G. W. F. (2018a). Using Anthropometric Indicator to Identify Hypertension in Adolescents: A Study in Sarawak, Malaysia. *International Journal of Hypertension*, 2018. <https://doi.org/10.1155/2018/6736251>
- Chen, Z., Smith, M., Du, H., Guo, Y., Clarke, R., Bian, Z., Collins, R., Chen, J., Qian, Y., Wang, X. and Chen, X. (2015). Blood pressure in relation to general and central adiposity among 500 000 adult Chinese men and women. *International Journal of Epidemiology*, 44(4), 1305-1319. <https://doi.org/10.1093/ije/dyv012>
- Couch, S. C., & Daniels, S. R. (2013). Non-pharmacologic treatment of pediatric hypertension. *Pediatric Hypertension* (pp. 529-537). Humana Press, Totowa, NJ.
- Cristina Beck, C., da Silva Lopes, A., & José Gondim Pitanga, F. (2011). Anthropometric Indicators as Predictors of High Blood Pressure in Adolescents.
- Crocker, P. R., Bailey, D. A., Faulkner, R. A., Kowalski, K. C., & McGrath, R. (1997). Measuring general levels of physical activity: preliminary evidence for the Physical Activity Questionnaire for Older Children. *Medicine and Science in Sports and Exercise*, 29(10), 1344-1349
- Cui, Y., Zhang, F., Wang, H., Zhao, L., Song, R., Han, M., & Shen, X. (2022). Temporal Associations between Tri-Ponderal Mass Index and Blood Pressure in Chinese Children: A Cross-Lag Analysis. *Nutrients*, 14(9). <https://doi.org/10.3390/nu14091783>
- Cutler, G. J., Flood, A., Hannan, P. J., Slavin, J. L., & Neumark-Sztainer, D. (2012). Association between major patterns of dietary intake and weight status in adolescents. *British Journal of Nutrition*, 108(2), 349-356.
- Dan, S. P., MT, M. N., & Zalilah, M. S. (2007). Sex and ethnic differentials in physical activity levels of adolescents in Kuantan. *Malaysian Journal of Nutrition*, 13(2), 109-120.
- Daniel, R. A., Haldar, P., Prasad, M., Kant, S., Krishnan, A., Gupta, S. K., & Kumar, R. (2020). Prevalence of hypertension among adolescents (10-19 years) in India: A systematic review and meta-analysis of cross-sectional studies. *PLoS ONE*, 15(10), e0239929. <https://doi.org/10.1371/journal.pone.0239929>
- de Boer, E. C., de Rooij, S. R., Olthof, M. R., & Vrijkotte, T. G. (2018). Sugar-sweetened beverages intake is associated with blood pressure and sympathetic nervous system activation in children. *Clinical Nutrition ESPEN*, 28, 232-235. <https://doi.org/10.1016/j.clnesp.2018.08.020>.
- de Lorenzo, A., Romano, L., di Renzo, L., Gualtieri, P., Salimei, C., Carrano, E., Rampello, T., & de Miranda, R. C. (2019). Triponderal mass index rather than body mass index: An indicator of high adiposity in Italian children and adolescents. *Nutrition*, 60(1), 41–47. <https://doi.org/10.1016/j.nut.2018.09.007>

- de Onis, M., Martínez-Costa, C., Núñez, F., Nguefack-Tsague, G., Montal, A., & Brines, J. (2013). Association between WHO cut-offs for childhood overweight and obesity and cardiometabolic risk. *Public Health Nutrition*, 16(4), 625–630. <https://doi.org/10.1017/S1368980012004776>
- Dehghan, P., & Abbasalizad Farhangi, M. (2020). Dietary acid load, blood pressure, fasting blood sugar and biomarkers of insulin resistance among adults: Findings from an updated systematic review and meta-analysis. *International Journal of Clinical Practice*, 74(4), e13471. <https://doi.org/10.1111/ijcp.13471>
- Derks, I. P., Kocevskaja, D., Jaddoe, V. W., Franco, O. H., Wake, M., Tiemeier, H., & Jansen, P. W. (2017). Longitudinal associations of sleep duration in infancy and early childhood with body composition and cardiometabolic health at the age of 6 years: the generation R study. *Childhood Obesity*, 13(5), 400-408. <https://doi.org/10.1089/chi.2016.0341>
- Devonshire, A. L., Hager, E. R., Black, M. M., Diener-West, M., Tilton, N., & Snitker, S. (2016). Elevated blood pressure in adolescent girls: Correlation to body size and composition. *BMC Public Health*, 16(1). <https://doi.org/10.1186/s12889-016-2717-6>
- Dong, Y., Jan, C., Zou, Z., Dong, B., Wang, Z., Yang, Z., Li, Y., Wen, B., Ma, Y., Song, Y., Ma, J., Sawyer, S. M., & Patton, G. C. (2019). Effect of overweight and obesity on high blood pressure in Chinese children and adolescents. *Obesity*, 27(9), 1503–1512. <https://doi.org/10.1002/oby.22562>
- Dulskiene, V., Kuciene, R., Medzioniene, J., & Benetis, R. (2014). Association between obesity and high blood pressure among Lithuanian adolescents: A cross-sectional study. *Italian Journal of Pediatrics*, 40(5), 102. <https://doi.org/10.1186/s13052-014-0102-6>
- Duncan, M. J., Mota, J., Vale, S., Santos, M. P., & Ribeiro, J. C. (2013). Associations between body mass index, waist circumference and body shape index with resting blood pressure in Portuguese adolescents. *Annals of Human Biology*, 40(2), 163-167.
- Duren, D. L., Sherwood, R. J., Czerwinski, S. A., Lee, M., Choh, A. C., Siervogel, R. M., & Chumlea, W. C. (2008). Body composition methods: Comparisons and interpretation. *Journal of Diabetes Science and Technology*, 2(6), 1139-1146.
- Erdal, İ., Yalçın, S. S., Aksan, A., Gençal, D., & Kanbur, N. (2020). How useful are anthropometric measurements as predictive markers for elevated blood pressure in adolescents in different gender? *Journal of Pediatric Endocrinology and Metabolism*, 33(9), 1203–1211. <https://doi.org/10.1515/jpem-2020-0175>
- Ewald, D. R., Bond, S. H., & Haldeman, L. A. (2017). Hypertension in low-income adolescents. *Global Pediatric Health*, 4, 2333794X17741819. <https://doi.org/10.1177/2333794X17741819>

- Eyre, E. L. J., Duncan, M. J., & Nevill, A. (2017). South Asian children have increased body fat in comparison to white children at the same body mass index. *Children*, 4(11), 102. <https://doi.org/10.3390/children4110102>
- Ezeudu, C. E., Chukwuka, J. O., Ebenebe, J. C., Igwe, W. C., & Egbuonu, I. (2018). Hypertension and prehypertension among adolescents attending secondary schools in urban area of South-east, Nigeria. *Pan African Medical Journal*, 31, 145 <https://doi.org/10.11604/pamj.2018.31.145.15994>
- Fagard, R. H., & Cornelissen, V. A. (2007). Effect of exercise on blood pressure control in hypertensive patients. *European Journal of Preventive Cardiology*, 14(1), 12–17.
- Farah, N. M. F., Yee, T. S., & Rasdi, H. F. M. (2019). Self-reported sleep quality using the Malay version of the Pittsburgh Sleep Quality Index (PSQI-M) in Malaysian adults. *International Journal of Environmental Research and Public Health*, 16(23), 4750. <https://doi.org/10.3390/ijerph16234750>
- Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using G*Power 3.1: Tests for correlation and regression analyses. *Behavior Research Methods*, 41, 1149–1160.
- Finkelstein, A., Havakuk, O., Steinvil, A., Arbel, Y., Halkin, A., Bazan, S., Abramowitz, Y., Keren, G., & Banai, S. (2013). Gender differences and clinical outcome in patients undergoing trans-femoral aortic valve implantation. *International Journal of Cardiology*, 168(5), 4854–4855. <https://doi.org/10.1016/j.ijcard.2013.07.066>
- Finkelstein, A., Havakuk, O., Steinvil, A., Arbel, Y., Halkin, A., Bazan, S., Abramowitz, Y., Keren, G., & Banai, S. (2013). Gender differences and clinical outcome in patients undergoing trans-femoral aortic valve implantation. *International Journal of Cardiology*, 168(5), 4854–4855. <https://doi.org/10.1016/j.ijcard.2013.07.066>
- Fishman, B., Zloof, Y., Orr, O., Tsur, A. M., Furer, A., Omer Gilon, M., Chodick, G., Leiba, A., Derazne, E., Tzur, D., Afek, A., Grossman, E., & Twig, G. (2021). The opposing trends of body mass index and blood pressure during 1977–2020; nationwide registry of 2.8 million male and female adolescents. *Cardiovascular Diabetology*, 20(1), 242. <https://doi.org/10.1186/s12933-021-01433-0>
- Fluss, R., Faraggi, D., & Reiser, B. (2005). Estimation of the Youden Index and its associated cutoff point. *Biometrical Journal*, 47(4), 458–472. <https://doi.org/10.1002/bimj.200410135>
- Flynn, J. T., Kaelber, D. C., & Baker-Smith, C. M. (2017). Clinical practice guideline for screening and management of high blood pressure in children and adolescents. *Pediatrics*, 140(3), e20171904. <https://doi.org/10.1542/peds.2017-1904>
- Fobian, A. D., Elliott, L., & Louie, T. (2018). A systematic review of sleep, hypertension, and cardiovascular risk in children and adolescents. *Current hypertension reports*, 20(5), 1–11.

- Freedman, D. S., Goodman, A., Contreras, O. A., Das Mahapatra, P., Srinivasan, S. R., & Berenson, G. S. (2012). Secular trends in BMI and blood pressure among children and adolescents: The Bogalusa Heart Study. *Pediatrics*, *130*(1). <https://doi.org/10.1542/peds.2011-3302>
- Freedman, D. S., Wang, J., Thornton, J. C., Mei, Z., Sopher, A. B., Pierson, R. N., Dietz, W. H., & Horlick, M. (2009). Classification of Body Fatness by Body Mass Index-for-Age Categories Among Children. In *Arch Pediatr Adolesc Med* (Vol. 163, Issue 9).
- Gangwisch, J. E. (2014). A review of evidence for the link between sleep duration and hypertension. *American Journal of Hypertension*, *27*(10), 1235–1242. <https://doi.org/10.1093/ajh/hpu071>
- George, C., Goedecke, J. H., Crowther, N. J., Jaff, N. G., Kengne, A. P., Norris, S. A., & Micklesfield, L. K. (2016). The role of body fat and fat distribution in hypertension risk in urban Black South African women. *PLoS One*, *11*(5), e0154894. <https://doi.org/10.1371/journal.pone.0154894>
- Goel, M., Pal, P., Agrawal, A., & Ashok, C. (2016). Relationship of body mass index and other life style factors with hypertension in adolescents. *Annals of Pediatric Cardiology*, *9*(1), 29–34. <https://doi.org/10.4103/0974-2069.171393>
- Gröber-Grätz, D., Widhalm, K., De Zwaan, M., Reinehr, T., Blüher, S., Schwab, K.O., Wiegand, S. and Holl, R.W. (2013). Body mass index or waist circumference: which is the better predictor for hypertension and dyslipidemia in overweight/obese children and adolescents? Association of cardiovascular risk related to body mass index or waist circumference. *Hormone Research in Paediatrics*, *80*(3), 170-178.
- Gui, Z. H., Zhu, Y. N., Cai, L., Sun, F. H., Ma, Y. H., Jing, J., & Chen, Y. J. (2017). Sugar-sweetened beverage consumption and risks of obesity and hypertension in Chinese children and adolescents: A national cross-sectional analysis. *Nutrients*, *9*(12), 1302. <https://doi.org/10.3390/nu9121302>
- Hall JE, do Carmo JM, da Silva AA, Wang Z, Hall ME. Obesity -induced hypertension: interaction of neurohumoral and renal mechanisms. *Circ Res* 2015; 116: 991 - 1006
- Hardy, S. T., Sakhuja, S., Jaeger, B. C., Urbina, E. M., Suglia, S. F., Feig, D. I., & Muntner, P. (2021). Trends in blood pressure and hypertension among US children and adolescents, 1999-2018. *JAMA Network Open*, *4*(4), e213917. <https://doi.org/10.1001/jamanetworkopen.2021.3917>
- He, H., Pan, L., Du, J., Jin, Y., Jia, P., & Shan, G. (2022). Body mass index was linked with multi-cardiometabolic abnormalities in Chinese children and adolescents: A community-based survey. *BMC Pediatrics*, *22*(1), 33. <https://doi.org/10.1186/s12887-021-03092-2>

- Heo, S., Kwon, S., Lee, Y. M., Shin, J. Y., & Lee, D. H. (2020). Comparison of trends in blood pressure and the prevalence of obesity among Korean and American adolescents: A 12-years cross-sectional study. *Journal of Preventive Medicine and Public Health*, 53(1), 45–55. <https://doi.org/10.3961/jpmp.19.198>
- Hosmer D. W., & Lemeshow, S.(2000). *Applied logistic regression* (Vol. 398). John Wiley & Sons.
- Hsu, C. Y., Lin, R. H., Lin, Y. C., Chen, J. Y., Li, W. C., Lee, L. A., Liu, K. H., & Chuang, H. H. (2020). Are body composition parameters better than conventional anthropometric measures in predicting pediatric hypertension? *International Journal of Environmental Research and Public Health*, 17(16), 5771. <https://doi.org/10.3390/ijerph17165771>
- Hwaung, P., Heo, M., Kennedy, S., Hong, S., Thomas, D. M., Shepherd, J., & Heymsfield, S. B. (2020). Optimum waist circumference-height indices for evaluating adult adiposity: An analytic review. *Obesity Reviews*, 21(1), e12947. <https://doi.org/10.1111/obr.12947>
- Jackson, S. L., Zhang, Z., Wiltz, J. L., Loustalot, F., Ritchey, M. D., Goodman, A. B., & Yang, Q. (2017). *Morbidity and Mortality Weekly Report Hypertension Among Youths-United States, 2001-2016*. <http://hyper.ahajournals.org/content/early/2017/11/10/>
- Ji, M., Zhang, S., & An, R. (2018). Effectiveness of A Body Shape Index (ABSI) in predicting chronic diseases and mortality: A systematic review and meta-analysis. *Obesity Reviews*, 19(5), 737-759. <https://doi.org/10.1111/obr.12666>
- Jiang, S. Z., Lu, W., Zong, X. F., Ruan, H. Y., & Liu, Y. (2016). Obesity and hypertension. *Experimental and Therapeutic Medicine*, 12(4), 2395–2399. <https://doi.org/10.3892/etm.2016.3667>
- Jiang, Y., Dou, Y.L., Xiong, F., Zhang, L., Zhu, G.H., Wu, T., Zhang, Y. and Yan, W.L. (2018). Waist-to-height ratio remains an accurate and practical way of identifying cardiometabolic risks in children and adolescents. *Acta Paediatrica*, 107(9), 1629-1634.
- Kaczmarek, M., Stawińska-Witoszyńska, B., Krzyżaniak, A., Krzywińska-Wiewiorowska, M., & Sivińska, A. (2015). Who is at higher risk of hypertension? Socioeconomic status differences in blood pressure among Polish adolescents: a population-based ADOPLNOR study. *European Journal of Pediatrics*, 174(11), 1461–1473. <https://doi.org/10.1007/s00431-015-2554-0>
- Kajale, N. A., Khadilkar, A. v, Chipplonkar, S. A., & Khadilkar, V. v. (2014). Body Fat Indices for Identifying Risk of Hypertension in Indian Children. In *INDIAN PEDIATRICS* (Vol. 555).
- Kar, S., & Khandelwal, B. (2015). Fast foods and physical inactivity are risk factors for obesity and hypertension among adolescent school children in east district of Sikkim, India. *Journal of natural science, biology, and medicine*, 6(2), 356.

- Katzmarzyk, P.T., Barreira, T.V., Broyles, S.T., Chaput, J.P., Fogelholm, M., Hu, G., Kuriyan, R., Kurpad, A., Lambert, E.V., Maher, C. and Maia, J. (2015). Association between body mass index and body fat in 9–11-year-old children from countries spanning a range of human development. *International journal of obesity supplements*, 5(2), S43-S46.
- Kelly RK, Magnussen CG, Sabin MA, Cheung M, Juonala M. Development of hypertension in overweight adolescents: a review. *Adolesc Health Med Ther* 2015; 6: 171 -87
- Kho, W., Cheah, W. L., & Hazmi, H. (2018). Elevated blood pressure and its predictors among secondary school students in Sarawak: a cross-sectional study. *Central European journal of public health*, 26(1), 16-21.
- Khoury, M., Manlihot, C., & McCrindle, B. W. (2013). Role of the waist/height ratio in the cardiometabolic risk assessment of children classified by body mass index. *Journal of the American College of Cardiology*, 62(8), 742-751.
- Kim, H. S. (2019). Simple universal cutoff point of waist-height ratio for metabolic risk in Korean children and adolescents. In *Korean Journal of Pediatrics* (Vol. 62, Issue 7, pp. 257–258). Korean Pediatric Society. <https://doi.org/10.3345/kjp.2019.00045>
- Kim, J. Y., Cho, H., & Kim, J. H. (2022). Difference in the Prevalence of Elevated Blood Pressure and Hypertension by References in Korean Children and Adolescents. *Frontiers in Medicine*, 9. <https://doi.org/10.3389/fmed.2022.793771>
- Knowles, G., Pallan, M., Thomas, G. N., Ekelund, U., Cheng, K. K., Barrett, T., & Adab, P. (2013). Physical activity and blood pressure in primary school children: A longitudinal study. *Hypertension*, 61(1), 70–75. <https://doi.org/10.1161/HYPERTENSIONAHA.112.201277>
- Koster, A., Caserotti, P., Patel, K.V., Matthews, C.E., Berrigan, D., Van Domelen, D.R., Brychta, R.J., Chen, K.Y. and Harris, T.B., 2012. Association of sedentary time with mortality independent of moderate to vigorous physical activity. *PLoS one*, 7(6), p.e37696.
- Krakauer, N. Y., & Krakauer, J. C. (2012). A new body shape index predicts mortality hazard independently of body mass index. *PLoS one*, 7(7), e39504.
- Kuciene, R., & Dulskiene, V. (2014). Associations of short sleep duration with prehypertension and hypertension among Lithuanian children and adolescents: A cross-sectional study. *BMC Public Health*, 14(1). <https://doi.org/10.1186/1471-2458-14-255>
- Kuciene, R., & Dulskiene, V. (2019). Associations between body mass index, waist circumference, waist-to-height ratio, and high blood pressure among adolescents: a cross-sectional study. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-45956-9>

- Kurnianto, A., Kurniadi Sunjaya, D., Ruluwedrata Rinawan, F., & Hilmanto, D. (2020). Prevalence of Hypertension and Its Associated Factors among Indonesian Adolescents. *International Journal of Hypertension*, 2020. <https://doi.org/10.1155/2020/4262034>
- Lee, J. W., Lim, N. K., Baek, T. H., Park, S. H., & Park, H. Y. (2015). Anthropometric indices as predictors of hypertension among men and women aged 40-69 years in the Korean population: The Korean Genome and Epidemiology Study. *BMC Public Health*, 15(1). <https://doi.org/10.1186/s12889-015-1471-5>
- Lee, S. T., Wong, J. E., Shanita, S. N., Ismail, M. N., Deurenberg, P., & Poh, B. K. (2015). Daily physical activity and screen time, but not other sedentary activities, are associated with measures of obesity during childhood. *International Journal of Environmental Research and Public Health*, 12(1), 146–161. <https://doi.org/10.3390/ijerph120100146>
- Lee, Y. Y., Kamarudin, K. S., & Wan Muda, W. A. M. (2019). Associations between self-reported and objectively measured physical activity and overweight/obesity among adults in Kota Bharu and Penang, Malaysia. *BMC Public Health*, 19(1). <https://doi.org/10.1186/s12889-019-6971-2>
- Leone, A., Vizzuso, S., Brambilla, P., Mameli, C., Ravella, S., de Amicis, R., Battezzati, A., Zuccotti, G., Bertoli, S., & Verduci, E. (2020). Evaluation of different adiposity indices and association with metabolic syndrome risk in obese children: Is there a winner? *International Journal of Molecular Sciences*, 21(11). <https://doi.org/10.3390/ijms21114083>
- Liew JK, Cheong XP, Law L, Teo WH, & Eng SS. (2019). *Prevalence and Factors Associated with Hypertension among Adolescents in Malaysia* (Vol. 18, Issue 1).
- Lo, K., Wong, M., Khalechelvam, P., & Tam, W. (2016). Waist-to-height ratio, body mass index and waist circumference for screening paediatric cardio-metabolic risk factors: a meta-analysis. *Obesity Reviews*, 17(12), (1258–1275). Blackwell Publishing Ltd. <https://doi.org/10.1111/obr.12456>
- Ma, C., Wang, R., Liu, Y., Lu, Q., Lu, N., Tian, Y., Liu, X., & Yin, F. (2016). Performance of obesity indices for screening elevated blood pressure in pediatric population Systematic review and meta-analysis. In *Medicine (United States)* (Vol. 95, Issue 39). Lippincott Williams and Wilkins. <https://doi.org/10.1097/MD.0000000000004811>
- Macdonald-Wallis, C., Solomon-Moore, E., Sebire, S. J., Thompson, J. L., Lawlor, D. A., & Jago, R. (2017). A longitudinal study of the associations of children's body mass index and physical activity with blood pressure. *PLoS ONE*, 12(12). <https://doi.org/10.1371/journal.pone.0188618>
- Maessen, M. F., Eijsvogels, T. M., Verheggen, R. J., Hopman, M. T., Verbeek, A. L., & Vegt, F. D. (2014). Entering a new era of body indices: the feasibility of a body shape index and body roundness index to identify cardiovascular health status. *PloS one*, 9(9), e107212.

- Makarem, N., Shechter, A., Carnethon, M. R., Mullington, J. M., Hall, M. H., & Abdalla, M. (2019). Sleep Duration and Blood Pressure: Recent Advances and Future Directions. In *Current Hypertension Reports* (Vol. 21, Issue 5). Current Medicine Group LLC 1. <https://doi.org/10.1007/s11906-019-0938-7>
- Malara, M., Kęska, A., Tkaczyk, J., & Lutosławska, G. (2015). Body shape index versus body mass index as correlates of health risk in young healthy sedentary men. *Journal of translational medicine*, 13(1), 1-5.
- Malavazos, A. E., Capitanio, G., Milani, V., Ambrogi, F., Matelloni, I. A., Basilio, S., Dubini, C., Sironi, F. M., Stella, E., Castaldi, S., Secchi, F., Menicanti, L., Iacobellis, G., Corsi Romanelli, M. M., Carruba, M. O., & Morricone, L. F. (2021). Tri-Ponderal Mass Index vs body Mass Index in discriminating central obesity and hypertension in adolescents with overweight. *Nutrition, Metabolism and Cardiovascular Diseases*, 31(5), 1613–1621. <https://doi.org/10.1016/j.numecd.2021.02.013>
- Malik, A. H., Akram, Y., Shetty, S., Malik, S. S., & Yanchou Njike, V. (2014). Impact of sugar-sweetened beverages on blood pressure. In *American Journal of Cardiology* (Vol. 113, Issue 9, pp. 1574–1580). Elsevier Inc. <https://doi.org/10.1016/j.amjcard.2014.01.437>
- Mccarthy, H. D., & Ashwell, M. (2006). A study of central fatness using waist-to-height ratios in UK children and adolescents over two decades supports the simple message-‘keep your waist circumference to half your height’. *International Journal of Obesity*.
- Ministry of Health (2018). Clinical practice guidelines: Management of hypertension. 5th edn. Ministry of Health Malaysia.
- Mohammed Nawi, A., Mohammad, Z., Jetly, K., Abd Razak, M. A., Ramli, N. S., Wan Ibadullah, W. A. H., & Ahmad, N. (2021). The Prevalence and Risk Factors of Hypertension among the Urban Population in Southeast Asian Countries: A Systematic Review and Meta-Analysis. In *International Journal of Hypertension* (Vol. 2021). Hindawi Limited. <https://doi.org/10.1155/2021/6657003>
- Mohan, B., Verma, A., Singh, K., Singh, K., Sharma, S., Bansal, R., Tandon, R., Goyal, A., Singh, B., Chhabra, S. T., Aslam, N., Wander, G. S., Roy, A., & Prabhakaran, D. (2019). Prevalence of sustained hypertension and obesity among urban and rural adolescents: A school-based, cross-sectional study in North India. *BMJ Open*, 9(9). <https://doi.org/10.1136/bmjopen-2018-027134>
- Mohd. Nor, H. A. (2020). Relation between Smartphone Addiction, Sleep Quality and Psychological Health among adolescents in Malaysia. *International Journal of Psychosocial Rehabilitation*, 24(4), 4288–4298. <https://doi.org/10.37200/ijpr/v24i4/pr201535>
- Mu, M., Xu, L. F., Hu, D., Wu, J., & Bai, M. J. (2017). Dietary Patterns and Overweight/Obesity: A Review Article. *Iranian journal of public health*, 46(7), 869-876.

- Nasution, A. T. P., Ramayati, R., Sofyani, S., Ramayani, O. Rr., & Siregar, R. (2017). Quality of sleep and hypertension in adolescents. *Paediatrica Indonesiana*, 56(5), 272. <https://doi.org/10.14238/pi56.5.2016.272-6>
- National Health and Morbidity Survey (NHMS). (2019). *Non-Communicable Diseases, Risk Factors & Other Health Problems*.
- Navarro-Solera, M., Carrasco-Luna, J., Pin-Arboledas, G., González-Carrascosa, R., Soriano, J. M., & Codoñer-Franch, P. (2015). Short sleep duration is related to emerging cardiovascular risk factors in obese children. *Journal of pediatric gastroenterology and nutrition*, 61(5), 571-576.
- Nevill, A. M., Reuter, C. P., Brand, C., Gaya, A. R., Mota, J., Renner, J. D. P., & Duncan, M. J. (2021). BMI fails to reflect the developmental changes in body fatness between boys and girls during adolescence. *International Journal of Environmental Research and Public Health*, 18(15). <https://doi.org/10.3390/ijerph18157833>
- Nkwana, M. R., Monyeki, K. D., & Lebelo, S. L. (2021). Body roundness index, a body shape index, conicity index, and their association with nutritional status and cardiovascular risk factors in South African rural young adults. *International Journal of Environmental Research and Public Health*, 18(1), 1–13. <https://doi.org/10.3390/ijerph18010281>
- Nor, N. K., Ghazali, A. H., & Ismail, J. (2019). Prevalence of overweight and obesity among children and adolescents with autism spectrum disorder and associated risk factors. *Frontiers in Pediatrics*, 7(FEB). <https://doi.org/10.3389/fped.2019.00038>
- Noubiap, J. J., Essouma, M., Bigna, J. J., Jingi, A. M., Aminde, L. N., & Nansseu, J. R. (2017). Prevalence of elevated blood pressure in children and adolescents in Africa: a systematic review and meta-analysis. *The Lancet Public Health*, 2(8), e375–e386. [https://doi.org/10.1016/S2468-2667\(17\)30123-8](https://doi.org/10.1016/S2468-2667(17)30123-8)
- Nuttall, F. Q. (2015). Body mass index: Obesity, BMI, and health: A critical review. In *Nutrition Today* (Vol. 50, Issue 3, pp. 117–128). Lippincott Williams and Wilkins. <https://doi.org/10.1097/NT.0000000000000092>
- Nyamdorj, R., Qiao, Q., Söderberg, S., Pitkäniemi, J.M., Zimmet, P.Z., Shaw, J.E., Alberti, K.G.M.M., Pauvaday, V.K., Chitson, P., Kowlessur, S. and Tuomilehto, J. (2009). BMI compared with central obesity indicators as a predictor of diabetes incidence in Mauritius. *Obesity*, 17(2), 342-348.
- Ononamadu, C. J., Ezekwesili, C. N., Onyeukwu, O. F., Umeoguaju, U. F., Ezeigwe, O. C., & Ihegboro, G. O. (2017). Comparative analysis of anthropometric indices of obesity as correlates and potential predictors of risk for hypertension and prehypertension in a population in Nigeria. *Cardiovascular Journal of Africa*, 28(2), 92–99. <https://doi.org/10.5830/CVJA-2016-061>

- Paciência, I., Araújo, J., & Ramos, E. (2016). Sleep duration and blood pressure: a longitudinal analysis from early to late adolescence. *Journal of Sleep Research*, 25(6), 702–708. <https://doi.org/10.1111/jsr.12433>
- Park, H. K., & Shim, Y. S. (2020). Distribution of tri-ponderal mass index and its relation to body mass index in children and adolescents aged 10 to 20 years. *The Journal of Clinical Endocrinology & Metabolism*, 105(3), e826-e834.
- Payab, M., Kelishadi, R., Qorbani, M., Motlagh, M.E., Ranjbar, S.H., Ardalan, G., Zahedi, H., Chinian, M., Asayesh, H., Larijani, B. and Heshmat, R. (2015). Association of junk food consumption with high blood pressure and obesity in Iranian children and adolescents: the CASPIAN-IV Study☆. *Jornal de pediatria*, 91, 196-205.
- Pazin, D. C., da Luz Kaestner, T. L., Olandoski, M., Baena, C. P., de Azevedo Abreu, G., Kuschnir, M. C. C., Bloch, K. V., & Faria-Neto, J. R. (2020). Association between abdominal waist circumference and blood pressure in Brazilian adolescents with normal body mass index: Waist circumference and blood pressure in Adolescents. *Global Heart*, 15(1). <https://doi.org/10.5334/GH.779>
- Pazin, D. C., Rosaneli, C. F., Olandoski, M., de Oliveira, E. R. N., Baena, C. P., Figueredo, A. S., Baraniuk, A. O., Kaestner, T. L. da L., Guarita-Souza, L. C., & Faria-Neto, J. R. (2017). Waist circumference is associated with blood pressure in children with normal body mass index: A cross-sectional analysis of 3,417 school children. *Arquivos Brasileiros de Cardiologia*, 109(6), 509–515. <https://doi.org/10.5935/abc.20170162>
- Peach, H., Gaultney, J. F., & Reeve, C. L. (2015). Sleep characteristics, body mass index, and risk for hypertension in young adolescents. *Journal of Youth and Adolescence*, 44(2), 271–284. <https://doi.org/10.1007/s10964-014-0149-0>
- Perng, W., Tang, L., Song, P. X., Goran, M., Tellez Rojo, M. M., Cantoral, A., & Peterson, K. E. (2019). Urate and nonanoate mark the relationship between sugar-sweetened beverage intake and blood pressure in adolescent girls: a metabolomics analysis in the element cohort. *Metabolites*, 9(5), 100.
- Peterson, C. M., Su, H., Thomas, D. M., Heo, M., Golnabi, A. H., Pietrobelli, A., & Heymsfield, S. B. (2017). Tri-ponderal mass index vs body mass index in estimating body fat during adolescence. *JAMA pediatrics*, 171(7), 629-636.
- Poh, B. K., Jannah, A. N., Chong, L. K., Ruzita, A. T., Ismail, M. N., & McCarthy, D. (2011). Waist circumference percentile curves for Malaysian children and adolescents aged 6.016.9 years. *International Journal of Pediatric Obesity*, 6(3–4), 229–235. <https://doi.org/10.3109/17477166.2011.583658>
- Poitras, V.J., Gray, C.E., Borghese, M.M., Carson, V., Chaput, J.P., Janssen, I., Katzmarzyk, P.T., Pate, R.R., Connor Gorber, S., Kho, M.E. and Sampson, M. (2016). Systematic review of the relationships between objectively measured physical activity and health indicators in school-aged children and youth. *Applied Physiology, Nutrition, and Metabolism*, 41(6), S197-S239.

- Powell-Wiley, T. M., Poirier, P., Burke, L. E., Després, J. P., Gordon-Larsen, P., Lavie, C. J., Lear, S. A., Ndumele, C. E., Neeland, I. J., Sanders, P., & St-Onge, M. P. (2021). Obesity and Cardiovascular Disease: A Scientific Statement from the American Heart Association. In *Circulation* (pp. E984–E1010). Lippincott Williams and Wilkins. <https://doi.org/10.1161/CIR.0000000000000973>
- Qin, Z., Xu, F., Ye, Q., Zhou, H., Li, C., He, J., Wang, Z., Hong, X. and Hou, X., 2018. Sugar-sweetened beverages and school students' hypertension in urban areas of Nanjing, China. *Journal of human hypertension*, 32(6), pp.392-396.
- Rampal L, KC Ng, Nur Izzati I, Farah Izzati Z, Mohammad Nazrul I, Faisal I, & Sharifah Zainiyah SY. (2011). *Prevalence of Hypertension Among Malay Adolescents in Putrajaya Secondary Schools, Malaysia, 2010*.
- Rocha, N. P., Milagres, L. C., Longo, G. Z., Ribeiro, A. Q., & Novaes, J. F. D. (2017). Association between dietary pattern and cardiometabolic risk in children and adolescents: a systematic review☆. *Jornal de pediatria*, 93, 214-222.
- Rodea-Montero, E. R., Evia-Viscarra, M. L., & Apolinar-Jiménez, E. (2014). Waist-to-height ratio is a better anthropometric index than waist circumference and BMI in predicting metabolic syndrome among obese mexican adolescents. *International Journal of Endocrinology*, 2014. <https://doi.org/10.1155/2014/195407>
- Roy, S., & Ghosh, J. (2019). Comparison of adiposity measures in the identification of adolescents with hypertension. *Sri Lanka J Child Health*, 48(1), 39-46.
- Sardinha, L. B., Santos, D. A., Silva, A. M., Grøntved, A., Andersen, L. B., & Ekelund, U. (2016). A comparison between BMI, waist circumference, and waist-to-height ratio for identifying cardio-metabolic risk in children and adolescents. *PLoS ONE*, 11(2). <https://doi.org/10.1371/journal.pone.0149351>
- Shariq, O. A., & Mckenzie, T. J. (2020). Obesity-related hypertension: A review of pathophysiology, management, and the role of metabolic surgery. *Gland Surgery*, 9(1), 80–93. <https://doi.org/10.21037/gs.2019.12.03>
- Shim, Y. S. (2019). The Relationship Between Tri-ponderal Mass Index and Metabolic Syndrome and Its Components in Youth Aged 10–20 Years. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-50987-3>
- Solomon-Moore, E., Salway, R., Emm-Collison, L., Thompson, J. L., Sebire, S. J., Lawlor, D. A., & Jago, R. (2020). Associations of body mass index, physical activity and sedentary time with blood pressure in primary school children from south-west England: A prospective study. *PLoS ONE*, 15(4). <https://doi.org/10.1371/journal.pone.0232333>
- Song, P., Zhang, Y., Yu, J., Zha, M., Zhu, Y., Rahimi, K., & Rudan, I. (2019a). Global Prevalence of Hypertension in Children: A Systematic Review and Meta-analysis. In *JAMA Pediatrics*, 173 (12), 1154–1163). American Medical Association. <https://doi.org/10.1001/jamapediatrics.2019.3310>

- Song, X., Jousilahti, P., Stehouwer, C.D.A., Söderberg, S., Onat, A., Laatikainen, T., Yudkin, J.S., Dankner, R., Morris, R., Tuomilehto, J. and Qiao, Q. (2015). Cardiovascular and all-cause mortality in relation to various anthropometric measures of obesity in Europeans. *Nutrition, Metabolism and Cardiovascular Diseases*, 25(3), 295-304.
- Soua, S., Ghammam, R., Maatoug, J., Zammit, N., ben Fredj, S., Martinez, F., & Ghannem, H. (2022). The prevalence of high blood pressure and its determinants among Tunisian adolescents. *Journal of Human Hypertension*. <https://doi.org/10.1038/s41371-022-00677-x>
- Souza, B. D. S., Cunha, D. B., Pereira, R. A., & Sichieri, R. (2016). Soft drink consumption, mainly diet ones, is associated with increased blood pressure in adolescents. *Journal of hypertension*, 34(2), 221-225.
- Sreeramareddy, C., Chew, W., Poulsaeman, V., Boo, N., Choo, K., & Yap, S. (2013). Blood pressure and its associated factors among primary school children in suburban Selangor, Malaysia: A cross-sectional survey. *Journal of Family and Community Medicine*, 20(2), 90–97. <https://doi.org/10.4103/2230-8229.114769>
- Stabouli, S., Papakatsika, S., & Kotsis, V. (2011). The role of obesity, salt and exercise on blood pressure in children and adolescents. *Expert review of cardiovascular therapy*, 9(6), 753-761.
- Stanaway, J. D., Afshin, A., Gakidou, E., Lim, S. S., Abate, D., Abate, K. H., Abbafati, C., Abbasi, N., Abbastabar, H., Abd-Allah, F., Abdela, J., Abdelalim, A., Abdollahpour, I., Abdulkader, R. S., Abebe, M., Abebe, Z., Abera, S. F., Abil, O. Z., Abraha, H. N., ... Murray, C. J. L. (2018). Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*, 392(10159), 1923–1994. [https://doi.org/10.1016/S0140-6736\(18\)32225-6](https://doi.org/10.1016/S0140-6736(18)32225-6)
- Su, T. T., Sim, P. Y., Nahar, A. M., Majid, H. A., Murray, L. J., Cantwell, M. M., Al-Sadat, N., & Jalaludin, M. Y. (2014). Association between self-reported physical activity and indicators of body composition in Malaysian adolescents. *Preventive Medicine*, 67, 100–105. <https://doi.org/10.1016/j.ypmed.2014.07.001>
- Sun, J., Yang, R., Zhao, M., Bovet, P., & Xi, B. (2021). Tri-Ponderal Mass Index as a Screening Tool for Identifying Body Fat and Cardiovascular Risk Factors in Children and Adolescents: A Systematic Review. In *Frontiers in Endocrinology* (Vol. 12). Frontiers Media S.A. <https://doi.org/10.3389/fendo.2021.694681>
- Tang, X., Liu, Y., Hu, J., Zhai, L., Jia, L., Ding, N., Ma, Y., & Wen, D. (2022). Association of waist circumference with blood pressure and familial dietary habits in preschool children: a cross-sectional study in northeastern China. *Italian Journal of Pediatrics*, 48(1). <https://doi.org/10.1186/s13052-022-01236-3>

- Tee, J. Y. H., Gan, W. Y., & Lim, P. Y. (2020). Comparisons of body mass index, waist circumference, waist-to-height ratio and a body shape index (ABSI) in predicting high blood pressure among Malaysian adolescents: A cross-sectional study. *BMJ Open*, 10(1). <https://doi.org/10.1136/bmjopen-2019-032874>
- Tee, J. Y. H., Gan, W. Y., Tan, K. A., & Chin, Y. S. (2018). Obesity and unhealthy lifestyle associated with poor executive function among Malaysian adolescents. *PLoS ONE*, 13(4), 1–17. <https://doi.org/10.1371/journal.pone.0195934>
- Thomas, D.M., Bredlau, C., Bosy-Westphal, A., Mueller, M., Shen, W., Gallagher, D., Maeda, Y., McDougall, A., Peterson, C.M., Ravussin, E. & Heymsfield, S.B. (2013). Relationships between body roundness with body fat and visceral adipose tissue emerging from a new geometrical model. *Obesity*, 21(11), 2264-2271.
- Tremblay, M.S., LeBlanc, A.G., Kho, M.E., Saunders, T.J., Larouche, R., Colley, R.C., Goldfield, G. and Gorber, S.C. (2011). Systematic review of sedentary behaviour and health indicators in school-aged children and youth. *International journal of behavioral nutrition and physical activity*, 8(1), 1-22.
- Tsioufis, C., Kyvelou, S., Tsiachris, D., Tolis, P., Hararis, G., Koufakis, N., Psaltopoulou, T., Panagiotakos, D., Kokkinos, P., & Stefanadis, C. (2011). Relation between physical activity and blood pressure levels in young Greek adolescents: The Leontio Lyceum Study. *European Journal of Public Health*, 21(1), 63–68. <https://doi.org/10.1093/eurpub/ckq006>
- UNICEF. (2017). *Children, food and nutrition : growing well in a changing world*.
- Uwaezuoke, S., Okoli, C., Ubesie, A., & Ikefuna, A. (2016). Primary hypertension among a population of Nigerian secondary school adolescents: Prevalence and correlation with anthropometric indices: A cross-sectional study. *Nigerian Journal of Clinical Practice*, 19(5), 649–654. <https://doi.org/10.4103/1119-3077.188706>
- Valdez, R. (1991). A simple model-based index of abdominal adiposity. *Journal of clinical epidemiology*, 44(9), 955-956.
- Wang, H., Liu, A., Zhao, T., Gong, X., Pang, T., Zhou, Y., Xiao, Y., Yan, Y., Fan, C., Teng, W., Lai, Y., & Shan, Z. (2017). Comparison of anthropometric indices for predicting the risk of metabolic syndrome and its components in Chinese adults: A prospective, longitudinal study. *BMJ Open*, 7(9). <https://doi.org/10.1136/bmjopen-2017-016062>
- Wang, L., Song, L., Liu, B., Zhang, L., Wu, M., Cao, Z., & Wang, Y. (2019). Trends and Status of the Prevalence of Elevated Blood Pressure in Children and Adolescents in China: a Systematic Review and Meta-analysis. In *Current Hypertension Reports* (Vol. 21, Issue 11). Current Medicine Group LLC 1. <https://doi.org/10.1007/s11906-019-0992-1>

- Wang, X., Dong, B., Ma, J., Song, Y., Zou, Z., & Arnold, L. (2020). Role of tri-ponderal mass index in cardio-metabolic risk assessment in children and adolescents: compared with body mass index. *International Journal of Obesity*, 44(4), 886–894. <https://doi.org/10.1038/s41366-019-0416-y>
- Wang, Y., Wang, L., Xue, H., & Qu, W. (2016). A review of the growth of the fast food industry in china and its potential impact on obesity. In *International Journal of Environmental Research and Public Health* (Vol. 13, Issue 11). MDPI. <https://doi.org/10.3390/ijerph13111112>
- Wellman, R. J., Sylvestre, M. P., Abi Nader, P., Chiolero, A., Mesidor, M., Dugas, E. N., Tougri, G., & O'Loughlin, J. (2020). Intensity and frequency of physical activity and high blood pressure in adolescents: A longitudinal study. *Journal of Clinical Hypertension*, 22(2), 283–290. <https://doi.org/10.1111/jch.13806>
- World Health Organization. (2021). Guideline for the pharmacological treatment of hypertension in adults: web annex A: summary of evidence.
- Wu, F., Buscot, M.J., Niinikoski, H., Rovio, S.P., Juonala, M., Sabin, M.A., Jula, A., Rönnemaa, T., Viikari, J.S., Raitakari, O.T. and Magnussen, C.G. (2020). Age-specific estimates and comparisons of youth tri-ponderal mass index and body mass index in predicting adult obesity-related outcomes. *The Journal of pediatrics*, 218, 198-203.
- Xu, Y., Yan, W., & Cheung, Y. B. (2015). Body shape indices and cardiometabolic risk in adolescents. *Annals of Human Biology*, 42(1), 70–75. <https://doi.org/10.3109/03014460.2014.903998>
- Yang, Y., Min, J., Chang, L., Chai, J., Song, Z., Zha, S., Zhang, M., Liu, H., & Yang, F. (2021). Prevalence trends of hypertension among 9–17 aged children and adolescents in Yunnan, 2017–2019: a serial cross-sectional surveillance survey. *BMC Public Health*, 21(1). <https://doi.org/10.1186/s12889-021-10258-1>
- Yuan, P., Qian, Z. (Min), Vaughn, M., Huang, J., Ward, P., Zhu, Y., Qin, X. di, Zhou, Y., Li, M., Xu, S., Zhang, Y. Z., Bao, W. W., Hao, Y. T., Zeng, X. W., & Dong, G. H. (2017). Comparison of body mass index with abdominal obesity for identifying elevated blood pressure in children and adolescents: The SNEC study. *Obesity Research and Clinical Practice*, 11(4), 406–413. <https://doi.org/10.1016/j.orcp.2016.08.006>
- Zaki, N. A. M., Sahril, N., Omar, M. A., Ahmad, M. H., Baharudin, A., & Nor, N. S. M. (2016). Reliability and validity of the physical activity questionnaire for older children (PAQ-C) in Malay language. *International Journal of Public Health Research*, 6(1), 670-676.
- Zhang, N., Chang, Y., Guo, X., Chen, Y., Ye, N., & Sun, Y. (2016). A body shape index and body roundness index: two new body indices for detecting association between obesity and hyperuricemia in rural area of China. *European journal of internal medicine*, 29, 32-36.

Zhao, Y., Wang, L., Xue, H., Wang, H., & Wang, Y. (2017). Fast food consumption and its associations with obesity and hypertension among children: results from the baseline data of the Childhood Obesity Study in China Mega-cities. *BMC Public Health*, 17(1), 933. <https://doi.org/10.1186/s12889-017-4952-x>

