



UNIVERSITI PUTRA MALAYSIA

**GROWTH PERFORMANCE, BODY CONFORMATION, CARCASS
CHARACTERISTICS AND PALATABILITY OF KEDAH-KELANTAN
CATLE AND ITS CROSSES**

DAHLAN BIN ISMAIL

FPV 1985 3

GROWTH PERFORMANCE, BODY CONFORMATION,
CARCASS CHARACTERISTICS AND PALATABILITY
OF KEDAH-KELANTAN CATTLE AND ITS CROSSES

by

Dahlan bin Ismail

A thesis submitted in partial fulfilment of the degree of Master
of Science in the Faculty of Veterinary Medicine and Animal
Science, Universiti Pertanian Malaysia.

July 1985



This thesis is dedicated to my father Ismail bin Md. Yunos and my mother Allahyarhammah Maimunah bte Yusoff who introduced me to this world



ACKNOWLEDGEMENTS

I wish to express my gratitude to Dr. Mohamed Mahyuddin bin Mohd. Dahan who provided continuous guidance, support and encouragement, not only in the preparation of this thesis, but also throughout the duration of my graduate studies.

Special thanks are extended to En. Ahmad Shokri Hj. Othman for his consultation in the statistical analysis of the data and Dr. Pravee Vijchulata for his valuable suggestions and constructive criticism in the final preparation of this manuscript.

Sincere appreciation is given to En. Shahril Hassan, En. Ahmad Tarmizi, En. Adnan Yassin and En. Chua Eng Chong for their assistance in the carcass evaluation and meat science works. Further acknowledgements are also due to Dr. Mohd. Arief Omar, En. Mohd. Nor Hamid, En. Yahya Mail, Puan Norpisah Abdul and En. Mohsodi Surip for their involvement directly or indirectly in conducting the research and collection of data in the beef cattle crossbreeding programme in MARDI Station near Kluang.

I would like to thank Dr. Syed Ali Abu Bakar, Director of Livestock Research in MARDI who has given me the permission and encouragement in conducting this research.



Further acknowledgements are due to Malaysian Agricultural Research and Development Institute (MARDI) for funding the research and to Universiti Pertanian Malaysia for providing the slaughtering and meat laboratory facilities.

This manuscript and the work it represents could not have been achieved without the patience and understanding of my wife, Hamidah and my two wonderful daughters, Nur Dalilah and Nur Khalidah.



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT	iii
TABLE OF CONTENTS	v
LIST OF TABLES	vii
LIST OF FIGURES	x
ABSTRACT	xi
CHAPTER I INTRODUCTION	1
CHAPTER II REVIEW OF LITERATURE	4
Genetic improvement of beef cattle	4
Breed evaluation of beef cattle	6
Growth performance	8
Body conformation	14
Carcass characteristics	18
Palatability	22
Carcass characteristics of local cattle	25
CHAPTER III EXPERIMENTAL PROCEDURE	27
CHAPTER IV RESULTS AND DISCUSSIONS	48
Growth performance	48
Liveweight-age regression coefficients	58
Body conformation and measurements	62
Temperament score	68
Carcass characteristics	71
Carcass composition	77
Wholesale cuts or special cuts	83



	Page
Correlations among carcass traits	83
Palatability	87
Predicting total lean meat	94
CHAPTER V SUMMARY AND CONCLUSIONS	100
LITERATURE CITED	106
APPENDICES	120
BIBLIOGRAPHICAL SKETCH	137
PUBLICATIONS	138



LIST OF TABLES

Table		Page
I	Kedah-Kelantan cows in the breeding herd	28
II	Mating system	29
III	Number of straightbred and crossbred KK cattle progenies	32
IV	Slaughter weight grouping of straightbred and crossbreds KK cattle	34
V	Means and standard deviation values for liveweight of male and female KK cattle and its crosses (pooled weight 1979-1982)	49
VI	Percent advantage in liveweight of F ₁ KK crossbreds compared with straightbred KK cattle	51
VII	Mean and standard deviation of adjusted 200-day weaning weight and means of actual weaning age of KK cattle and its crosses (pooled 1979-1981 calves)	54
VIII	Average daily gain of KK cattle and its crossbreds	56
IX	Table of T-values from comparison of liveweight regression lines between adjusted means of male and female KK cattle and its crossbreds (pooled weight)	59
X	Comparison of regression lines of liveweight between KK and its crossbreds	61
XI	Mean and standard error values for the starved weight, conformation score and body measurement of KK cattle and its crossbreds	63
XII	Correlation coefficients between starved weight, conformation score and body measurement of KK cattle and its crossbreds	67
XIII	Temperament scores for KK cattle and its crossbreds	69
XIV	Liveweight, carcass weight and dressing percentage at different weight groups of KK and KK crossbreds bull	72



Table	Page
XV	Carcass characteristics of KK and KK crossbreds cattle 74
XVI	Mean and standard deviation values of meat components in KK cattle and its crossbreds carcass 78
XVII	Mean values of fat components in KK cattle and its crossbreds carcass 80
XVIII	Correlation coefficients between meat, fat and body measurements 82
XIX	Correlation coefficients between starved weight, age, conformation score and carcass traits 84
XX	Relationship between starved weight, age and some carcass traits of KK cattle and its crossbreds 86
XXI	Comparison of mean values for palatability traits, shear force value, ultimate muscle pH, cooking loss, age group and temperament score . 88
XXII	Correlation coefficients between palatability score, shear force value, starved weight and age 91
XXIII	Relationship between ultimate pH of meat and some palatability traits, shear force value and cooking loss of <u>Longissimus dorsi</u> muscle of KK and its crossbreds 92
XXIV	The best prediction equations for dependent variable 'total lean meat' from some of the body measurements (all breedtypes) 95
XXV	The best one variable model for prediction of total lean meat for all breedtypes 96
XXVI	The best prediction equations for dependent variable 'total lean meat' from some of the carcass traits (all breedtypes) 98



LIST OF TABLES IN APPENDIX

Appendix		Page
I	Live body measurements and conformation	120
II	Carcass traits measurements	122
III	Beef palatability, shear force value and cooking data	126
IV	External and internal organs (offals)	127
V	Correlation coefficients between total lean meat and offals	130
VI	Correlation coefficients between palatability scores, starved weight, age, pH and temperament score of KK cattle	131
VII	Mean values of special cuts for KK and KK crossbreds	132
VIII	Chemical composition of <u>Longissimus dorsi</u> muscle	136
IX	Means, standard deviation, minimum and maximum values of palatability scores, W-B values, pH, and temperament scores (all breed types)	137



LIST OF FIGURES

Figure		Page
1	Live body measurements of cattle	36
2	The procedure of fabricating the beef cattle ...	41
	A. Forequarter	
	B. Hindquarter	



An abstract of the thesis presented to the Senate of Universiti Pertanian Malaysia in partial fulfilment of the requirements for the Degree of Master of Science.

GROWTH PERFORMANCE, BODY CONFORMATION,
CARCASS CHARACTERISTICS AND PALATABILITY
OF KEDAH-KELANTAN CATTLE AND ITS CROSSES

by

Dahlan bin Ismail

July 1985

Supervisor: Mohamed Mahyuddin bin Mohd. Dahan, Ph.D.

Faculty: Veterinary Medicine and Animal Sciences

The growth performance of 423 heads of male and female Kedah-Kelantan (KK) cattle and its F₁ crosses were recorded from the same grazing management. Comparisons were made between progenies of KK cows sired by straightbred KK, American Brahman (Br), Polled Hereford (Hr), Friesian (Fr) and Jersey (Jv).

The main objectives of this experiment was to evaluate the simplest crossbreeding system of KK cattle by straight cross between KK female and exotic male for the advantage of heterosis in the growth performance, conformation, carcass characteristics and meat palatability in the F₁ generation.

The present result showed that all the F₁ crossbred heifers and bulls were superior in the growth performance than the



straightbred KK at all ages. All crossbred calves had significantly higher ($P < 0.01$) birth weight than KK. Brahman x KK (BK) calves were the heaviest at birth, followed by Friesian x KK (FK), Hereford x KK (HK) and Jersey x KK (JK), respectively. The overall percent advantage of crossbreds at birth weight over KK was 27.13%. FK was the heaviest at the adjusted 200-day weight, followed by HK, BK, JK and KK respectively. All crossbreds calves were showing better pre-weaning growing ability than the KK. HK showed superiority in post weaning growth performance than the other crossbreds. Ranking of sire breeds by using the slope of regression equation of growth-age lines as the estimate of growth rate, generally showed consistency for the breedtype among years. Under similar management unit and one type of maternal environment, the result showed that breed sire, especially temperate sires, had contributed the main effect on all the growth traits of these calves.

HK showed higher conformation score due to the tendency of the beef breed crosses to fatten at lighter weight than the dairy breed crosses. HK and FK had the longest body, FK was the tallest at wither, BK was the tallest at the hip, KK was the shortest cattle. HK was the shortest among the crossbreds at the hip.

There is evidence for breed difference in the temperament. This study concluded that Bos indicus cattle and crosses (Brahman x KK) were having poor temperament than half-bred cattle (FK, HK and JK).



The result of carcass evaluation concluded that breed of sire had no significant effect on factor associated with dressing percentage but had significant effect ($P < 0.05$) on the carcass weight per day of age (CWDA). All the crossbreds showed higher CWDA than KK in all slaughter weight group. HK showed the highest CWDA and 21 to 44% superior than KK as the animal increases in slaughter weight. Animals sired by beef type (Hr and Br) produced higher percentage of hindquarter than the crosses from dairy type (Fr and Jy). HK showed the biggest ($P < 0.05$) loin eye area in all slaughter weight group.

KK bulls showed lower bone composition and higher lean composition than other breedtype under similar weight group. Fat composition in the carcass of Zebu cattle (KK and BK) was higher than Zebu x temperate cattle.

Comparison of palatability scores between breed type revealed no significant difference but the scores significantly decreased ($P < 0.05$) as the cattle increased in age from 28-month to 40-month. Among breedtypes, FK and BK showed higher shear force values than JK, HK and KK. The present result concluded that ultimate muscle pH of all breedtypes have some effect on the flavour, overall palatability, tenderness and cooking loss.

The best single body measurement for prediction of total lean meat of half carcass was the measurement of heart girth and the best 4 combination of body measurements found were heart



girth, width of loin, flank girth and width of head. Carcass measurements were also used for predictions of total lean meat. The result showed that forequarter weight, loin eye area, meat to bone ratio, carcass length and dressing percentage are the best 5 variables that should be used in the combination.



Abstrak

Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia sebagai memenuhi sebahagian dari keperluan untuk Ijazah Master Sains.

GROWTH PERFORMANCE, BODY CONFORMATION,
CARCASS CHARACTERISTICS AND PALATABILITY OF
KEDAH-KELANTAN CATTLE AND ITS CROSSES

by

Dahlan bin Ismail

Julai 1985

Penyelia: Mohamed Mahyuddin bin Mohd. Dahan, Ph.D.

Fakulti: Kedokteran Veterinar dan Sains Peternakan

Prestasi tumbesaran 423 ekor lembu Kedah-Kelantan (KK) dan kacukan-kacukannya generasi F_1 , jantan dan betina telah direkodkan di bawah sistem ragutan yang sama. Pertandingan dilakukan di antara anak-anak lembu diibukan oleh baka KK dan baka jantannya dari baka-baka tulin KK, American Brahman (Br), Polled Hereford (Hr), Friesian (Fr) dan Jersey (Jy).

Tujuan utama kajian ini ialah untuk menilai sistem kacukan lembu KK betina dengan baka jantan eksotik untuk kebaikan prestasi tumbesaran, conformasi (bentuk), sifat-sifat karkas dan palatabiliti (penerimaan) daging pada anak-anak kacukan generasi F_1 .



Keputusan kajian ini menunjukkan kesemua anak-anak lembu kacukan F_1 (jantan dan betina) lebih tinggi prestasi tumbesarnya dari lembu KK tulin. Kesemua anak-anak kacukan menunjukkan berat lahir yang lebih tinggi dari KK ($P < 0.01$). Brahman x KK (BK) ialah yang terberat pada berat lahir diikuti oleh Friesian x KK (FK), Hereford x KK (HK) dan Jersey x KK (JK) berturutan. Keseluruhannya, baka-baka kacukan menunjukkan 27.13% lebih berat dari baka KK tulin pada berat lahir. HK adalah terberat pada berat anggaran 200-hari diikuti pula oleh HK, BK, JK dan KK berturutan. Kesemua anak-anak lembu kacukan menunjukkan kadar tumbesaran sebelum bercerai susu yang lebih baik dari anak lembu KK tulin. HK menunjukkan prestasi tumbesaran selepas bercerai susu yang tertinggi dari baka-baka kacukan lain. Tiap-tiap jenis baka bapa menunjukkan aturan kecerunan persamaan regrasi tumbesar-umur yang seakan-akan serupa bagi tiap-tiap jenis baka yang sama pada tahun-tahun kajian. Di bawah sistem pengurusan yang sama dan satu jenis baka ibu, keputusan menunjukkan baka bapa terutama bapa dari jenis Bos taurus, telah memainkan peranan yang penting terhadap kesan sifat tumbesaran anak-anak lembu ini.

HK menunjukkan 'score' conformasi yang tertinggi kerana keupayaan baka pedaging kacukan menggemuk pada umur yang lebih rendah dari baka kacukan. HK dan FK mempunyai badan yang terpanjang, FK tertinggi di bahu, BK tertinggi di pinggang, KK adalah lembu yang terendah, HK adalah baka kacukan yang terendah di pinggang dan di bahu.



Dari kajian ini juga terdapat bukti menunjukkan perbezaan antara baka di dalam sifat kelakuan (temperament). Kajian ini memutuskan bahawa lembu baka Bos indicus dan kacukannya (BK) menghadapi sifat kelakuan yang kurang terkawal dari lembu baka kacukan separa (FK, HK dan JK).

Keputusan uraikaji karkas menunjukkan baka bapa tidak menunjukkan kesan terhadap faktor-faktor bersangkutan dengan peratus pemotongan, tetapi berkesan ($P < 0.05$) pada berat karkas setiap hari umur (CWDA). Kesemua kacukan menunjukkan CWDA yang lebih tinggi dari KK dalam kesemua kumpulan penyembelihan. HK menunjukkan CWDA yang tertinggi dan lebih tinggi dari KK sebanyak 21 hingga 44% mengikut bertambahnya berat penyembelihan. Lembu yang menggunakan bapa jenis pedaging (Hr dan Br) menghasilkan peratus suku belakang karkas yang lebih tinggi dari bapa jenis tenusu (Fr dan Jy). HK menunjukkan luas daging bahu pinang (loin eye area) yang terbesar dalam kesemua kumpulan pemotongan ($P < 0.05$). Baka KK jantan menunjukkan kandungan tulang yang rendah dan kandungan daging yang tinggi dari jenis baka lain dalam kumpulan berat yang sama. Kandungan lemak lembu jenis Bos indicus (KK dan BK) lebih tinggi dari lembu Bos indicus x Bos taurus.

Perbandingan 'score' penerimaan daging (palatability score) antara jenis-jenis baka menunjukkan tidak ada kesan yang nyata tetapi score menurun ($P < 0.05$) apabila umur bertambah dari 28-bulan ke 40-bulan. Di antara jenis baka, FK dan BK menunjukkan daya keliatan (shear force) yang tinggi dari JK, HF dan KK.

Keputusan ini menunjukkan pfi otot tercapai (ultimate pfi) dari kesemua jenis baka ada mempunyai kesan terhadap rasa, keseluruhan penerimaan daging, kelembutan dan kehilangan berat semasa dimasak.

Ukuran badan yang terbaik untuk persamaan ramalan jumlah daging dari separuh karkas adalah ukuran lilitan dada, dan gabungan 4 ukuran badan yang terbaik ialah lilitan dada, lebar 'loin' lilitan pinggang dan lebar kepala. Ukuran-ukuran karkas juga digunakan untuk persamaan ramalan jumlah daging. Keputusan menunjukkan berat suku belakang, luas daging batang pinang, nisbah daging ke tulang, panjang karkas dan peratus perpotongan adalah 5 variabel terbaik patut digunakan secara gabungan.



CHAPTER I

INTRODUCTION

Beef production in tropical regions of the world is generally low compared to that in temperate zones. Although the developing countries (mostly in the tropical regions) have 70% of the world's cattle and buffalo population but they produce only 34% of the world's beef; thus they have unexploited resources (Jasiorowski, 1976). The bovine meat production from developing countries was only 15.5 million tons in 1982 compared to 31.2 million tons from developed countries (Asian Livestock, 1983). In general, the international beef export and import trade is dominated by the highly developed countries with only Argentina among the developing countries playing an important role on the export side.

In Malaysia, beef production is very low in terms of quantity and quality. The indigenous Kedah-Kelantan cattle are less productive in terms of growth, meat and milk yield. Their population is relatively small and there is currently a shortage of breeding females for multiplication. More than 90% of the cattle producer or rearer are the smallholders. Beef production amongst smallholders is mainly a supplementary activity. The farmer usually keeps one to a few heads of cattle or buffaloes which are integrated with the whole pattern of his cropping system. In 1980, the estimated local production of bovine meat



and consumption is around 13,000 and 20,000 tons, respectively (Anon, 1981). Thus, there is still insufficient production of meat to meet the national requirement.

There is an attempt to increase cattle production in this country but the Government have encountered many constraints. The first constraint is the availability of adequate number of improved genetic stock of cattle. Generally, animals which have evolved in a tropical environment are well adapted to that environment in terms of survival of the species but may not be well adapted in terms of providing optimum quantities of beef. For example, the indigenous Kedah-Kelantan (KK) cattle, although they have survived well in this country in terms of high fertility and total adaptation to the environment, these cattle are very slow in growth. Conversely, temperate breeds which have been evolved principally for beef production in temperate areas and have undergone various selection programmes, may not be well adapted for survival or production in tropical areas. The animals either do not survive or do not breed, or their performance is much poorer here than in their country of origin (Mason and Buvanendran, 1982). Breed development for meat and milk production is not only time consuming but also location specific. There is, therefore, a need for the genetic variability which is available in the indigenous breeds of cattle and in new production to be carefully assessed so that the kind of cattle propagated or developed are suitable for our conditions and environment.

While crossbreeding may have advantages for commercial production, straightbred stocks will be needed for commercial production on operations where crossbreeding is not practical and as source of breeding stock for crossbreeding operations. Little is known about the performance of some of the crossbred cattle in this country. Baharin (1978) reported that most of the trials conducted at UPM, MARDI, Institut Haiwan and commercial farms do indicate encouraging performance of the crossbreeds. However, most of the estimates are derived from a small sample of data. There is no information concerning the expected performance, carcass characteristics and the acceptability of the carcass of the crossbred in the local market. The availability of the information on the conformation and palatability of local breed and crossbreeds cattle under improved pasture condition is also scarce.

Realising the benefits of crossbreeding in beef production in terms of hybrid vigour in F_1 progeny, straight crossing between KK cows and purebred exotic breed bulls will be carried out in this study.

The objectives of this study are: (1) to study the growth performance, body conformation, carcass characteristics and meat palatability of the KK cattle and its crosses; (2) to provide information on temperamental behaviour of these cattle and (3) to develop prediction equations for estimating lean quantity in these cattle based on live body measurements and carcass measurements.

CHAPTER II

REVIEW OF LITERATURE

a) Genetic improvement of beef cattle

There exist two methods for genetic improvement: (1) selection which changes the additive genetic potential and (2) crossbreeding which is based on both additive and non-additive potential of the population. There is no doubt that both methods can and must be used in developing countries. But selection programmes is normally time consuming and requires sound production testing scheme. Genetic improvement through selection programmes for cattle in the developed countries have been advanced to the present stage, chiefly because the infrastructural elements necessary for planned breeding programmes such as intensive performance recording and A.I. are available. The recent advances have also been made in population genetics and statistics which have made it possible to develop genetic improvement schemes applicable on a national scale. In contrast, infrastructural elements for national breeding programmes are lacking or are inadequate in most parts of the tropics.

The quickest way to improve the productivity of commercial cattle in the tropics is to exploit hybrid vigour through systematic crossbreeding (Koger, 1963). There are several reasons for using crossbreeding in the tropics, apart from