



UNIVERSITI PUTRA MALAYSIA

**UTILIZATION OF SETARIA SPHACELATA VARIETY SPLENDIDA
BY KEDAH-KELANTAN CATTLE**

ROMZIAH SIDIK BUDIONO

FPV 1985 6



UTILIZATION OF SETARIA SPHACELATA VARIETY SPLENDIDA
BY KEDAH-KELANTAN CATTLE

by

ROMZIAH SIDIK BUDIONO

A thesis submitted in partial fulfilment of the
requirements for the degree of Doctor of Philosophy
in the Faculty of Veterinary Medicine and Animal Science
Universiti Pertanian Malaysia

May 1986



It is hereby certified that we have read this thesis entitled A Study on the Performance of Kedah-Kelantan Cattle Grazed on Setaria Sphacelata Variety Splendida by Romziah Sidik Budiono, and in our opinion it is satisfactory in terms of scope, quality and presentation as partial fulfilment of the requirements for the degree of Doctor of Philosophy.

(Chairman Board of Examiners)

(External Examiner)

(Internal Examiner)

(Supervisor)



Dedicated to the memory of my late father and father-in-law

H.M. Sidik and Margono Wiryodirdjo



ACKNOWLEDGEMENT

Praise be to the Almighty Allah. Had it not been His will, the completion of this study would not have been possible.

The author wishes to express her sincere gratitude and appreciation to all those who collaborated in preparation of this dissertation. This study would have come into form without the concerted efforts and/or valuable contributions of the following people and institutions.

- The advisors of her supervisory committee, Prof. Dr. Syed Jalaluddin bin Syed Salim, Dr. Pravee Vijchulata, Dr I.R. Lane and Prof. M.R. Jainudeen, for patiently and intelligently guiding her throughout the completion of her study;
- Prof. Dr. Abdul Latiff Ibrahim, Dean, Faculty of Veterinary Medicine and Animal Science, Dr. Tengku Azmi Tengku Ibrahim, Head, Department of Animal Science, and Assoc. Prof. Dr. Alang Perang Abdul Rahman Zainuddin, Dean of Post Graduate Studies, University Pertanian Malaysia,
- The Malaysian Public Services Department for the financial support.
- Dr. Baljid Singh and Mr. Tukiran of UPM Dairy and Beef Unit for offering the land and assisting her in organizing and helping with the sample collection in order to make this work possible.
- Dr. Idris Abdol and Mr. Ismail Sulong of Universiti Pertanian Malaysia for allowing the fencing of the ranch.
- All workers in fields II, XV and XVI of UPM farm for helping her in the collection of samples.
- Mrs. Nurhani Abdullah, Mr. Azahar Mohd. Noor, Mr. Ibrahim Mohsin, Mr. Rahman Hassan, Miss Roosmina Umar, Mrs. Rohana Atan, Mr. Syahril and Mr. Halim for their assistance in laboratory work.
- Dr. Zulkifli and Mr. Ibrahim of Soil Science Department of UPM for allowing her to use the laboratory facilities to analyse mineral elements.



- Mr. Yap Cheng Kee for helping her analyse data.
- Mr. and Mrs. Andrew Alex Tuen, Dr. Kassim Hamid and Mr. Mustapha Mamad, for their support in preparing the manuscript.
- Miss Armi Shamsuar and Miss Fadzlun Mohd. Yusof for the typing of this thesis.
- To her mothers, Mrs. Sidik and Mrs. Margono Wiryodirdjo for their encouragement.
- Her husband, Mr. Budiono Wiryodirdjo, for his inspiration, encouragement, patience and understanding during the execution of this task.

The author is greatly indebted to all mentioned that she wished they be absolved completely from blame for any omissions and commissions in this study. She would also like to express her thanks to all others not mentioned here. This, however, does not in anyway diminish her appreciation and gratitude to their valuable contribution to this study.

ROMZIAH SIDIK BUDIONO



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT	iii
TABLE OF CONTENTS	v
LIST OF TABLES	ix
LIST OF FIGURES	xii
LIST OF ABBREVIATIONS	xiii
ABSTRACT	xiv
CHAPTER 1 - INTRODUCTION	1
CHAPTER 2 - LITERATURE REVIEW	5
Forage Production	5
Factor Affecting Forage Yield	5
Climate	6
Soil	7
Forage Species	8
Management	9
Forage Quality	10
Forage Nutritive Value	11
Chemical Composition	12
Digestibility	15
Fermentation product.....	16
Factors Affecting Forage Nutritive Values	18
Plant Species	18
Stage of Growth	19



	Page
Soil Fertility	19
Fertilizers Rate	20
Management	21
Environment	21
<u>Setaria sphacelata</u> Variety Splendida	22
Morphology and Growth	22
Establishment, Persistence and Management	22
Yield and nutritive values	23
Toxicity factors, in <u>Setaria sphacelata</u> variety splendida	24
Utilization of <u>Setaria sphacelata</u> variety splendida	25
Grazing Management	26
Forage Intake	26
Regulation of Feed Intake	27
Measuring Feed Intake in Grazing Animals	28
Stocking Rate	29
Effects of Supplementation	30
Energy Supplement	30
Protein and Non-Protein Nitrogen Supplement	31
Mineral Supplement	32



	Page
CHAPTER 3 - EXPERIMENT I	36
THE EFFECTS OF CUTTING HEIGHTS AND CUTTING INTERVALS ON YIELD AND NUTRITIVE VALUES OF <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA	36
Introduction	36
Material and Method	37
Results.....	40
Discussion	65
CHAPTER 4 - EXPERIMENT II	
A STUDY ON THE DIGESTIBILITY OF <u>SETARIA</u> <u>SPHACELATA</u> VARIETY SPLENDIDA	66
Introduction	66
Material and Method	66
Results	68
Discussion	73
CHAPTER 5 - EXPERIMENT III	
THE EFFECT OF STOCKING RATES, GRAZING SYSTEMS AND UREA MOLASSES SUPPLEMENTAL BLOCK ON THE PERFORMANCE OF KEDAH-KELANTAN BULLS GRAZING <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA	76
Introduction	76
Material and Method	77
Results	80
Discussion	76



	Page
CHAPTER 6 - EXPERIMENT IV	
EFFECTS OF MINERAL AND PALM KERNEL MEAL SUPPLEMENT ON THE PERFORMANCE OF KEDAH- KELANTAN BULLS GRAZING <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA	103
Introduction	103
Material and Method	104
Results	106
Discussion	120
CHAPTER 7 - GENERAL DISCUSSION	124
CHAPTER 8 - SUMMARY AND CONCLUSION	126
BIBLIOGRAPHY	128
APPENDICES	



LIST OF TABLES

Table No.		Page
3.1	EFFECTS OF CUTTING HEIGHT AND INTERVAL ON DM YIELD OF LEAF, STEM AND WHOLE PLANT	42
3.2	EFFECT OF CUTTING HEIGHT AND INTERVAL ON CHEMICAL COMPOSITION AND DMD OF LEAF, STEM AND WHOLE PLANT FOR PRIMARY HARVEST	44
3.3	EFFECT OF CUTTING INTERVAL ON DM YIELD, CHEMICAL COMPOSITION AND DMD OF LEAF, STAM AND WHOLE PLANT FOR PRIMARY GROWTH	48
3.4	EFFECT OF CUTTING HEIGHT AND INTERVAL ON THE CUMULATIVE 12-WEEK DM YIELD	54
3.5	EFFECTS OF CUTTING HEIGHTS AND CUTTING INTERVAL ON CHEMICAL COMPOSITION OF LEAF, STEM AND WHOLE PLANT AVERAGED OVER THE TRIAL PERIOD	57
3.6	EFFECTS OF CUTTING INTERVAL ON DM YIELD, CHEMICAL COMPOSITION, DMD AND DDM OF LEAF, STEM AND WHOLE PLANT GROWTH AVERAGED OVER THE TRIAL PERIOD	59
4.1	CHEMICAL COMPOSITION OF FIVE WEEK OLD SUN DRIED <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA (MEAN + SD)	69
4.2	LIVE WEIGHT GAIN, <u>IN VIVO</u> DMI, APPARENT DMI, OMI, APPARENT OMI AND DIGESTIBLE OMI IN KEDAH-KELANTAN BULLS CONSUMING SUN DRIED <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA CUT AT 5-WEEK OLD	71
4.3	N RETENTION OF KEDAH-KELANTAN BULLS CONSUMING SUN DRIED <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA CUT AT 5-WEEK OLD	72
4.4	ESTIMATED ME FOR MAINTENANCE AND PRODUCTION (MJ/h/d)	72
4.5	COMPARISON BETWEEN DETERMINED DMI AND ESTIMATED DMI USING THE MAFF EQUATION	73



	Page
5.1 MEAN PERCENTAGE OF DM, CP, NDF, ASH AND DMD FOR DIFFERENT PARTS OF <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA	81
5.2 EFFECTS OF STOCKING RATES, GRAZING SYSTEMS AND STOCKING RATES ON DM,CP, NDF, ASH AND DMD	82
5.3 DRY MATTER PRODUCTION OF <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA DURING THE TRIAL (138 DAYS)	85
5.4 DIGESTIBLE DRY MATTER OF <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA DURING THE TRIAL	85
5.5 DAILY DMI PER ANIMAL UNDER VARYING GRAZING SYSTEMS AND STOCKING RATE	87
5.6 TOTAL DAILY CP INTAKE (g) PER ANIMAL UNDER VARYING GRAZING SYSTEMS AND STOCKING RATES....	88
5.7 PREDICTED DAILY ME (MJ/h/d) REQUIREMENTS FOR MAINTENANCE AND PRODUCTION PER ANIMAL UNDER VARYING GRAZING SYSTEMS AND STOCKING RATES	89
5.8 PREDICTED ME (MJ/ha) INTAKE FROM THE PASTURE FOR MAINTENANCE AND PRODUCTION UNDER VARYING GRAZING SYSTEMS AND STOCKING RATES	90
5.9 AVERAGE DAILY GAIN (g/h) OF ANIMALS DURING THE TRIAL (138 DAYS)	94
5.10 PRODUCTION OF ANIMAL PER UNIT AREA (kg/ha/d)	94
5.11 LIVE WEIGHT CHANGE OVER 138 DAYS (kg/d)	95
6.1 CHEMICAL COMPOSITION OF <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA (DM BASIS) IN THE EXPERIMENTAL PADDOCKS*	107
6.2 DMD, DM AND DDM PRODUCTION OF <u>SETARIA SPHACELATA</u> VARIETY SPLENDIDA CUMULATIVE OVER 132 DAYS IN THE EXPERIMENTAL PADDOCKS*	109

	Page
6.3	EFFECTS OF SUPPLEMENTATION ON THE DAILY DMI OF EXPERIMENTAL ANIMALS 110
6.4	EFFECTS OF SUPPLEMENTATION ON THE DAILY CP INTAKE 111
6.5	DAILY LIVE WEIGHT GAIN PER ANIMAL AND PER UNIT OF LAND 112
6.6	EFFECTS OF TREATMENTS ON TOTAL RUMINAL VFA, AMMONIA N AND PH 114
6.7	EFFECTS OF SUPPLEMENTATION ON CARCASS TRAITS* 115
6.8	EFFECTS OF SUPPLEMENTATION ON THE CONCEN- TRATION OF CA, P, MG, ZN AND CU IN BLOOD PLASMA 117
6.9	EFFECTS OF SUPPLEMENTATION ON ASH, CA, P AND MG IN BONE* 119
6.10	EFFECTS OF SUPPLEMENTATION ON CONCENTRA- TION OF CU AND MN IN THE LIVER 120



LIST OF FIGURES

Figure No.		Page
1	SCHEME OF COMPARISON BETWEEN PROXIMATE AND VAN SOEST SYSTEM OF FORAGE FRACTION	14
3.1	EFFECT OF CUTTING HEIGHTS ON MEAN DM YIELD OF LEAF AT 5 (*), 10 (□) AND 20 (●) CM CUTTING HEIGHT	41
3.2	EFFECT OF CUTTING INTERVAL ON MEAN DM YIELD OF LEAF (■), STEM (○) AND WHOLE PLANT (⊙)	50
3.3	EFFECT OF CUTTING INTERVAL ON MEAN CRUDE PROTEIN CONTENT IN LEAF (⊠), STEM (○) AND WHOLE PLANT (⊡)	51
3.4	EFFECT OF CUTTING INTERVAL ON MEAN NEUTRAL DETERGENT FIBRE (NDF) IN LEAF (⊠) AND STEM (○)	51
3.5	EFFECT OF CUTTING INTERVAL ON MEAN ASH CONTENT IN LEAF (⊠), STEM (○) AND WHOLE PLANT (⊙)	52
3.6	EFFECT OF CUTTING INTERVAL ON MEAN DRY MATTER DIGESTIBILITY (DMD) IN LEAF (⊠), STEM (○) AND WHOLE PLANT (⊙)	52
3.7	CUMULATIVE DM YIELD AND DIGESTIBLE DRY MATTER OF LEAF (L), STEM (S) AND WHOLE PLANT (W) OVER 12 WEEKS	62
5.1	LAYOUT OF THE EXPERIMENTAL PADDOCKS, ASSIGNMENTS OF ANIMALS TO TREATMENTS AND THE PATTERN OF CHANGE OVER OF THE PADDOCKS IN THE MIDDLE OF THE TRIAL	78
5.2	DRY MATTER AND DIGESTIBLE DRY MATTER OF <u>SETARIA SPHACELATA</u> VARIETY <u>SPLENDIDA</u> UNDER VARYING GRAZING SYSTEMS AND STOCKING RATES	84
5.3	RELATIONSHIP OF STOCKING RATE, PRODUCTION PER ANIMAL AND PRODUCTION PER UNIT AREA	92



LIST OF ABBREVIATION

mg	=	micro gram
g	=	gram
kg	=	kilogram
h	=	head
ha	=	hectare
m ²	=	meters square
d	=	day
MJ	=	mega joule
ml	=	mili liter
m mol	=	mili molecule
y	=	year



An abstract of the thesis presented to the Senate of Universiti
Pertanian Malaysia as partial fulfilment of the requirements
for the Degree of Doctor of Philosophy

UTILIZATION OF SETARIA SPHACELATA VARIETY SPLENDIDA
BY KEDAH-KELANTAN CATTLE

by

Romziah Sidik Budiono

May 1986

Supervisor : Professor Dr. Syed Jalaluddin bin Syed Salim

Co-Supervisor : Dr. Pravee Vijchulata

Faculty : Veterinary Medicine and Animal Science

There is limited information on beef production from pasture in Malaysia, particularly on optimization of land for maximum production. Setaria sphacelata variety splendida is a tropical grass species which has been successfully established in Malaysia as a fodder for cattle. However, optimum stocking rates and grazing systems for Setaria sphacelata variety splendida pasture for maximum productivity for beef cattle has not been established.

The objective of this study was to determine the average daily gain per animal and total gain per hectare for Kedah-Kelantan cattle, the indigenous beef cattle in Malaysia. For this purpose experiments were conducted to determine the dry



matter yield, chemical composition and digestibility using the nylon bag technique (Experiment I) and in vivo digestibility of a Setaria sphacelata variety splendida pasture (Experiment II). Later experiments examined the effects of stocking rate, grazing system and urea-mollases supplementation (Experiment III) and energy and/or mineral supplementation (Experiment IV) on daily gain and total gain/ha of Kedah-Kelantan bulls grazed on a Setaria sphacelata variety splendida pasture.

Cutting heights (5, 10 and 20 cm) had no effect on yield, chemical composition and dry matter digestibility (DMD) of Setaria sphacelata variety splendida. Crude protein and dry matter digestibility declined with increasing cutting intervals. Total dry matter (DM) yield was highest with a 12-week cutting interval. The in vivo apparent dry matter digestibility was 63.1% and the mean value for nitrogen balance was 21.4 g/h/d for bulls consuming sun dried Setaria sphacelata variety splendida cut at 5 weeks old.

Total live weight gain was significantly higher (3 kg vs 2 kg/ha/d) ($P < 0.05$) at a stocking rate of 10 animals/ha (High) than 5 animals/ha (Low) although the daily gain was significantly better on low than on the high stocking rate (425 vs 311 g/d). Type of grazing system did not influence average daily gain or total gain per hectare. Urea-mollases supplementation did not improve the daily gain of animals at



the high stocking rate. Palm kernel cake alone or in combination with mineral supplementation increased growth rates of bulls grazing continuously on a Setaria sphacelata variety splendida pasture at a stocking rate 6.25 animals/ha.

From these results it may be recommended that a 6-week cutting interval for a cut and carry system or a 5-week interval for a grazing system optimises forage yield and DDM for a Setaria sphacelata variety splendida pasture. The optimum stocking rate for this pasture grazed continuously or rotationally is 12.6 Kedah-Kelantan bulls per hectare which is equivalent to 4.6 animal unit/ha. There is little benefit of a urea-mollases supplementation at a stocking rate of 10 animals/ha. The possibility, however, exists that supplementation may be beneficial at higher stocking rates.

Abstrak tesis dikemukakan kepada Senat Universiti Pertanian
Malaysia bagi memenuhi sebahagian daripada syarat-syarat Ijazah
Falsafah Kedoktoran

UTILISASI RUMPUT SETARIA SPHACELATA
VARIETY SPLENDIDA OLEH LEMBU KEDAH-KELANTAN

oleh

ROMZIAH SIDIK BUDIONO

Mei 1986

Penyelia : Prof. Dr. Syed Jalaluddin Syed Salim
Penyelia Bersama : Dr. Pravee Vijchulata
Fakulti : Kedoktoran Veterinar dan Sains Peternakan

Di Malaysia, terdapat keterangan yang kurang tentang pengeluaran lembu pedaging berhubung dengan penggunaan padang rumput terutamanya dalam menentukan penggunaan padang yang optima untuk pengeluaran yang paling tinggi. Setaria sphacelata variety splendida adalah sejenis rumput di kawasan tropik yang telah berhasil ditanam di Malaysia untuk digunakan sebagai bahan makanan lembu. Walau bagaimana pun kadar muatan yang optimum serta sistem ragutan untuk menghasilkan pengeluaran daging yang tinggi dari rumput Setaria sphacelata variety splendida belum diselidiki.

Tujuan penyelidikan ini adalah untuk menentukan purata kenaikan berat badan setiap ekor lembu dan kenaikan berat badan



yang terkumpul dari setiap hektar padang rumput. Bagi maksud ini, eksperimen-eksperimen telah dijalankan dalam menentukan bahan kering, kadar bahan kimia dan nilai pencernaan dengan cara menggunakan teknik bag nylon (Eksperimen I) dari rumput Setaria sphacelata variety splendida. Eksperimen seterusnya ialah untuk mengkaji kesan kadar muatan, sistem ragutan dan pemberian makanan tambahan yang berupa campuran urea dan molasses (Eksperimen III) dan pemberian tambahan tenaga dan/atau mineral (Eksperimen IV) terhadap kenaikan berat badan satu ekar atau satu hektar lembu Kedah-Kelantan yang meragut rumput Setaria sphacelata variety splendida.

Paras pemotongan (5, 10 dan 20 cm) tidak memberi kesan pada hasil bahan kering, kadar bahan kimia dan nilai pencernaan rumput Setaria sphacelata variety splendida. Protin kasar dan nilai pencernaan menurun dengan meningkatnya jarak waktu pemotongan. Bahan kering yang terkumpul sangat tinggi pada jarak pemotongan setiap 12 minggu. Nilai pencernaan sebenar adalah 63.1% dan nilai purata keseimbangan nitrogen sebesar 21.4 g/ekor/hari untuk lembu jantan yang makan rumput kering Setaria sphacela variety splendida yang dipotong semasa berumur 5 minggu.

Kenaikan berat badan yang terkumpul nyata tertinggi (3 dibandingkan dengan 2 kg/ha/hari) ($P < 0.05$) pada tingkat kadar muatan 10 ekor/ha (tinggi) dibandingkan dengan 5 ekor/ha (rendah) walaupun kenaikan berat badan nyata lebih baik pada

kadar muatan yang rendah dibandingkan dengan kadar muatan yang tinggi (425 dibandingkan dengan 311 g/hari). Jenis sistem ragutan tidak mempengaruhi nilai purata kenaikan berat badan satu ekor atau satu hektar. Makanan tambahan urea-molasses tidak dapat meningkatkan kenaikan berat badan lembu yang dipelihara dengan kadar muatan tinggi (10 ekor/ha). Bungkil kelapa sawit sahaja atau campuran mineral dengan bungkil kelapa sawit dapat meningkatkan kecepatan pertumbuhan lembu jantan yang meragut rumput Setaria sphacelata variety splendida secara berterusan dengan kadar muatan 6.25 ekor/ha.

Hasil penyelidikan menunjukkan bahawa jarak pemotongan setiap 6 minggu sekali dapat dilaksanakan untuk sistem potong dan bawa atau jarak pemotongan setiap 5 minggu dapat dilaksanakan untuk sistem ragutan bagi mendapatkan nilai optima daripada hasil perumputan dan bahan kering yang tercerna bagi rumput Setaria sphacelata variety splendida. Nilai optimum kadar muatan untuk padang rumput yang diragut secara berterusan atau bergilir adalah 12.6 ekor/ha atau sama dengan 4.6 ekor unit/hektar. Hanya sedikit keuntungan yang didapati daripada makanan tambahan urea dan molasses pada lembu dengan kadar muatan 10 ekor/ha. Kemungkinan hal ini dapat menguntungkan apabila pemberian makan diberikan pada lembu dengan kadar muatan yang lebih tinggi lagi.



CHAPTER I

INTRODUCTION

Ruminants are capable of digesting cell wall components of grass through microbial functions, therefore grass become the primary source of feed for cattle. Grass quality and output of production from cattle are closely related. The animals require nutrients from the grasses for body maintenance, work, reproduction of meat and milk, etc. Grasses itself take nutrients from the soil.

Efficient animal production from pasture is basically determined by the intake of forage, the nutritive values of the ingested materials and the availability of the forage biomass. Dry matter yield of the forage is negatively correlated to its feed values (Raymond, 1969; Moore, 1980). While biomass produced per unit area increases with maturity, its nutritive values, especially, crude protein and digestibility fall. Hence, forage consumption by animals decline.

Stocking rate and grazing system, are important management factors that may influence production per animal and total animal production per unit area of pasture land. Stocking rate affects individual animal performance by an amount which will vary from year to year and from one season of the year to another (Morley, 1981). The optimum stocking rate can be



estimated by predicting the relationship between stocking rate and animal production per head, per hectare in terms of live weight gain based on an optimum grazing pressure (Mott, 1960).

If the pasture is undergrazed or overgrazed the animal production per head will be reduced and potential production of an animal underestimated (Mott, 1960). When there is a good supply of pasture, stocking rate has little effect on individual animal performance, since there is enough fodder for each animal. As the stocking rate of the pasture increased, the individual animal output starts to decrease, as animals begin to compete with each other for the available forage supply and have less opportunity to select the most nutritious parts of the pastures (Humphreys, 1978).

The optimum stocking rate of a pasture, however, varies according to the specific grass species available together with its managements. Since different breeds of animal may impose different demands on pasture because their feed requirements may differ and they may graze in different ways (Morley and Speeding, 1968) type, species and physiological difference of the animals grazed on the pasture can also influence the optimum carrying capacity of the paddock.

In situations where nutrients available from the pasture cannot meet the requirement of the animals, improvement can be obtained by supplementing the grazing animals with energy, protein or non protein nitrogen (Alden, 1981) and/or minerals (McDowell and Conrad, 1977).



Proper management of the pasture for compromising the various factors is therefore essential and is found to be specific for individual forage species (Moore and Mott, 1973; Whiteman, 1980).

Setaria Sphacelata is a tropical grass which grows well in areas of reasonably high rainfall, usually over 700 mm (Luck, 1979). Although dry matter of leaves is low and crude protein content averages at 10%, the forage is palatable. The grass regenerates rapidly after grazing (defoliation) or cutting (Bogdan, 1979F). This grass has been successfully introduced into Malaysia as a forage crop for cattle. The information on its nutritive values and its potential as a pasture grass under various grazing management system for Kedah-Kelantan (KK) cattle, is rather limited.

Kedah-Kelantan cattle are the main indigenous beef cattle breed in Malaysia. The breed is well adapted to the environmental conditions. They are used mainly for beef and draught (Devendra et al, 1973). Although the breed is known for its high fertility, the live weight gain of KK cattle, however, is relatively low in comparison to the improved beef breeds such as Brahman or Droughtmaster (Devendra, 1973). It is possible that the low daily live weight gain of KK cattle may, in part, reflect the substandard nutrition and management. Therefore, with proper management and nutrition, daily live weight gain of this breed might be improved.

