

UNIVERSITI PUTRA MALAYSIA

PATHOLOGICAL EFFECTS RESULTING FROM
BIOACCUMULATION OF CRUDE OIL ON IMMUNOLOGICAL
PARAMETERS IN RED SNAPPER (*LUTJANUS*ARGENTIMACULATUS) FORSSAL AND BLACK TIGER SHRIMP
(PENAEUS MONODOM) FABRICIUS

SOHRAB AKBARI

FPV 1997 7



PATHOLOGICAL EFFECTS RESULTING FROM BIOACCUMULATION OF CRUDE OIL ON IMMUNOLOGICAL PARAMETERS IN RED SNAPPER (LUTJANUS ARGENTIMACULATUS) FORSSAL AND BLACK TIGER SHRIMP (PENAEUS MONODON) FABRICIUS

BY

SOHRAB AKBARI

Dissertation Submitted in Fulfilment of the Requirements for the Degree of Doctor of Philosophy in the Faculty of Veterinary Medicine and Animal Science, Universiti Putra Malaysia

June 1997



To all fellow mankind, who believe in God, and in doing good deeds



ACKNOWLEDGEMENTS

I have completed this programme, because there were those who gave me support and assistance. There are many of them whom I cannot thank enough; yet I cannot solely lay claim to this success without naming a few of them.

I would like to thank Prof. Dr. Mohammed Shariff, my committee chairman, whose extraordinary help provided me such occasion to pursue my work with this great passion.

I am also thankful to Prof. Dr. Law Ah Theem, a member of my supervisory committee for his valuable discussions on toxicity of crude oil and going through the manuscript.

My gratitude also goes to Dr. Abd. Rahman Bin Manas, a member of my supervisory committee, whose expertise gave me enough impetus to go through crude oil analysis.

I am likewise grateful to Dr. John Foo Tet Wei, Department of Zoology, National University of Singapore, for his technical help in cortisol radioimmunoassay.

I would also like to thank technical staff members of Gas Chromatography and Gas Chromatography-Mass Spectrometer



unit, Chemistry Department, Faculty of Science, and Electron Microscope Unit, Faculty of Veterinary Medicine and Animal Science, University Putra Malaysia for their technical assistance.

I will be fail in my duty if I do not record my thanks to Shiraz University, Shiraz, Iran for their encouragement and financial support in my programme.

I am deeply indebted and grateful to Ayatollah H. A. Nejabat whose exceptional ethos on Allah provides me much inspiration and motivation to pursue my daily activities. My personal thanks are also due to my wife Z. Nejabat, my son Mohammed Javad, and my daughters Razih, Maryam, and Marzih who prayed to Allah for me to have this favourable outcome in my study.



TABLE OF CONTENTS

		Page
ACKN	OWLEDGEMENTS	iii
LIST	OF TABLES	x
LIST	OF FIGURES	xxvi
LIST	OF PLATES	xxx
LIST	OF SPECIES	xxxv
LIST	OF ABBREVIATIONS	xxxviii
ABST	RACT	хl
ABST	RAK	xliv
CHAP	TER	
I	GENERAL INTRODUCTION	1
II	REVIEW OF LITERATURE	9
	What is Crude Oil?	9
	Source of Oil Pollution	12
	Fate of Spilled Oil	14
	Uptake, Biotransformation and Effects of Petroleum Hydrocarbons in Fish and Invertebrates	18
	Uptake and Bioaccumulation of	10
	Petroleum Hydrocarbons	19
	Biotransformation	22
	Effects of Petroleum Hydrocarbons	26
	Factors Modulating Uptake, Biotransformation and Effects of Petroleum Hydrocarbons in Aquatic Animals	43
	THE BOURDLINE BUILDING IS A A A A A A A A A A A A A A A A A A	4)



	Used in the Study	48
	Research Problems	50
III	ACUTE TOXICITY OF WATER SOLUBLE FRACTION OF CRUDE OIL TO RED SNAPPER (LUTJANUS ARGENTIMACULATUS) AND BLACK TIGER SHRIMP (PENAEUS MONODON)	52
	Materials and Methods	54
	Collection and Maintenance of Experimental Animals	54
	Fish	54
	Shrimp	55
	Test Oil and Exposure Apparatus	56
	Hydrocarbon Analysis	58
Reco	very of Hydrocarbons 64	
	Treatment of Discharged Oily Seawater	65
	Method for Estimation of LC ₅₀ Values of Crude Oil	65
	Results	66
	Adjustment of the Flow-through System	66
	Analysis of Crude Oil	67
	Estimated Values of LC ₅₀	76
	Discussion	83
IV	ACCUMULATION OF NAPHTHALENES IN LIVER AND MUSCLE OF RED SNAPPER (LUTJANUS ARGENTIMACULATUS) AND IN HEAD AND ABDOMEN OF BLACK TIGER SHRIMP (PENAEUS MONODON) EXPOSED TO SUBLETHAL CONCENTRATIONS OF CRUDE OIL.	91
	Materials and Methods	94
	Collection and Maintenance of Test	9.4



	r 1511	24
	Shrimp	94
	Test Oil and Exposure Apparatus	95
	Sampling of Test Animals	96
	Fish	96
	Shrimp	97
	Analysis of Naphthalenes in the Target Tissues	97
	Analysis of Naphthalenes in Exposure Seawater	99
	Recovery of Naphthalenes in the Tissue Samples	99
	Statistical Analysis	100
	Results	100
	Accumulation of Naphthalenes in the Fish Tissues	103
	Accumulation of Naphthalenes in the Shrimp Tissues	107
	Comparison Between Equilibrium Con- centrations of Naphthalenes in the Tissues of Experimental Animals	111
	Discussion	116
V	EFFECTS OF SUBLETHAL CONCENTRATIONS OF CRUDE OIL ON CORTISOL, BIOCHEMICAL AND CELLULAR INDICES OF RED SNAPPER (LUTJANUS ARGENTIMACULATUS) BLOOD	122
	Materials and Methods	125
	Haematological Study	125
	Measurement of Cortisol Levels in plasma	129
	Statistical Analysis	130
	Results	131
	Cortisol and Biochemical Parameters	131



	Red Blood Cells (RBC) and Related Parameters	136
	White Blood Cells (WBC) and Diff- erential Cell Counts	139
	Discussion	145
VI	EFFECTS OF SUBLETHAL CONCENTRATIONS OF CRUDE OIL ON PHENOLOXIDASE, BIOCHEMICAL AND CELLULAR INDICES OF BLACK TIGER SHRIMP (PENAEUS MONODON) HAEMOLYMP	157
	Materials and Methods	160
	Total and Differential Haemocyte Count	160
	Assay for Phenoloxidase (PO) Activity	162
	Statistical Analysis	163
	Results	164
	Total Haemocyte (THC) and Differential Cell Counts	164
	Changes in Phenoloxidase Activities	169
	Discussion	173
VII	EFFECTS OF SUBLETHAL CONCENTRATIONS OF CRUDE OIL ON CELLULAR AND ULTRACELLULAR STRUCTURES OF IMMUNOLOGICALLY IMPORTANT TISSUES OF BLACK TIGER SHRIMP (PENAEUS MONODON)	182
	Material and Methods	185
	Fixation of Samples	185
	Light Microscopy	185
	Transmission and Scanning Electron Microscopy	186
	Tissue Processing	186
	Light Microscopy	186
	Transmission and Scanning Electron	105



Results	188
Physiological Behaviour and Gross Appearance	188
Histopathological Alterations	188
Discussion	219
VIII GENERAL DISCUSSION AND RECOMMENDATION	231
BIBLIOGRAPHY	245
APPENDICES	266
Part A (Original Data)	267
Part B (Statistics)	310
Part C (Standard and Control Curves)	360
BIOGRAPHICAL SKETCH	366



LIST OF TABLES

Table	e	Pag
1	Estimated World Input of Petroleum Hydrocarbons to the Sea (Mt/Year)	13
2	Solubility of Polyaromatic Hydrocarbons in Distilled Water at 25-27°C	16
3	Pathways for the Environment Fate of Crude Oil	18
4	Comparative Acute Toxicity, Measured as 48 h LC ₅₀ , of Water Soluble Fraction and Oil in Water Dispersions of Three Oils to Several Species of Marine Animals	31
5	Reported Values on the Changes in Cortisol, Biochemical and Cellular Parameters of Fish Exposed to Petroleum Hydrocarbons	36
6	Reported Values on the Changes in Phenol- oxidase, Biochemical and Cellular Par- ameters of invertebrates Exposed to diff- erent stressors	39
7	Percentages of Water Soluble Fractions of Crude Oil Before Its Exposure to the Test Animals in Experimental Tanks for Estimation LC ₅₀ Values	66
8	Standard Aromatic Hydrocarbons and Their Recovery Factors (Mean ± SD), Used for Quantification Analysis of Water soluble Fractions in Exposure Seawater	73
9	Standard Aliphatic Hydrocarbons and Their Recovery Factors (Mean ± SD), Used for Quantification Analysis of Water Soluble Fractions in Exposure Seawater	75
10	Concentrations of Soluble Aliphatic and Aromatic Hydrocarbons in the Exposed Water, Used for Estimation of LC50 Values for L. argentimaculatus and P. monodon. Concentrations Are in mg/l	77
	Concentrations are in mg/1000000000000000000000000000000000000	



11	Mortality of <i>L. argentimaculatus</i> Exposed to Water Soluble Fraction of Crude Oil During 96 h	78
12	Mortality of <i>P. monodon</i> Exposed to Water Soluble Fraction of Crude Oil During 96 h.	78
13	Estimated LC ₅₀ Values of Water Soluble Fraction of Malaysian Tapis Blend Crude Oil for <i>L. argentimaculatus</i> and <i>P. monodon</i> , using Trimmed Spearman-Karber Method	79
14	Water Quality Parameters in the Exposure Tanks of <i>L. argentimaculatus</i> and <i>P. monodon</i> During 96 h Acute Toxicity Test	82
15	Quantitative Analysis of Naphthalenes in the Exposed Seawater During Sublethal Toxicity Test in L. argentimaculatus and P. monodon	101
16	Standard Naphthalenes and Their Recovery Factors (Mean ± SD), Used for Quantification Analysis of Naphthalenes in Tissue Samples of P. monodon and L. argentimaculatus	102
17	Water Quality Parameters in the Exposure Tanks of <i>L. argentimaculatus</i> and <i>P. monodon</i> During Sublethal Toxicity Test	115
18	The Results of Histopathological Changes in different Organs of <i>P. monodon</i> Exposed to Three Different Concentrations of Crude Oil	218
19	General Properties of Malaysian Tapis Blend Crude Oil	267
20	Aromatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Red Snapper (Lutjanus argentimaculatus) (Tank No. C2)	268
21	Aromatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Red Snapper (Lutjanus argentimaculatus) (Tank No. C3)	269
22	Aromatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Red Snapper (Lutjanus argentimaculatus)	270



23	posed Water During Acute Toxicity Test of Red Snapper (Lutjanus argentimaculatus) (Tank No. C5)	271
24	Aromatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Red Snapper (Lutjanus argentimaculatus) (Tank No. C6)	272
25	Aromatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Black Tiger Shrimp (Penaeus monodon) (Tank No. C2)	273
26	Aromatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Black Tiger Shrimp (Penaeus monodon) (Tank No. C3)	274
27	Aromatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Black Tiger Shrimp (Penaeus monodon) (Tank No. C4)	275
28	Aromatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Black Tiger Shrimp (Penaeus monodon) (Tank No. C5)	276
29	Aromatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Black Tiger Shrimp (Penaeus monodon) (Tank No. C6)	277
30	Aliphatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Red Snapper (Lutjanus argentimaculatus) (Tank No. C2)	278
31	Aliphatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Red Snapper (Lutjanus argentimaculatus) (Tank No. C3)	278
32	Aliphatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Red Snapper (Lutjanus argentimaculatus) (Tank No. C4)	279
33	Aliphatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Red Snapper (Lutjanus argentimaculatus)	270



34	posed Water During Acute Toxicity Test of Red Snapper (Lutjanus argentimaculatus) (Tank No. C6)	280
35	Aliphatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Black Tiger Shrimp (Penaeus monodon) (Tank No. C2)	281
36	Aliphatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Black Tiger Shrimp (Penaeus monodon) (Tank No. C3)	281
37	Aliphatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Black Tiger Shrimp (Penaeus monodon) (Tank No. C4)	282
38	Aliphatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Black Tiger Shrimp (Penaeus monodon) (Tank No. C5)	282
39	Aliphatic Hydrocarbon Fractions in the Exposed Water During Acute Toxicity Test of Black Tiger Shrimp (Penaeus monodon) (Tank No. C6)	283
40	Estimated LC ₅₀ Values of Water Soluble Fraction of Crude Oil for Red Snapper (<i>Lutjanus argentimaculatus</i>), Using Trimmed Spearman-Karber Method	284
41	Estimated LC ₅₀ Values of Water Soluble Fraction of Crude Oil for Black Tiger Shrimp (Penaeus monodon), Using Trimmed Spearman- Karber Method	285
42	Quantitative Analysis of Naphthalenes (mg) in 500 ml of Seawater During Sublethal Toxicity Test of Red Snapper (Lutjanus argentimaculatus)	286
43	Quantitative Analysis of Naphthalenes (mg) in 500 ml of Seawater During Sublethal Toxicity Test of Black Tiger Shrimp (Penaeus monodon)	287
44	Accumulation of Naphthalenes (mg/10g wet samples) in the liver of Red Snapper (Lutjanus argentimaculatus) During Sub-	200



45	Accumulation of Naphthalenes (mg/10g wet samples) in the Muscle of Red Snapper (Lutjanus argentimaculatus) During Sublethal Toxicity Test	289
46	Accumulation of Naphthalenes (mg/10g wet samples) in the Head Tissues of Black Tiger Shrimp (Penaeus monodon) During Sublethal Toxicity Test	290
47	Accumulation of Naphthalenes (mg/10g wet samples) in the Muscle of Black Tiger Shrimp (Penaeus monodon) During Sublethal Toxicity Test	291
48	Fluctuation of Plasma Cortisol (ng/ml of plasma) in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	292
49	Fluctuation of Plasma Glucose (mg/100 ml of plasma) in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	293
50	Fluctuation of Plasma Cholesterol (mg/100 ml of plasma) in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test.	294
51	Fluctuation of Total Plasma Protein (g/100 ml of plasma) in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	295
52	Fluctuation of Red Blood Cell Counts (10 /ml of blood) in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	296
53	Fluctuation of the Size of Red Blood Cell in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	297
		201



54	of blood) in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	298
55	Fluctuation of Haematocrit (%) Values in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	299
56	Fluctuation of White Blood Cells in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	300
57	Fluctuation of the Percentage of Blood Lymphocytes in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	301
58	Fluctuation of the Percentage of Blood Thrombocytes in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	302
59	Fluctuation of the Percentage of Blood Neutrophils in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	303
60	Fluctuation of the Total Haemocyte Counts in Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 386 h Sublethal Toxicity Test	304
61	Fluctuation of the Percentage of Granular Cells in Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 386 h Sublethal Toxicity Test	305



62	Fluctuation of the Percentage of Semi- granular Cells in Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 386 h Sublethal Toxi- city Test	306
63	Fluctuation of the Percentage Hyaline Cells in Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 386 h Sublethal Toxicity Test	307
64	Fluctuation of the Haemocyte Phenoloxidase Activity (A ABC/min/mg protein) in Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 386 h Sublethal Toxicity Test	308
65	Fluctuation of the Total Plasma Protein in mg/ml of plasma in Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 386 h Sublethal Toxicity Test	309
66	Regression Analysis of Concentration-Mortality Response for Fish (<i>Lutjanus argentimaculatus</i>) at 12 h Post Exposure in Acute Toxicity Test	310
67	Regression Analysis of Concentration-Mortality Response for Shrimp (<i>Penaeus monodon</i>) at 12 h Post Exposure in Acute Toxicity Test	310
68	Regression Analysis of Concentration- Mortality Response for Fish (Lutjanus argentimaculatus) at 48 h Post Exposure in Acute Toxicity Test	311
69	Regression Analysis of Concentration- Mortality Response for Shrimp (<i>Penaeus monodon</i>) at 48 h Post Exposure in Acute Toxicity Test	311
70	Regression Analysis of Concentration- Mortality Response for Fish (<i>Lutjanus</i> argentimaculatus) at 96 h Post Exposure in Acute Toxicity Test	312
71	Regression Analysis of Concentration-Mortality Response for Shrimp (<i>Penaeus monodon</i>) at 96 h Post Exposure in Acute Toxicity Test	312



Multiple Range Analysis for Bioaccumulation of Naphthalenes in the Liver of Red Snapper (Lutjanus argentimaculatus) Exposed to Three Different Concentrations, a Dose-response Analysis	12	Analysis of Variance for Bloaccumulation of Naphthalenes in the Liver of Red Snapper (Lutjanus argentimaculatus) Exposed to Three Different Concentrations of Naphthalenes During 96 h Sublethal Toxicity Test	313
tion of Naphthalenes in the Liver of Red Snapper (Lutjanus argentimaculatus) Exposed to Naphthalenes During 96 h Sublethal Toxicity Test, a Time-response Analysis. 314 75 Analysis of Variance for Bioaccumulation of Naphthalenes in the Muscle of Red Snapper (Lutjanus argentimaculatus) Exposed to Three Different Concentrations of Naphthalenes During 96 h Sublethal Toxicity Test	73	tion of Naphthalenes in the Liver of Red Snapper (Lutjanus argentimaculatus) Exposed to Three Different Concentrations,	313
tion of Naphthalenes in the Muscle of Red Snapper (Lutjanus argentimaculatus) Exposed to Three Different Concentrations of Naphthalenes During 96 h Sublethal Toxicity Test	74	tion of Naphthalenes in the Liver of Red Snapper (Lutjanus argentimaculatus) Ex- posed to Naphthalenes During 96 h Subleth-	314
tion of Naphthalenes in the Muscle of Red Snapper (Lutjanus argentimaculatus) Exposed to Three Different Concentrations, a Dose-response Analysis	75	tion of Naphthalenes in the Muscle of Red Snapper (Lutjanus argentimaculatus) Ex- posed to Three Different Concentrations of Naphthalenes During 96 h Sublethal	315
tion of Naphthalenes in the Muscle of Red Snapper (Lutjanus argentimaculatus) Exposed to Naphthalenes During 96 h Sublethal Toxicity Test (Level), a Time-response Analysis	76	tion of Naphthalenes in the Muscle of Red Snapper (Lutjanus argentimaculatus) Ex- posed to Three Different Concentrations,	315
tion of Naphthalenes in the Head Tissues of Black Tiger Shrimp (Penaeus monodon) Exposed to Three Different Concentrations of Naphthalenes During 384 h Sublethal Toxicity Test	77	tion of Naphthalenes in the Muscle of Red Snapper (Lutjanus argentimaculatus) Exposed to Naphthalenes During 96 h Sublethal Toxicity Test (Level), a Time-response	316
tion of Naphthalenes in the Head Tissues of Black Tiger Shrimp (Penaeus monodon) Exposed to Three Different Concentrations,	78	tion of Naphthalenes in the Head Tissues of Black Tiger Shrimp (Penaeus monodon) Exposed to Three Different Concentrations of Naphthalenes During 384 h Sublethal	317
	79	tion of Naphthalenes in the Head Tissues of Black Tiger Shrimp (Penaeus monodon)	317



80	Multiple Range Analysis for Bioaccumulation of Naphthalenes in the Head Tissues of Black Tiger Shrimp (Penaeus monodon) Exposed to Naphthalenes During 384 h Sublethal Toxicity Test, a Time-response Analysis	318
81	Analysis of Variance for Bioaccumulation of Naphthalenes in the Muscle of Black Tiger Shrimp (Penaeus monodon) Exposed to Three Different Concentrations of Naphthalenes During 384 h Sublethal Toxicity Test	319
82	Multiple Range Analysis for Bioaccumulation of Naphthalenes in the Muscle of Black Tiger Shrimp (Penaeus monodon) Exposed to Three Different Concentrations, a Dose-response Analysis	319
83	Multiple Range Analysis for Bioaccumulation of Naphthalenes in the Muscle of Black Tiger Shrimp (Penaeus monodon) Exposed to Naphthalenes During 384 h Sublethal Toxicity Test, a Time-response Analysis	320
84	Analysis of Variance for Fluctuations of Plasma Cortisol in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	321
85	Multiple Range Analysis for Fluctuations of Plasma Cortisol in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Doseresponse Analysis	321
86	Multiple Range Analysis for Fluctuations of Plasma Cortisol in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Time-response Analysis	322
87	Analysis of Variance for Fluctuations of Plasma Glucose in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity	
	Test	323



00	Plasma Glucose in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Dose-response Analysis	323
89	Multiple Range Analysis for Fluctuations of Plasma Glucose in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Time-response Analysis	324
90	Analysis of Variance for Fluctuations of Plasma Cholesterol in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	325
91	Multiple Range Analysis for Fluctuations of Plasma Cholesterol in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Dose-response Analysis	325
92	Multiple Range Analysis for Fluctuations of Plasma Cholesterol in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Time-response Analysis	326
93	Analysis of Variance for Fluctuations of Total Plasma Protein in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	327
94	Multiple Range Analysis for Fluctuations of Total Plasma Protein in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Dose-response Analysis	327
		321



95	of Total Plasma Protein in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Time-response Analysis	328
96	Analysis of Variance for Fluctuations of Red Blood Cell Counts in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	329
97	Multiple Range Analysis for Fluctuations of Red Blood Cell Counts in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Dose-response Analysis	330
98	Multiple Range Analysis for Fluctuations of Red Blood Cell Counts in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Time-response Analysis.	331
99	Analysis of Variance for Fluctuations of Red Blood Cell Size in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	331
100	Multiple Range Analysis for Fluctuations of Red Blood Cell Size in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Dose-response Analysis.	332
101	Multiple Range Analysis for Fluctuations of Red Blood Cell Size in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Time-response	
	Analysis	332



102	Haemoglobin in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	333
103	_	333
104	Analysis of Variance for Fluctuations of Haematocrit in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	334
105	Multiple Range Analysis for Fluctuations of Haematocrit in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Dose-response Analysis	334
106	Analysis of Variance for Fluctuations of White Blood Cell Counts in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	335
107	Multiple Range Analysis for Fluctuations of White Blood Cell counts in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96h Sublethal Toxicity Test, a Dose-response Analysis	335
108	Multiple Range Analysis for Fluctuations of White Blood Cell counts in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude oil During 96 h Sublethal Toxicity Test, a Time-response	
	Analysis	336



109	Analysis of Variance for Fluctuations of Lymphocytes in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	337
110	Multiple Range Analysis for Fluctuations of Lymphocytes in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96h Sublethal Toxicity Test, a Dose-response Analysis	337
111	Multiple Range Analysis for Fluctuations of Lymphocytes in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Time-response Analysis	338
112	Analysis of Variance for Fluctuations of Thrombocytes in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	339
113	Multiple Range Analysis for Fluctuations of Thrombocytes in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations (Level) of Water Soluble Fractions of Crude Oil During 96h Sublethal Toxicity Test, a Dose-response Analysis	339
114	Multiple Range Analysis for Fluctuations of Thrombocytes in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Time-response Analysis	340
115	Analysis of Variance for Fluctuations of Neutrophils in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 96 h Sublethal Toxicity Test	341
116	Multiple Range Analysis for Fluctuations of Neutrophils in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Dose-response Analysis	341
	- · · · · · · · · · · · · · · · · · · ·	



117	Multiple Range Analysis for Fluctuations of Neutrophils in Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test, a Time-response Analysis	342
118	A Correlation Analysis of Transformed (square root of Y+0.5) variation in Blood Parameters and Bioaccumulation of Naphthalenes in Liver and Muscle of Red Snapper (Lutjanus argentimaculatus) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 96 h Sublethal Toxicity Test	343
119	Analysis of Variance for Fluctuations of Haemocytes of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 384 h (Time) Subacute Toxicity Test	346
120	Multiple Range Analysis for Fluctuations of Haemocytes of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 384 h Sublethal Toxicity Test, a Dose-response Analysis	346
121	Multiple Range Analysis for Fluctuations of Haemocytes of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 384 h sublethal Toxicity Test, a Time-response Analysis	347
122	Analysis of Variance for Fluctuations of Granular Cells of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 384 h Subacute Toxicity Test	348
123	Multiple Range Analysis for Fluctuations of Granular Cells of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 384 h sublethal Toxicity Test, a Dose-response Analysis	348



131	Analysis of Variance for Fluctuations of Haemocyte Phenoloxidase Activity of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 384 h Subacute Toxicity Test	354
132	Multiple Range Analysis for Fluctuations of Haemocyte Phenoloxidase Activity of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 384 h sublethal Toxicity Test, a Dose-response Analysis	354
133	Multiple Range Analysis for Fluctuations of Haemocyte Phenoloxidase Activity of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 384 h sublethal Toxicity Test, a Timeresponse Analysis	355
134	Analysis of Variance for Fluctuations of Total Plasma Protein of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fraction of Crude Oil During 384 h Subacute Toxicity Test	356
135	Multiple Range Analysis for Fluctuations of Total Plasma Protein of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 384 h sublethal Toxicity Test, a Dose-response Analysis	356
136	Multiple Range Analysis for Fluctuations of Total Plasma Protein of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 384 h sublethal Toxicity Test, a Time-response Analysis	357
137	A Correlation Analysis of Transformed (square root of Y+0.5) Variations of Haemolymph Parameters and Bioaccumulation of Naphthalenes in the Head and Muscle of Black Tiger Shrimp (Penaeus monodon) Exposed to Different Concentrations of Water Soluble Fractions of Crude Oil During 384 h Sublethal Toxicity Test	358

