

ORIGINAL ARTICLE

Allergic Rhinitis and Atopic Dermatitis in Association with Home Environment among Adolescents from Urban and Suburban Areas

Khairul Nizam Mohd Isa¹, Juliana Jalaludin^{2,3*}, Jamal Hisham Hashim⁴, Zailina Hashim²

¹ Environmental Health Research Cluster (EHRC), Environmental Healthcare Section, Institute of Medical Science Technology, Universiti Kuala Lumpur, 43000 Kajang, Selangor, Malaysia

² Department of Environmental and Occupational Health, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, UPM, 43400 Serdang, Selangor, Malaysia

³ Emerging Pollutant, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

⁴ Department of Health Sciences, Faculty of Engineering and Life Science, Universiti Selangor, Shah Alam Campus, Seksyen 7, 40000 Shah Alam, Selangor, Malaysia

ABSTRACT

Introduction: This study aimed to determine the associations between home environment exposures and allergic rhinitis (AR) and atopic dermatitis (AD) among adolescents living in Hulu Langat, Selangor, Malaysia. **Method:** This comparative cross-sectional study involved adolescents residing in both suburban and urban areas within Hulu Langat district. Information on allergic rhinitis and home environment exposure was gathered from 546 adolescents aged 14-year-old using the International Study of Asthma and Allergies in Children (ISAAC) questionnaire. **Results:** The prevalence of AR and AD for urban and suburban adolescents were 59.8% and 40.2%, and 64.9% and 35.1%, respectively. A logistic regression analysis showed that frequency ETS exposure (OR=1.12, 95% CI= 1.02-1.24), furry pet (OR=1.20, 95%CI= 1.17-1.24), indoor painting (OR=1.26, 95%CI= 1.01-1.61), floor renovation (OR=1.70, 95%CI= 1.25-2.32), water leakage (OR=1.22, 95%CI= 1.19-1.26) and mold odor in the last 12 months (OR=1.51, 95%CI= 1.06-2.15) were associated with AR among adolescents living in urban areas. Nevertheless, floor renovation (OR=1.91, 95%CI= 1.01-4.04), water leakage exposure (OR=1.81, 95%CI= 1.03-3.19) and other odor last 12 months (OR=2.37, 95%CI= 1.32-4.25) exposures were linked to an increased likelihood of AR among adolescents from the suburban area. **Conclusion:** The findings suggest that various home environmental exposures play a significant role in the occurrence of AR among adolescents, emphasizing the need for improved home environment management to reduce allergy-related health risks.

Malaysian Journal of Medicine and Health Sciences (2026) 22(SUPP3): 62-68. doi:10.47836/mjmhs.22.s3.10

Keywords: Allergic rhinitis, atopic dermatitis, adolescents, urban, suburban

Corresponding Author:

Juliana Jalaludin, PhD

Email: juliana@upm.edu.my

Tel: +603-9769 2401

INTRODUCTION

With a rapid development of economy and infrastructure in urban and suburban regions, allergic diseases have increased worldwide. In developed countries, the prevalence of allergic rhinitis (AR) and atopic dermatitis (AD) estimated in the range from 12% to 20% and 0% to 24%, respectively [1,2]. Review summarised that 2.23% of the world population were suffered from AD in year 2019 [3]. Furthermore, a review estimated the prevalence of AR in Malaysia to be 7.1%, with approximately 16.1% among adolescents aged 13-14 years [4]. Allergic diseases exert a significant clinical

and socioeconomic burden on individuals. Studies have demonstrated that AR and AD are linked to a higher likelihood of developing various respiratory diseases conditions like asthma, airway hyperreactivity (AHR), and chronic obstructive pulmonary disease (COPD) [5].

Numerous reviews have shown that exposure to home environmental factors such as indoor painting, renovations, water leakage and dampness, mold, environmental tobacco smoke (ETS) exposure, pets and indoor pollutants can influence the development of AR and AD in adolescents [6,7]. However, the association between these home environmental factors and prevalence of AR and AD in urban and suburban settings has not been widely studied [8–10]. This scarcity of research on the effects of home environmental factors in urban and suburban environments on AR and AD provides an opportunity for further investigation into

these relationships. Moreover, review reported that geographical differentiation influenced the development and exacerbation of AR and AD among children and adults [11]. Additionally, urbanization, lifestyle, home environmental factors, and climate variations can significantly impact these conditions differently across locations and regions [12]. To the best of our knowledge, no previous study has been conducted in tropical country to determine the association between home environmental factors and AR and AD among adolescents from both urban and suburban areas. Therefore, understanding the relationship between these factors is crucial for assessing the impact in the society, planning health service, implementing preventive strategies.

This study aimed to determine the associations between home environment exposures and AR and AD among adolescents living in Hulu Langat Selangor. Investigating these associations in a Malaysian setting, characterized by its diverse urban-suburban landscape and ongoing development, will provide valuable insights into the specific risk factors relevant to this population.

METHODOLOGY

Study population

This cross-sectional study included 547 fourteen-year-old adolescents from 8 secondary schools located in the Hulu Langat district, Selangor, Malaysia. This district was selected due to its diverse urban and suburban regions and ongoing industrial and property development projects. Several areas within Hulu Langat (Kajang, Ampang Jaya, and Batu 9 Cheras) are classified as urban, with a total population of approximately 882,000 in 2010. These areas were designated as urban based on population size (over 80,000 individual within a 5-10 km radius area), central location, functional services, health and educational service levels, and economic growth potential. Suburban areas were defined as townships located on the outskirts of Kuala Lumpur, Kajang, Ampang Jaya (Ulu Langat), and Batu 9 Cheras, characterized by residential areas and fewer commercial activities than urban areas [13,14]. Data collection was conducted between August and November 2018. The study protocol was approved by the Universiti Putra Malaysia Ethics Committee for Research Involving Human Subjects (JKEUPM) (JKEUPM-2018-189), and written consent was obtained from guardians prior to the study's commencement.

Assessment of demographic and health data

A self-administered questionnaire, adapted from the International Study of Asthma and Allergies in Childhood (ISAAC), was used to gather data on allergic conditions, home environmental factors and demographic characteristics. For this study, a participant was considered to have allergic rhinitis (AR) if they answered "yes" to experiencing sneezing, a runny nose,

or a blocked nose when not affected by a cold or flu over the past 12 months. Atopic dermatitis (AD) was defined by a positive response regarding a history of diagnosis and symptoms of an itchy rash lasting at least six months. Based on the provided addresses, adolescents' home locations were categorized as either urban or suburban. The completed questionnaires were cross-checked, verify responses, and clarify any uncertainties during in-person interviews conducted by the researchers.

Allergy skin test

Skin prick tests were administered to all participating adolescents after obtaining consent and assent. Five common aeroallergens, namely cat (*Felis domesticus*), house dust mite (*Dermatophagoides pteronyssinus* and *Dermatophagoides farina*), and fungi (*Cladosporium herbarium* and *Alternaria alternate*) (ALK-Abelley, Madrid, Spain) were tested on the anterior forearm. For quality control, histamine (10 mg/mL) acted as the positive control, whereas glycerol-saline functioned as the negative control. The allergic reaction was measured after 15 minutes by quantifying the mean diameter of the wheal using ruler. This procedure was conducted by trained medical assistants following the Australasian Society of Clinical Immunology and Allergy guidelines. Individuals were classified as atopic if they exhibited a positive skin prick reaction to one or more allergens tested [15].

Statistical analysis

Descriptive test analyses were performed by using the Statistical Package for the Social Sciences (SPSS) version 25.0 to examine the characteristics of study population characterised by urban and suburban areas. A comparative analysis of prevalence between urban and suburban areas for demographic characteristics, reported symptoms, and smoking status was performed using the chi-square test. The associations between home environmental factors and allergic conditions (AR and AD) were assessed using logistic regression, controlling for demographic (gender and ethnicity), smoking status, clinical variables (diagnosed asthma, atopic and parental allergy/asthma). The regression models were analysed using Stata/MP version 15.1 in which a two-sided alternative hypothesis at 5% significance level was used.

RESULTS

The personal characteristics of adolescents

Table I shows the comparison of personal characteristics between adolescents stay in urban and suburban areas. The majority of participants were Malay (87.9% overall), but the distribution differed significantly between urban and suburban locations. Suburban areas had a higher proportion of Malay participants (40.8%) compared to urban areas (59.2%). High proportion of doctor-diagnosed asthma were reported by urban adolescents (65.0%) than from suburban (35.0%), nevertheless, the

observed difference did not reach statistical significance ($p > 0.05$). Likewise, no statistically significant were observed in positive allergy prick test responses (atopy), parental allergy/asthma, smoking, or parental/sibling smoking between adolescents living in urban and suburban settings ($p > 0.05$). Allergic rhinitis (AR) was reported by 55.1% (301) of the overall study population, with similar rates observed in suburban (55.7%) and urban (54.8%) areas ($p > 0.05$). Meanwhile atopic dermatitis (AD) was reported by 14.8% (81) of the total study population, with 14.4% in suburban and 15.1% in urban areas ($p > 0.05$).

Table I: Personal characteristics of adolescents in urban and suburban areas

Characteristics	Overall (n = 546)	Suburban (n = 201)	Urban (n = 345)	p-value
Gender				
Male	217 (39.7)	66 (30.4)	151 (69.6)	0.012*
Female	329 (60.3)	135 (41.0)	194 (59.0)	
Ethnicity				
Malay	480 (87.9)	196 (40.8)	284 (59.2)	< 0.001**
Non-Malay	66 (12.1)	5 (7.6)	61 (92.4)	
Doctor-diagnosed asthma (Yes)	60 (11.0)	21 (35.0)	39 (65.0)	0.933
Atopic (Yes)	313 (57.3)	110 (35.1)	203 (64.9)	0.349
Allergic rhinitis (AR) (Yes)	301 (55.1)	112 (55.7)	189 (54.8)	0.832
Atopic dermatitis (AD) (Yes)	81 (14.8)	29 (14.4)	52 (15.1)	0.838
Parental allergy/asthma (Yes)	179 (32.8)	74 (41.3)	105 (58.7)	0.126
Smoking (Yes)	31 (5.7)	9 (29.0)	22 (71.0)	0.355
Parental/sibling smoking (Yes)	314 (57.5)	125 (39.8)	189 (60.2)	0.477

* $p < 0.05$; ** $p < 0.001$

Association analysis

Table II presents the findings of logistic regression analyses conducted separately for adolescents residing in urban and suburban areas. After controlling for key demographic, smoking status, clinical variables, we found floor renovation, water leakage and other odor last 12 months were significantly associated with AR in adolescents from suburban areas with the odd values of 1.91 (95% CI = 1.01-4.04), 1.81 (95% CI = 1.03-3.19) and 2.37 (95% CI = 1.32-4.25), respectively. In the model analysis for adolescents from urban, home environmental factor of painting, floor renovation, water leakage, mold odor last 12 months were also significantly associated with AR with the odd values of 1.26 (95% CI = 1.01-1.61), 1.70 (95% CI = 1.25-2.32), 1.22 (95% CI = 1.19-1.26), and 1.51 (95% CI = 1.06-2.15), respectively. Notably, the same model also shows that EST exposure on every day (OR = 2.32, 95% CI = 1.40-3.86) and 1-4 times/week day (OR = 1.12, 95% CI = 1.02-1.24) were

significantly associated with an increased likelihood of AR. Furthermore, adolescents from urban who keeping furry pet at home were more likely to reported AR (OR = 1.20, 95% CI = 1.17-1.24).

Table II: Association (OR) of AR and home environmental factors in adolescent from urban and suburban area

Home Environmental Factor	Suburban		Urban	
	OR	95% CI	OR	95% CI
ETS at home	1.08	0.60-1.96	1.20	0.77-1.86
Frequency ETS exposure				
Never	1	-	1	-
1-3 times/month	1.09	0.15-8.21	0.53	0.11-2.46
1-4 times/week	1.28	0.52-3.12	1.12	1.02-1.24
Everyday	1.09	0.32-3.64	2.32	1.40-3.86
Furry pet	0.88	0.54-1.43	1.20	1.17-1.24
Indoor painting last 12 months	1.33	0.48-3.66	1.26	1.01-1.61
Floor renovation last 12 months	1.91	1.01-4.04	1.70	1.25-2.32
Water leakage last 12 months	1.81	1.03-3.19	1.22	1.19-1.26
Mold odor last 12 months	2.69	0.14-5.12	1.51	1.06-2.15
Other odor last 12 months	2.37	1.32-4.25	1.14	0.42-2.78

OR = Odd ratio; 95% CI = 95% Confidence interval; R² = Coefficient of determination

The associations between home environmental factors and AD were used similar statistical analysis, logistic regression stratified by house areas. Surprisingly, all the home environmental factors tested in the models analysis were not statistically associated with the AD reported by adolescents from both areas ($p > 0.05$) (Table III).

Table III: Association (OR) of AD and home environmental factors in adolescent from urban and suburban area

Home Environmental Factor	Suburban		Urban	
	OR	95% CI	OR	95% CI
ETS at home	1.10	0.42-2.85	1.62	0.40-1.95
Frequency ETS exposure				
Never	1	-	1	-
1-3 times/month	0.65	0.06-7.15	0.82	0.30-2.26
1-4 times/week	0.67	0.21-2.10	1.06	0.18-2.06
Everyday	1.08	0.35-2.46	1.16	0.68-2.51
Furry pet	1.53	0.77-3.06	1.28	0.95-1.72
Indoor painting last 12 months	1.43	0.50-4.06	0.85	0.43-1.64
Floor renovation last 12 months	2.99	0.61-4.05	2.41	0.53-10.95
Water leakage last 12 months	0.76	0.32-1.77	1.01	0.53-1.91
Mold odor last 12 months	0.46	0.04-6.01	1.02	0.58-2.03
Other odor last 12 months	0.51	0.09-3.01	1.22	0.13-1.11

DISCUSSION

This study employed a cross-sectional study design to investigate the association of home environmental factors and allergic rhinitis and atopic dermatitis among adolescents from urban and suburban areas. Briefly, logistic regression analysis revealed that some of the home environmental factors including indoor painting, floor renovation, and water leakage last 12 months were significantly associated with AR and AD in adolescents from both urban and suburban areas. EST frequency exposure and keeping furry pets at home were also more likely associated with AR among adolescents from urban areas. These findings help clarify the prevalence and allergic reactions among adolescents, especially those living in urban and suburban settings. This information can assist in developing and increase awareness to the major publics on the environmental factors that potentially trigger and exacerbate the allergic sensitization among adolescents in urban and suburban areas.

In the present study, a higher prevalence of doctor-diagnosed asthma among adolescents living in urban areas (65%) compared to those in suburban areas (35%). Compared to other studies conducted in Northwestern Iran and China, the proportion of adolescents diagnosed with asthma were also higher among adolescents from urban than suburban areas [16,17]. Similarly, a recent study conducted in Australia strongly identified that the risk of asthma increase from inner city to the outer suburban and this trend is attributed by areal socioeconomic and distance from healthcare facilities [18]. Furthermore, a review concluded that the development of asthma in children and adolescents is influenced by several urban factors, including indoor environmental allergens and pollutants exposure, obesity, chronic stress and social determinants like poverty and housing quality [19].

Furthermore, adolescents from urban and suburban areas exhibited similar prevalence rates of AR and AD over the past 12 months and six months, respectively. This suggests that AR and AD are common conditions among adolescents across both study areas. This finding was similar to those reported by Pisithkul et al. [20] and Saeki et al. [21] among adolescents aged 12-25 years in Northern Thailand and Japan, respectively. Surprisingly, a recent systemic review reported that the prevalence of AD among adolescents in different continents ranged between 6.8% and 57.6%, which directly and indirectly impacts quality of life, behaviour, mental, family, school life and social group [22].

Findings from the logistic regression analysis in this study indicated a positive association between exposure to ETS and AR among adolescents in urban areas. This finding aligns with previous research conducted among adolescents in Korea [23]. Furthermore, recent research

suggests that tobacco smoke exposure induces significant neutrophil migration in response to inflammation, increases myeloperoxidase enzyme (MPO) activity, and leads to tissue damage through the production of hypochlorous acid (HOCl) [24]. Previous studies and review show mixed results on the relationship between exposure to furry pets and the development of allergic responses [25,26]. Many studies observed risk factors of furry pet exposure for allergic reactions, whereas other studies have shown protective effects and no associations [6,27,28].

We also observed that the reported risk of AR among adolescents from both urban and suburban areas was attributed to indoor painting, floor renovation and water leakage in the last 12 months. Additionally, we found that the presence of mold odor inside the house within the last 12 months, as reported by adolescents residing in urban areas, was associated with AR. Numerous studies have shown that visible mold and damp conditions in the home increase the risk of AR among children, adolescents, and adults [29,30]. Possible explanations for this association include water leakage creating dampness and promoting mold growth. Exposure to mold can then lead to allergic reactions and stimulate immune system activation [31,32]. Exposure to mold has the potential to cause allergic responses and initiate immune system activity.

Our finding that indoor painting is a risk factor for AR aligns with previous studies [33,34]. Indoor painting emissions such as volatile organic compounds (VOCs), heavy metals and solvents generally lead to short-term chemical exposure. It is suggested that exposure to VOCs like propylene glycol which are emitted from acrylic paints, may contribute to the development or exacerbation of AR. This occurs through the activation of the immune system, specifically by promoting a Th2-dominant immune response over Th1 function [35].

Moreover, the observed association between floor renovation and AR in adolescents from both urban and suburban areas aligns with findings reported by Ren et al. [36] in Wuhan, China. The same authors also mentioned that the chemical emission from home renovation materials including formaldehyde, phthalate esters, VOCs, and artificial synthetic materials could be the risk factors for AR, eczema and asthma. In this study, we also found that exposure to 'other odor' was a contributing risk factor for AR among adolescents in suburban areas. The results align with those reported in earlier studies from Japan and Northern Thailand [37,38]. Indoor odors, which may originate from chemicals found in cleaning products, cooking fumes, pet dander, air fresheners, and furniture, are often indicative of inadequate ventilation [39].

Nevertheless, in the present study, no home environmental factors were found to be significantly associated with

AD among adolescents in either the urban or suburban groups. In contrast to earlier studies reported that several home environmental factors including dampness, own furry pets, house renovation were positively associated with AD in children and adolescents [40]. A possible explanation for this discrepancy is that early-life indoor and outdoor environmental exposures, along with lifestyle factors from childhood to adolescence, particularly in urban environments may shape the development of innate and adaptive immune responses. This is supported by recent research demonstrating that early-life exposures, including animals, time spent outdoors, house locations and home environment factors, influence the upregulation of immune cell transcriptomes [41]. Furthermore, the clinical manifestation of allergic reactions may emerge towards the end of adolescence and typically peaks between the second and fourth decades of life [42].

This study had several limitations. First, clinical diagnoses of AR and AD were not performed. Second, recall bias may have influenced the observed associations. However, standardized questionnaires were used to collect information on AR and AD, and trained enumerators validated the responses after questionnaire completion. Finally, the use of a cross-sectional study design restricts interpretation of causal relationships.

CONCLUSION

In conclusion, our regression model demonstrated that floor renovation, water leakage and other odor within the past 12 months are significant risk factors for allergic rhinitis (AR) among adolescents residing in both urban and suburban areas. These insights are valuable for shaping targeted health interventions and treatment plans for these adolescents. Specifically, preventive measures such as avoidance of ETS exposure, implementation of adequate ventilation systems to ensure continuous air circulation and prioritizing the use of construction materials and furnishings with low emissions of organic compounds are crucial for effectively reducing pollutant exposure and enhancing indoor air quality for residents. Reducing pollutant exposure can be effectively achieved through the strategic implementation of robust ventilation systems and the careful selection of low-emitting building materials and furnishings.

ACKNOWLEDGEMENT

We gratefully acknowledge the contributions of the adolescents involved in this research and the teachers for their ongoing support and gracious hospitality. The authors wish to acknowledge Dr. Mohammed Abdulrazzaq Jabbar for his skillful technical assistance during the allergy skin prick test.

REFERENCES

1. Kogias C, Drylli A, Panagiotakos D, Douros K, Antonogeorgos G. Allergic rhinitis systematic review shows the trends in prevalence in children and adolescents in Greece since 1990. *Allergies*. 2023;3:220–8. doi:10.3390/allergies3040014.
2. Volke A, Toompere K, Laisaar K-T, Oona M, Tisler A, Johansson A, et al. 12-month prevalence of atopic dermatitis in resource-rich countries: A systematic review and meta-analysis. *Sci Rep*. 2022;12:1–13. doi:10.1038/s41598-022-19508-7.
3. Faye O, Flohr C, Kabashima K, Ma L, Paller AS, Rapelanoro FR, et al. Atopic dermatitis: A global health perspective. *J Eur Acad Dermatology Venereol*. 2024;38:801–11. doi:10.1111/jdv.19723.
4. Prasad V, Abdullah M, Nordin F, Subha ST. Prevalence, causes and treatments of allergic rhinitis in Malaysia: A literature review. *Egypt J Otolaryngol*. 2022;38:170. doi:10.1186/s43163-022-00361-4.
5. Xu X, Liu X, Li J, Deng X, Dai T, Ji Q, et al. Environmental risk factors, protective factors, and biomarkers for allergic rhinitis: a systematic umbrella review of the evidence. *Clin Rev Allergy Immunol*. 2023;65:188–205. doi:10.1007/s12016-023-08964-2.
6. Wang J, Zhang Y, Li B, Zhao Z, Huang C, Zhang X, et al. Asthma and allergic rhinitis among young parents in China in relation to outdoor air pollution, climate and home environment. *Sci Total Environ*. 2021;751:141734. doi:10.1016/j.scitotenv.2020.141734.
7. Ma X, Xie Z, Zhou Y, Shi H. Prevalence and risk factors of atopic dermatitis in Chinese children aged 1–7 years: A systematic review and meta analysis. *Front Public Heal*. 2024;12:1–8. doi:10.3389/fpubh.2024.1404721.
8. Silverberg JI, Barbarot S, Gadkari A, Simpson EL, Weidinger S, Mina-Osorio P, et al. Atopic dermatitis in the pediatric population. *Ann Allergy, Asthma Immunol*. 2021;126:417–428.e2. doi:10.1016/j.anai.2020.12.020.
9. Li S, Cao S, Duan X, Zhang Y, Gong J, Xu X, et al. Household mold exposure in association with childhood asthma and allergic rhinitis in a northwestern city and a southern city of China. *J Thorac Dis*. 2022;14:1725–37. doi:10.21037/jtd-21-1380.
10. Mohd Isa KN, Jalaludin J, Elias SM, Mohamed N, Hashim JH, Hashim Z. Evaluation of the relationship between fractional exhaled nitric oxide (FeNO) with indoor PM10, PM2.5 and NO2 in suburban and urban schools. *Int J Environ Res*. 2022;19:1–12. doi:doi:10.3390/ijerph19084580.
11. Shin J, Kim K, Kim H, Ko H, Kim B, Kim M, et al. Geographic differences in atopic dermatitis risk between urban and rural area: A systematic review and meta-analysis. *JDDG J Der Dtsch Dermatologischen Gesellschaft*. 2023;21:973–82.

doi:10.1111/ddg.15135.

12. Zhang X, Zhang M, Sui H, Li C, Huang Z, Liu B, et al. Prevalence and risk factors of allergic rhinitis among Chinese adults: A nationwide representative cross-sectional study. *World Allergy Organ J.* 2023;16:100744. doi:10.1016/j.waojou.2023.100744.
13. Aida N, Sasidhran S, Kamarudin N, Aziz N, Puan CL, Azhar B. Woody trees, green space and park size improve avian biodiversity in urban landscapes of Peninsular Malaysia. *Ecol Indic.* 2016;69:176–83. doi:10.1016/j.ecolind.2016.04.025.
14. Jabatan Perancangan Bandar & Desa. Garis panduan perancangan kemudahan masyarakat (Pindaan 2013). Kuala Lumpur: 2014.
15. ASCIA (Australasian Society of Clinical Immunology and Allergy). Skin prick testing for the diagnosis of allergic disease - A manual for practitioners. ASCIA. 2016:1–39.
16. Sabeti Z, Ansarin K, Seyedrezazadeh E, Asghari M, Jafarabadi, Zafari V, et al. A comparison of asthma prevalence in adolescents living in urban and semi-urban areas in Northwestern Iran. *Hum Ecol Risk Assess An Int J.* 2021;27:2051–2068. doi:10.1080/10807039.2021.1944051.
17. Norback D, Lu C, Wang J, Zhang Y, Li B, Zhao Z, et al. Asthma and rhinitis among Chinese children- Indoor and outdoor air pollution and indicators of socioeconomic status (SES). *Environ Int.* 2018;115:1–8. doi:10.1016/j.envint.2018.02.023.
18. Cameron E, Mo J, Yu C. A health inequality analysis of childhood asthma prevalence in urban Australia. *J Allergy Clin Immunol.* 2024;154:285–96. doi:10.1016/j.jaci.2024.01.023.
19. Grant TL, Wood RA. The influence of urban exposures and residence on childhood asthma. *Pediatr Allergy Immunol.* 2022;33:e13784. doi:10.1111/pai.13784.
20. Pisithkul T, Pisithkul T, Lao-Araya M. Impact of air pollution and allergic status on health-related quality of life among university students in northern Thailand. *Int J Environ Res Public Health.* 2024;21:452. doi:10.3390/ijerph21040452.
21. Saeki H, Ohya Y, Nawata H, Arima K, Inukai M, Rossi AB, et al. Impact of the family and household environment on pediatric atopic dermatitis in Japan. *J Clin Med.* 2023;12:2988. doi:10.3390/jcm12082988.
22. Vicho-de-la-Fuente N, Martinez-Santos A, Rodriguez-Gonzalez R, Florez B, Sheaf G, Coyne I. Impact of atopic dermatitis on adolescents and families: A mixed-method systematic review. *J Adv Nurs.* 2024;1–28. doi:10.1111/jan.16652.
23. Lee A, Lee SY, Lee K-S. Association of secondhand smoke exposure with allergic multimorbidity in Korean adolescents. *Sci Rep.* 2020;10:16409. doi:10.1038/s41598-020-73430-4.
24. Yigit E, Yuksel H, Ulman C, Yilmaz O. Nasal effects of environmental tobacco smoke exposure in children with allergic rhinitis. *Respir Med.* 2025;236:107886. doi:10.1016/j.rmed.2024.107886.
25. Onwusereaka CO, Jalaludin J, Isa KNM, Nordin SBA, Abubakar S, Choo VCP. Targeted metagenomics identification of microbiome in preschools exposed to air pollutants and their association with respiratory health symptom, allergy and eczema. *Air Qual Atmos Heal.* 2024;17:1777–1793. doi:10.1007/s11869-024-01545-y.
26. AlShatti KA, Ziyab AH. Pet-keeping in relation to asthma, rhinitis, and eczema symptoms among adolescents in Kuwait: A cross-sectional study. *Front Pediatr.* 2020;8:331. doi:10.3389/fped.2020.00331.
27. Huang S, Garshick E, Weschler LB, Hong C, Li J, Li L, et al. Home environmental and lifestyle factors associated with asthma, rhinitis and wheeze in children in Beijing, China. *Environ Pollut.* 2020;256:113426. doi:10.1016/j.envpol.2019.113426.
28. Lu C, Liu Z, Liao H, Yang W, Li Q, Liu Q. Effects of early life exposure to home environmental factors on childhood allergic rhinitis: Modifications by outdoor air pollution and temperature. *Ecotoxicol Environ Saf.* 2022;244:114076. doi:10.1016/j.ecoenv.2022.114076.
29. Suhaimi NF, Jalaludin J, Mohd Juhari MA. The impact of traffic-related air pollution on lung function status and respiratory symptoms among children in Klang Valley, Malaysia. *Int J Environ Health Res.* 2022;32:535–46. doi:10.1080/09603123.2020.1784397.
30. Lee S, Ryu S-H, Sul WJ, Kim S, Kim D, Seo S. Association of exposure to indoor molds and dampness with allergic diseases at water-damaged dwellings in Korea. *Sci Rep.* 2024;14:1–17. doi:10.1038/s41598-023-50226-w.
31. Mendell MJ, Macher JM, Kumagai K. Measured moisture in buildings and adverse health effects: A review. *Indoor Air.* 2018;28:488–99. doi:10.1111/ina.12464.
32. Yu W, Zhang Y, Li B, Zhang Y, Du C, Zhao Z, et al. Increased risk of respiratory and allergic diseases in preschoolers from building envelope dampness exposure: Repeated cross-sectional surveys in multicity China. *Build Environ.* 2023;241:110433. doi:10.1016/j.buildenv.2023.110433.
33. Bluyssen PM, Zhang D, Ortiz M. Self-reported rhinitis and headaches of students from universities in Taiwan, Chile, Suriname, China and the Netherlands, and its association with their home environment. *Intell Build Int.* 2022;14:679–89. doi:10.1080/17508975.2021.1964424.
34. Wang J, Janson C, Malinovschi A, Holm M, Franklin KA, Modig L, et al. Asthma, allergic rhinitis and atopic dermatitis in association with home environment - The RHINE study. *Sci Total Environ.* 2022;853:158609. doi:10.1016/j.scitotenv.2022.158609.
35. Barbato A, Zampieri B, Marsili G. Effects of home

- wall painting on respiratory and allergic diseases in children. *Pediatr Respir J*. 2023;1:127–36. doi:10.56164/PediatrRespirJ.2023.29.
36. Ren X, Liu Y, Hu L, Zhang Y, Xu H, Shi Y, et al. Associations between home renovation and asthma, allergic rhinitis, and eczema among preschool children in Wuhan, China. *Int J Environ Health Res*. 2022;32:2298–308. doi:10.1080/09603123.2021.1955832.
37. Prapamontol T, Norback D, Thongjan N, Suwannarin N, Somsunun K, Ponsawansong P, et al. Asthma and rhinitis in wet and dry season among students in upper Northern Thailand: the role of building dampness and household air pollution. *Int J Environ Health Res*. 2023;33:710–22. doi:10.1080/09603123.2022.2047902.
38. Takaoka M, Suzuki K, Norback D. Sick building syndrome among junior high school students in Japan in relation to the home and school environment. *Glob J Health Sci*. 2015;8:165–77. doi:10.5539/gjhs.v8n2p165.
39. Maung TZ, Bishop JE, Holt E, Turner AM, Pfrang C. Indoor air pollution and the health of vulnerable groups: a systematic review focused on particulate matter (PM), volatile organic compounds (VOCs) and their effects on children and people with pre-existing lung disease. *Int J Environ Res Public Health*. 2022;19:8752. doi:10.3390/ijerph19148752.
40. Cheng J, Wu JJ, Han G. Epidemiology and characterization of atopic dermatitis in East Asian populations: A systematic review. *Dermatol Ther (Heidelb)*. 2021;11:707–17. doi:10.1007/s13555-021-00516-w.
41. Lunjani N, Ambikan AT, Hlela C, Levin M, Mankahla A, Heldstab-Kast JI, et al. Rural and urban exposures shape early life immune development in South African children with atopic dermatitis and nonallergic children. *Allergy*. 2024;79:65–79. doi:10.1111/all.15832.
42. Nur Husna SM, Tan H-TT, Md Shukri N, Mohd Ashari NS, Wong KK. Allergic rhinitis: A clinical and pathophysiological overview. *Front Med*. 2022;9:1–10. doi:10.3389/fmed.2022.874114.