

Instant Noodles from Climate-Resilient Crops: Nutritional Quality, Sensory Acceptance, and Satiety of Sago–Bambara Groundnut Noodles Compared with Conventional Wheat Noodles

Yoanita R. Indrining Tyas^{1†}, Eny Palupi^{1*†}, Zuraidah Nasution¹, Desy R. Tarigan¹,
Alit Pangestu², Norhaizan M. Esa³

¹Department of Nutrition Science, Faculty of Medicine and Nutrition, Bogor Agricultural University, Bogor 16680, Indonesia

²Research Center for Food Technology and Processing, National Research and Innovation Agency, Yogyakarta 55861 Indonesia

³Department of Nutrition and Dietetics, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

Climate-resilient crops, such as sago (*Metroxylon sagu* Rottb.) and Bambara groundnut (*Vigna subterranea* (L.) Verdc.), offer promising solutions to enhance food security, nutritional quality, and sustainability under climate change. This study aimed to develop instant noodles based on sago starch enriched with Bambara groundnut flour and to evaluate their physicochemical properties, nutritional composition, sensory acceptability, and satiety response in comparison with conventional wheat-based instant noodles. Four different ratios of sago starch to Bambara groundnut flour were used 100:0 (F0), 70:30 (F1), 60:40 (F2), and 50:50 (F3) (w/w). The results demonstrated that sago–Bambara groundnut noodles exhibited a significantly higher total dietary fiber content (9.4–12.2 g/100 g dry matter basis, db) than sago noodles (4.5 g/100 g db) and wheat noodles (2.5 g/100 g db). However, sensory evaluation in the hedonic test showed that increasing the ratio of Bambara groundnut flour in the formula decreased noodle sensory acceptability. Therefore, formula F2 (60:40, w/w) was selected as the optimal. The quantitative descriptive sensory analysis (QDA) demonstrated that the noodles made according to this formula exhibited stronger off-beany and beany aroma, savory and starchy taste, off-beany and beany aftertaste, hardness and graininess mouthfeel compared to the sago noodles. Notably, the satiety index measurement using the visual analogue scale (VAS) questionnaire showed that the sago–Bambara groundnut noodles sustained a 50% fullness level for a significantly longer time (135 min) than wheat noodles (77 min) and demonstrated a higher satiety index (122% vs. 90%). These findings demonstrate that instant noodles formulated entirely from climate-resilient crops can enhance nutritional quality and satiety compared to the conventional wheat-based instant noodles, while remaining sensorially acceptable.

Keywords: cereal products, high-fiber food, *Metroxylon sagu*, satiety index, *Vigna subterranea*

INTRODUCTION

Obesity, as a multifactorial metabolic disorder characterized by excess fat accumulation due to an energy imbalance, has become a global nutritional problem, which according to the World Health Organization (WHO), afflicts one in eight people in the world [WHO, 2024]. One primary contributing factor to obesity is the frequent consumption of ultra-processed food, such as instant noodles

[Septiani & Sugiatmi, 2026]. Instant noodles are processed foods that contain high calories, saturated fats, and sodium but are generally low in essential nutrients, such as fiber, vitamins, and minerals and are often associated with a high glycemic response and low satiety response [Huh *et al.*, 2017; Salihi *et al.*, 2025]. Conventional instant noodles are predominantly produced using refined wheat flour. The increasing reliance on wheat, particularly in regions

*Corresponding Author:

E-mail: enypalupi@apps.ipb.ac.id (Dr. E. Palupi)

†These authors had equal contribution to this work

Submitted: 22 January 2026

Accepted: 28 April 2026

Published on-line: 25 May 2026



© Copyright: © 2026 Author(s). Published by InLife Institute of Animal Reproduction and Food Research, Polish Academy of Sciences. This is an open access article licensed under the Creative Commons Attribution 4.0 License (CC BY 4.0) (<https://creativecommons.org/licenses/by/4.0/>)

where it is not locally produced, raises concerns regarding food system vulnerability [Farhan *et al.*, 2025]. This vulnerability is exacerbated as climate change poses additional challenges to global food security [Begna & Wakweya, 2026], emphasizing the urgent need to diversify staple food sources with climate-resilient crops [Benitez-Alfonso *et al.*, 2023].

Sago (*Metroxylon sago* Rottb.) and Bambara groundnut (*Vigna subterranea* (L.) Verdc.) are underutilized crops that exhibit strong adaptability to marginal environments, drought tolerance, and stable yields under adverse climatic conditions [Bintoro *et al.*, 2018; Tan *et al.*, 2020]. Sago is a traditional source of starch in some tropical regions that contains high levels of carbohydrates, approximately 84.7 g per 100 g of starch [Moshawih *et al.*, 2025], but has a low protein content (0.9 g/100 g starch) [Ministry of Health, 2018]. Consequently, food products developed primarily from sago tend to have a limited protein content; for instance, previous studies have reported that sago-based noodles contain only 3.07 g protein/100 g [Dewita *et al.*, 2019]. In contrast, Bambara groundnut is a legume rich in protein (17–25 g/100 g), with a relatively balanced amino acid profile [Maphosa *et al.*, 2022]. Therefore, the partial substitution of sago starch with Bambara groundnut flour is expected to significantly improve the nutritional quality of sago-based products, particularly by increasing their protein content. The complementary nutritional characteristics of these two climate-resilient crops suggest their potential for the development of nutritionally-enhanced staple foods.

Previous research has demonstrated the effectiveness of legume substitutions in noodle formulations; for instance, substituting wheat flour with 25% chickpea flour increased the protein content of noodles from 12.79 to 15.31 g/100 g [Bayomy & Alamri, 2022]. However, research on the integration of fully climate-resilient crops into instant noodle formulations remains limited. In particular, the combined use of sago starch and Bambara groundnut flour in instant noodles has not been extensively investigated, especially regarding the enhancement of protein and dietary fiber levels and satiety-related outcomes. Given the growing interest in functional convenience foods that support appetite regulation and metabolic health, the evaluation of satiety responses alongside nutritional composition is becoming increasingly relevant.

Therefore, this study aimed to develop instant noodles based on sago starch enriched with Bambara groundnut flour and compare their nutritional quality and satiety properties with those of conventional wheat-based instant noodles. By linking climate resilience, nutritional improvement, and consumer-relevant functional outcomes, this study contributes to the development of sustainable and healthier instant noodle alternatives.

To achieve these objectives, this study was conducted in several stages, including (1) formulation of sago noodles with Bambara groundnut flour, (2) assessment of their physical characteristics and nutritional content, (3) their sensory evaluation, and (4) their satiety index measurement.

MATERIALS AND METHODS

■ Materials

There were two main materials that have been used in this research, *i.e.*, sago (*M. sago*) starch and Bambara groundnut (*V. subterranea*) beans. The Bambara groundnut was purchased from the traditional market in Bogor, Indonesia. As many as 40 kg of beans have been used in a single run of Bambara groundnut flour. The sago starch (20 kg) was procured from the local farmer in Banggai, Central Sulawesi, Indonesia. Commercial all-salt (Refina, UnichemCandi Indonesia, Sidoarjo, Indonesia), sodium tripolyphosphate (Sodium tripolyphosphate, Aditya Birla Chemicals, Samut Prakan, Thailand), and wheat-based instant noodles (Indomie, Indofood, Jakarta, Indonesia) were purchased at a local market in Bogor, Indonesia.

■ Production of sago and Bambara groundnut noodles

The production of the sago and Bambara groundnut noodles consisted of two main stages, *i.e.*, sago flour and Bambara groundnut flour preparation, and noodle preparation. To produce sago flour, sago starch was sun-dried for 18 h. Dried sago starch was ground and sieved using an 80-mesh screen to obtain sago flour. To produce Bambara groundnut flour, the beans were soaked for 12 h. Then, they were boiled at 90°C for 10 min to inactivate lipoxygenase and eliminate the beany flavor [Chong *et al.*, 2019]. The beans were dried using a cabinet dryer (CTM SRL Manufacture, Milan, Italy) at 90°C at 10 rpm. Dried beans were milled using a pin disc mill and sieved using an 80-mesh screen to obtain Bambara groundnut flour.

The development of the sago and Bambara groundnut noodle formulations was conducted as a trial-and-error process based on the Agustia *et al.* [2016] recipe before producing the final noodle product. Formulas of noodles with different flour composition used in this study have been adjusted to the Indonesian National Standard to meet the requirements of instant noodle quality standards [BSN, 2018] and the high-fiber claims for processed food products [Indonesian Food and Drug Authority, 2022]. The formulas contained different ratios of sago flour (SF) and Bambara groundnut flour (BGF): F0 (100% SF), F1 (70% SF:30% BGF, w/w), F2 (60% SF:40% BGF, w/w), and F3 (50% SF:50% BGF, w/w) (Table 1).

Sago flour was mixed into 100 mL of water and cooked at 90°C for 1 min until it thickened to produce a gelatinized sago starch. Then, it was mixed with Bambara groundnut flour and other ingredients, including sodium tripolyphosphate (0.5% of recipe ingredients, w/w) and salt (0.5% of recipe ingredients, w/w) using a mixer (Miyako HM-620, Jakarta, Indonesia) until a smooth dough was formed. A dough made from wheat flour, instead of SF and BGF, was also prepared. The noodles were formed using a roll press (Ardin CM2020, Jakarta, Indonesia) with a length of ± 15 cm and dried at 105°C for 60 min in an oven (Kirin KBO 190RA, Jakarta, Indonesia). The dried noodles were boiled at 100°C for 7–9 min and allowed to cool to room temperature prior to analysis.

Table 1. Formulas of noodles with different ratios of sago flour (SF) and Bambara groundnut flour (BGF).

Ingredient	Unit	F0 (100:0)	F1 (70:30)	F2 (60:40)	F3 (50:50)
Sago flour	g (%)	100 (49.5)	70 (34.7)	60 (29.7)	50 (24.8)
Bambara groundnut flour	g (%)	0 (0)	30 (14.9)	40 (19.8)	50 (24.8)
Salt (NaCl)	g (%)	1 (0.5)	1 (0.5)	1 (0.5)	1 (0.5)
Sodium tripolyphosphate	g (%)	1 (0.5)	1 (0.5)	1 (0.5)	1 (0.5)
Water	mL	100 (49.5)	100 (49.5)	100 (49.5)	100 (49.5)

F0, formula of noodles made from SF; F1, formula of noodles made from SF and BGF in a 70:30 (w/w) ratio; F2, formula of noodles made from SF and BGF in a 60:40 (w/w) ratio; and F3, formula of noodles made from SF and BGF in a 50:50 (w/w) ratio.

■ Analysis of physical characteristics

The physical characteristics of the noodles were assessed by evaluating their color and texture. The color was measured using a chromameter (AMT511, Amtast, Lakeland, FL, USA), calibrated with a provided set of black and white plates. The CIELab color space was defined by L^* , a^* , and b^* values, which represent lightness (dark–bright), redness (green–red), and yellowness (blue–yellow), respectively.

The firmness was measured using a texture analyzer (TA.XT Plus, Stable Micro Systems, Godalming, UK). Three strands of cooked noodles were placed parallel to each other on the platform and positioned perpendicular to the Asian Noodle Rig (HDP/ANR) probe. Compression was initiated at a pre-test speed of 1 mm/s, a test speed of 2 mm/s, and a post-test speed of 10 mm/s with a compression distance of 9.9 mm and a strain level of 10%. The test was activated by a trigger force of 0.049 N and trigger distance of 2 mm until the sample was compressed with a force of 0.981 N and break sensitivity of 0.098 N. The firmness value was obtained from the maximum force required to compress the noodle strands, expressed in Newton (N).

■ Nutrient content determination

The nutrient content determination of sago and Bambara groundnut noodles included determining the moisture content using the gravimetric method (AOAC International method no. 2005:925.10), ash content using the gravimetric method (AOAC 2005:923.03), protein content using the Kjeldahl method (AOAC 2005:955.04), lipid content using the Soxhlet method (AOAC 2005:920.39c), carbohydrate content using the by-difference method, and dietary fiber content (total, soluble and insoluble) using the enzymatic gravimetric method [AOAC, 2005]. The available carbohydrate content was calculated as the difference between the carbohydrate content and the dietary fiber content. The results of nutrient content were expressed on a dry matter basis (db) of noodles.

■ Sensory evaluation

The research protocol of sensory evaluation has been reviewed and approved by the Human Research Ethics Committee of Bogor Agricultural University in Bogor, Indonesia, with the approval

number 1633/IT3.KEPMSM IPB/SK/2025. The sensory evaluation consisted of two tests: a hedonic acceptance test and a quantitative descriptive analysis.

The sensory attributes assessed in the acceptance test included appearance, aroma, taste, mouthfeel, aftertaste, and overall acceptability of noodles. Sensory attributes were assessed using a 9-point hedonic scale ranging from 1 (dislike extremely) to 9 (like extremely), based on the level of preference for each sensory attribute assessed. Thirty semi-trained panelists (23 women and 7 men) aged 18–40 years participated in the study. The inclusion criteria for panelists in the sensory evaluation were as follows: willingness to participate in the sensory evaluation; consistency in decision-making; absence of ear, nose, and throat disorders; normal color vision; no allergies to ingredients contained in the noodle formulations; required to fast for 1 h prior to the sensory evaluation; avoid consuming pungent foods before sensory evaluation; and prohibited using strongly scented cosmetics such as perfumes during the sensory evaluation [ISO 8586:2014].

The quantitative descriptive sensory analysis (QDA) was carried out only on the sago flour noodles (F0; as control) and optimal noodle formula made from SF and BGF in ratio 40:60, w/w (F2). The selection of noodle formula for QDA and satiety index measurement was made by considering the most relevant results from the nutritional composition analysis and hedonic sensory evaluation.

The QDA involved 11 trained panelists who were selected using the Ishihara test, matching test, triangle test, and ranking test. The QDA procedure was carried out in two main stages: a focus group discussion (FGD) to identify sensory attributes, followed by an assessment of the intensity of each attribute. The sensory attributes assessed included aroma, taste, mouthfeel, and aftertaste. The sensory attributes were assessed using an intensity scale ranging from 0 (very weak) to 10 (very strong) [ISO 8586:2014].

■ Satiety index measurement

The research protocol of satiety index measurement has been reviewed and approved by the Human Research Ethics Committee of Bogor Agricultural University in Bogor, Indonesia, with the approval number 1633/IT3.KEPMSM IPB/SK/2025.

Table 2. Subject characteristics for satiety index assessment.

Subject code	Sex	Age (years)	BMI (kg/m ²)	FBS (mg/dL)
01	Female	25	21.3	85
02	Female	22	21.4	89
03	Female	24	21.9	79
04	Female	23	23.3	88
05	Female	24	19.1	90
06	Female	23	19	91
07	Female	21	21.6	96
08	Female	20	20.9	88
09	Female	24	22.9	94
10	Female	19	22.3	94
11	Male	24	21.7	97
12	Male	24	20.9	99
13	Male	22	21.7	92
14	Male	21	22.1	79
15	Male	21	22.8	83
16	Male	25	22.5	86
17	Male	24	21	93
Mean		22.71	21.55	89.59
SD		1.79	1.18	5.86
%CV		7.8	5.4	6.5

BMI, body mass index; FBS, fasting blood sugar; SD, standard deviation; %CV, coefficient of variation.

Seventeen subjects participated in this study, including ten women (58.8%) and seven men (41.2%) with an average age of 22.91 ± 1.30 years (Table 2), who were selected using a purposive sampling method. The inclusion criteria for the subjects of this study were as follows: (1) man or woman aged between 20 and 25 years, (2) a normal body mass index (BMI) ($18.5\text{--}22.9\text{ kg/m}^2$), (3) a normal fasting blood sugar level ($<100\text{ mg/dL}$), (4) no allergies or food intolerance, (5) engaged in light to moderate physical activity, and (6) normal dietary behavior. The exclusion criteria for the subjects of this study were (1) pregnancy or breastfeeding (for female subjects), (2) active smokers, (3) consuming alcohol, (4) experiencing digestive disorders and/or eating disorders, and (5) undergoing medications that can affect eating habits. One subject was excluded due to human error in completing the questionnaire. The satiety scores of the excluded subject exhibited abnormal fluctuations that deviated significantly from those of the other subjects, resulting

in outlier data for the satiety index measurement. A sample size of 17 was considered acceptable for a satiety index study [Forde, 2018]. The homogeneity of subject characteristics was determined using the coefficient of variation percentage (%CV). If the $\%CV \leq 5\%$ for body mass index and $\%CV < 33\%$ for fasting blood sugar, then the subject characteristics were considered homogenous [Jamaiyah *et al.*, 2010; Mo *et al.*, 2021]. The %CV was calculated using Equation (1) [Ulijaszek & Kerr, 1999]:

$$\%CV = \frac{\text{Standard deviation}}{\text{Mean}} \times 100\% \quad (1)$$

The satiety index measurement was conducted on four different days with a 6-day washout period between the tested foods to ensure that the physiological condition had returned to normal [Nolan *et al.*, 2016]. Subjects were required to fast for 8–10 h before the test. After fasting, they consumed isocaloric food (240 kcal) and one cup of water (220 mL) within a maximum of 15 min. The isocaloric foods given to the subjects consisted of standard and test foods. White bread was used as a standard food, according to Holt *et al.* [1995]. The tested foods were F0 (100% SF), F2 (60% SF:40% BGF, w/w), and wheat noodles. According to their energy content, the portion of F0 was 55.9 g, that of F2 was 55.1 g, that of wheat noodles was 37.4 g, and that of white bread was 100.0 g. The tested foods were served with 250 mL of broth and seasonings. Then, the four parameters of the satiety index: (1) hunger score, (2) fullness score, (3) desire to eat score, and (4) food quantity score, were measured using the 100-mm visual analogue scale (VAS) questionnaire at 0, 30, 60, 90, 120, 150, and 180 min after consumption of the tested foods.

The scores of hunger, fullness, desire to eat, and food quantity were plotted on a graph to calculate the area under the curve (AUC). The curve was scaled in units of sampling time (counted in hours) and fullness index, which was expressed as a ruler scale in mm. The AUC was calculated using the trapezoid method and expressed as mmxh. The satiety index was calculated by comparing the AUC of the tested food with that of the standard food for each subject. Then, the satiety index was calculated using Equation (2) based on Holt *et al.* [1995]:

$$\text{Satiety index} = \frac{\text{AUC of tested food}}{\text{AUC of standard food}} \times 100\% \quad (2)$$

Statistical analysis

The data for each parameter were presented as mean and standard deviation of triplicated measurements/determinations. Statistical analysis was performed using Excel 2016 for Windows (Microsoft, Redmond, WA, USA) and IBM SPSS Statistics (version 26.0; IBM Corp., Armonk, NY, USA). The results of the determination of the physical characteristics, nutritional composition, hedonic sensory evaluation, and satiety index were analyzed using the one-way analysis of variance (ANOVA), followed by the Duncan's multiple range test. However, the results of the quantitative descriptive analysis were analyzed using the independent t-test. The differences were considered significant at $p < 0.05$.

RESULTS AND DISCUSSION

■ Physical characteristics

The firmness values of noodles made from sago and Bambara groundnut flours are shown in **Table 3**. The incorporation of Bambara groundnut flour into the recipe significantly ($p < 0.05$) decreased the firmness of the noodles compared to those made from sago flour. However, the firmness exhibited no significant difference ($p \geq 0.05$) among the F1, F2, and F3 samples. Similar results were reported by Singthong *et al.* [2026], according to whom the noodles substituted with 50% kidney bean flour had significantly decreased firmness compared to the rice and tapioca flour noodles. The incorporation of legume flour can disrupt the starch structure, which causes firmness reduction [Singthong *et al.*, 2026]. The protein and fiber in legume flour form physical barriers around starch granules and aggressively compete for available water. These macromolecules restrict the swelling required for starch gelatinization, disrupting the formation of the starch gel network. This process may weaken the structural integrity of the noodle matrix, resulting in softer noodles [Singthong *et al.*, 2026].

The color coordinate values of the noodles made from sago and Bambara groundnut flours are shown in **Table 3**. Additionally, the appearance of cooked noodles is presented in **Figure 1**. The substitution of the Bambara groundnut flour significantly ($p < 0.05$) decreased the lightness (L^*), redness (a^*), and yellowness (b^*) values of noodle color compared to those made from

sago flour (**Table 3**). Similar findings were reported by Mashau *et al.* [2022], who showed that the crust and crumb of steamed breads fortified with Bambara groundnut flour had significantly lower L^* and a^* values compared to the steamed wheat bread, although an increase in b^* value was noted, in contrast to the noodles in our study. The darker color of the noodles might be due to the presence of color flavonoids and tannins in the Bambara groundnut seeds, especially in the seed coat [Palamae *et al.*, 2024; Pretorius *et al.*, 2023].

■ Nutritional composition

The nutritional composition of the sago and Bambara groundnut flours as well as noodles made from these flours is presented in **Table 4**. Data on wheat noodles is also shown in **Table 4**. The energy value and nutrient content of the noodles made from sago and Bambara groundnut flour were significantly different ($p < 0.05$) from those of wheat noodles. The contents of carbohydrates, available carbohydrates, and total dietary fiber of the noodles made from sago and Bambara groundnut flour were significantly higher ($p < 0.05$) compared to wheat noodles. However, the energy value and contents of protein and lipids of the noodles made from sago and Bambara groundnut flour were significantly lower ($p < 0.05$) than those of the wheat noodles. The use of Bambara groundnut flour in the noodle formula significantly ($p < 0.05$) increased the energy value and contents of protein, lipids, total dietary fiber, soluble dietary fiber, insoluble

Table 3. Physical characteristics of noodles made from sago flour (SF) and Bambara groundnut flour (BGF) used in different ratios.

Physical characteristic	F0 (100:0)	F1 (70:30)	F2 (60:40)	F3 (50:50)
L^*	80.8±0.7 ^a	77.3±0.5 ^b	76.2±0.7 ^c	74.9±0.4 ^d
a^*	10.0±0.4 ^a	6.3±0.2 ^b	6.0±0.3 ^{bc}	5.7±0.2 ^c
b^*	15.0±1.1 ^a	9.1±0.3 ^b	8.6±0.6 ^b	7.9±0.5 ^b
Firmness (N)	53.0±1.2 ^a	33.9±4.7 ^b	28.5±1.2 ^b	27.4±0.0 ^b

Results are shown mean ± standard deviation. Different superscript letters in each row indicate significant differences at $p < 0.05$, analyzed using Duncan's multiple range test. F0, noodles made from SF; F1, noodles made from SF and BGF in a 70:30 (w/w) ratio; F2, noodles made from SF and BGF in a 60:40 (w/w) ratio; and F3, noodles made from SF and BGF in a 50:50 (w/w) ratio; L^* , lightness; a^* , redness; b^* , yellowness.



Figure 1. Appearance of noodles made from sago flour (SF) and Bambara groundnut flour (BGF) used in different ratios. F0, noodles made from SF; F1, noodles made from SF and BGF in a 70:30 (w/w) ratio; F2, noodles made from SF and BGF in a 60:40 (w/w) ratio; and F3, noodles made from SF and BGF in a 50:50 (w/w) ratio.

Table 4. Nutritional composition of wheat noodles and noodles made from sago flour (SF) and Bambara groundnut flour (BGF) used in different ratios.

Nutritional composition	SF	BGF	F0 (100:0)	F1 (70:30)	F2 (60:40)	F3 (50:50)	Wheat noodles
Energy (g/100 g db)	401±0.0 ^f	424±0.3 ^b	393±0.3 ^g	403±0.3 ^e	406±0.2 ^d	409±0.7 ^c	481±0.1 ^a
Protein (g/100 g db)	0.4±0.0 ^g	18.7±0.0 ^a	2.1±0.1 ^f	6.5±0.0 ^e	8.6±0.0 ^d	10.6±0.2 ^c	12.0±0.1 ^b
Lipid (g/100 g db)	0.5±0.0 ^f	7.6±0.0 ^b	0.0±0.0 ^g	2.9±0.0 ^e	3.6±0.1 ^d	4.5±0.1 ^c	17.3±0.0 ^a
Carbohydrate (g/100 g db)	98.6±0.0 ^a	70.2±0.0 ^f	96.0±0.2 ^b	87.9±0.1 ^c	84.7±0.2 ^d	81.8±0.3 ^e	69.4±0.1 ^g
Available carbohydrate (g/100 g db)	–	–	91.7±0.3 ^a	78.5±0.1 ^b	73.6±0.1 ^c	69.7±0.1 ^d	66.9±0.1 ^e
Total dietary fiber (g/100 g db)	–	–	4.5±0.2 ^d	9.4±0.1 ^c	11.1±0.3 ^b	12.2±0.1 ^a	2.5±0.0 ^e
Soluble dietary fiber (g/100 g db)	–	–	1.1±0.2 ^d	2.1±0.0 ^c	2.5±0.1 ^b	2.8±0.1 ^a	–
Insoluble dietary fiber (g/100 g db)	–	–	3.4±0.0 ^d	7.3±0.1 ^c	8.5±0.4 ^b	9.4±0.2 ^a	–
Moisture (g/100 g wb)	14.5±0.1 ^a	9.7±0.0 ^b	5.5±0.3 ^e	7.3±0.3 ^c	6.7±0.3 ^d	4.6±0.2 ^f	6.6±0.0 ^d
Ash (g/100 g db)	0.5±0.0 ^f	3.5±0.0 ^a	1.9±0.1 ^d	2.8±0.8 ^c	3.1±0.7 ^b	3.2±0.2 ^b	1.3±0.0 ^e

Results are shown mean ± standard deviation. Different superscript letters in each row indicate significant differences at $p < 0.05$, analyzed using Duncan's multiple range test. F0, noodles made from SF; F1, noodles made from SF and BGF in a 70:30 (w/w) ratio; F2, noodles made from SF and BGF in a 60:40 (w/w) ratio; and F3, noodles made from SF and BGF in a 50:50 (w/w) ratio; db, dry basis; wb, wet basis.

dietary fiber, and ash of the noodles compared to the control sample made from sago flour. Meanwhile, the substitution of sago starch with Bambara groundnut flour significantly ($p < 0.05$) decreased the carbohydrate and available carbohydrate contents of the noodles compared to the control sample made from sago flour. Similar results were reported by Makhuvha *et al.* [2024], who found that substituting wheat flour with 25% Bambara groundnut flour in pasta significantly increased the protein, lipid, fiber, and ash contents, while decreasing the carbohydrate content compared to wheat flour pasta. Based on the Indonesian Food and Drug Authority regulation [2022] regarding the supervision of food labeling and advertisements, the developed noodle formulas demonstrate significant potential for nutritional claims. The F0 noodle formula meets the claim criteria for a 'source of fiber' food product (≥ 3 g/100 g), while F1, F2, and F3 formulas for a 'high-fiber' food product (≥ 6 g/100 g). High-fiber foods are associated with higher satiety index scores, leading to a more sustained feeling of fullness [Gerstein *et al.*, 2004]. Consequently, enhancing satiety through fiber intake serves as a strategic approach to mitigating the risk of excess energy intake and subsequent weight gain [Hervik & Svihus, 2019; Savastano *et al.*, 2014].

■ Sensory characteristics

The results of the sensory characteristics assessment in the hedonic acceptance test of the noodles made from sago flour and Bambara groundnut flour are presented in Table 5. The scoring of the hedonic attributes of F1 and F2 did not differ significantly ($p \geq 0.05$). However, the higher proportion of Bambara groundnut flour in the noodle formula resulted in the taste, mouthfeel, aftertaste, and overall acceptance of F3 being

Table 5. Sensory characteristics in the hedonic acceptance test of noodles made from sago flour (SF) and Bambara groundnut flour (BGF) used in different ratios.

Sensory attribute	F1 (70:30)	F2 (60:40)	F3 (50:50)
Appearance	5.9±1.7 ^a	6.1±1.4 ^a	5.2±1.5 ^a
Aroma	6.0±1.1 ^a	6.3±1.1 ^a	5.7±1.3 ^a
Taste	5.6±1.7 ^{ab}	6.0±1.3 ^a	5.1±1.4 ^b
Mouthfeel	5.1±1.8 ^{ab}	5.6±1.4 ^a	4.5±1.4 ^b
Aftertaste	5.4±1.4 ^{ab}	5.8±1.1 ^a	5.0±1.3 ^b
Overall acceptance	5.9±1.4 ^{ab}	6.2±1.2 ^a	5.2±1.5 ^b

Results are shown mean ± standard deviation. Different superscript letters in each row indicate significant differences at $p < 0.05$, analyzed using Duncan's multiple range test. F1, noodles made from SF and BGF in a 70:30 (w/w) ratio; F2, noodles made from SF and BGF in a 60:40 (w/w) ratio; and F3, noodles made from SF and BGF in a 50:50 (w/w) ratio.

significantly ($p < 0.05$) lower rated by panelists than F2. These noodles did not differ significantly ($p \geq 0.05$) only in the appearance and aroma. The overall result of the sensory hedonic acceptance test indicates that the noodles made from sago flour and Bambara groundnut flour received hedonic scores between 5 and 6, indicating they fell within the 'like moderately' category and were generally accepted by the panelists.

Since noodles F1 and F2 received higher scores in the hedonic acceptance test than F3 and, at the same time, F2 had a more favorable nutrient composition than F1 (especially a higher dietary fiber content), the F2 noodles were selected for QDA and satiety index measurement.

Table 6. Sensory attributes and intensity standards used in the quantitative descriptive analysis of noodles.

Attribute		Definition	Standards	Intensity
Aroma	Off-beany	The aroma associated with uncooked or raw beans	Bambara groundnut flour	10
	Onion	The aroma associated with fried onions	Onion oil	10
	Beany	The aroma associated with cooked or boiled beans	Boiled Bambara groundnut	10
Taste	Salty	The salt taste on the tongue associated with sodium chloride	0.2% NaCl solution	2
			0.8% NaCl solution	8
	Savory	The savory taste on the tongue associated with monosodium glutamate	0.18% MSG solution	2
			0.72% MSG solution	8
	Starchy	The starch taste on the tongue associated with sago flour	Sago flour	10
Aftertaste	Onion	The fried onion taste on the tongue	Onion oil	10
	Off-beany	The intensity of taste lingering in the mouth for 15 s after swallowing associated with uncooked or raw beans	Bambara groundnut flour	10
	Beany	The intensity of taste lingering in the mouth for 15 s after swallowing associated with cooked or boiled beans	Boiled Bambara groundnut	10
	Starchy	The intensity of taste lingering in the mouth for 15 s after swallowing associated with starch or flour	Sago flour	10
Mouthfeel	Springiness	The rate of return of the sample to its original shape after chewing	Wheat noodle	10
	Hardness	The force required to bite the sample completely with the molars	Wheat noodle	10
	Graininess	Degree to which grains are perceived in the mouth	Bambara groundnut flour	10

Thirteen sensory attributes were selected by the panel in QDA of the sago flour noodles and noodles made from sago flour and Bambara groundnut flour. These sensory attributes and intensity standards used are listed in **Table 6**. Scores of eight attributes exhibited significant differences ($p < 0.05$) between F0 and F2, comprising two aroma, two taste, two aftertaste, and two mouthfeel attributes (**Figure 2**). The substitution of sago starch with 40% (w/w) Bambara groundnut flour resulted in a stronger off-beany aroma, beany aroma, savory taste, starchy taste, off-beany aftertaste, beany aftertaste, hardness mouthfeel, and graininess mouthfeel of the noodles compared to the sago noodles. The stronger beany aroma of noodles made from both flours compared to the sample without Bambara groundnut flour may be due to activity of lipoxygenase of Bambara groundnut [Kudre & Benjakul, 2013]. Lipoxygenase catalyzes the oxidation of polyunsaturated fatty acids, such as linoleic acid, resulting in aldehyde compounds that produce a beany aroma [Kudre & Benjakul, 2013]. Although blanching can inactivate the lipoxygenase [Chong *et al.*, 2019], the decrease in enzyme activity depends on the temperature and time [Estiasih *et al.* 2025]. According to Nguyen *et al.* [2025], lipoxygenase activity remains at 32–63% after heating at 80–90°C for 25 min. Consequently, the sub-optimal heating temperatures and times lead to the incomplete inactivation of the lipoxygenase enzyme. Furthermore, the substitution of Bambara groundnut flour significantly enhances the savory taste of the noodles, due to its high glutamic acid content

(18.31 g/100 g) [Aremu *et al.*, 2025]. This amino acid is present naturally in protein-rich foods that produce a savory taste [Jahanshiri *et al.*, 2022]. Moreover, it also enhances the graininess of the noodles, because the insoluble fiber content exceeding 2% is known to induce a gritty texture in the mouth [Aussan-asuwannakul *et al.*, 2022; Chakraborty *et al.*, 2019], which is further exacerbated by insufficiently fine flour sieving during the processing [lju *et al.*, 2025].

■ Satiety index

The satiety index is a measure that evaluates the level of satiety produced by a test food compared to that produced by a standard food, and is expressed as a percentage [Forde, 2018]. The nutrient content of the tested noodles and wheat bread used as a standard food *per* 240 kcal, satiety scores, and satiety indices are shown in **Table 7**. A significant difference ($p < 0.05$) was found in the satiety index between sago noodles (F0), noodles made from sago and Bambara groundnut flours (F2) and wheat noodles. The satiety index values of the tested foods, in the order of wheat noodles, sago noodles as control sample, and sago noodles with Bambara groundnut flour, were 90%, 113%, and 122%, respectively. The satiety index of the noodles made from sago and Bambara groundnut flours was significantly higher than that of the wheat noodles, indicating that the substitution of sago starch with 40% (w/w) Bambara groundnut flour resulted in a greater sense of fullness. As shown in **Table 7**, the sago noodles with and without Bambara groundnut flour




Figure 2. Results of quantitative descriptive sensory analysis (QDA) of noodles made from sago flour (F0) and sago and Bambara groundnut flours used in a 60:40 (w/w) ratio (F2). Asterisks indicate significant differences at $p < 0.05$, analyzed using independent t -test.

Table 7. Nutrient composition of a portion (240 kcal) of white bread, wheat noodles, and noodles made from sago flour (F0), and from sago and Bambara groundnut flours in a ratio of 60:40, w/w (F2), as well as satiety scores and satiety index for these food products.

Parameter	White bread	F0 (100:0)	F2 (60:40)	Wheat noodles
Nutrient content				
Protein (g)	9.9±0.0 ^a	1.1±0.1 ^d	4.4±0.0 ^c	5.3±0.1 ^b
Lipid (g)	4.7±0.0 ^b	0.0±0.0 ^d	1.9±0.1 ^c	7.7±0.0 ^a
Carbohydrate (g)	48.4±3.6 ^a	48.0±0.2 ^b	40.5±0.2 ^c	30.8±0.1 ^d
Available carbohydrate (g)	45.4±0.1 ^a	45.8±0.4 ^a	35.2±0.1 ^b	29.7±0.1 ^c
Total dietary fiber (g)	3.0±0.1 ^b	2.4±0.1 ^c	5.7±0.2 ^a	1.1±0.0 ^d
Soluble dietary fiber (g)	0.7±0.0 ^b	0.6±0.1 ^c	1.3±0.0 ^a	–
Insoluble dietary fiber (g)	2.3±0.1 ^b	1.8±0.0 ^c	4.4±0.2 ^a	–
Moisture (mL)	35.4±0.4 ^a	34.6±0.0 ^b	36±0.0 ^a	25.4±0.0 ^c
Weight per 240 kcal (g)	100	55.9	55.1	37.4
Satiety score				
Hunger (mmxh)	44.6±6.9 ^b	38.6±7.9 ^c	36.4±8.9 ^c	51.4±17.9 ^a
Fullness (mmxh)	54.8±7.1 ^b	61.6±8.2 ^a	65.8±8.2 ^a	46.6±17.3 ^b
Desire to eat (mmxh)	45.9±5.9 ^b	38.5±8.7 ^c	36.9±8.0 ^c	55.5±16.7 ^a
Food quantity (mmxh)	45.5±7.4 ^b	37.9±8.9 ^c	37.3±11.0 ^c	54.2±15.2 ^a
Satiety index (%)	Reference	113±15 ^b	122±17 ^a	90±31 ^c

Results are shown mean ± standard deviation. Different superscript letters in each row indicate significant differences at $p < 0.05$, analyzed using Duncan's multiple range test.

significantly ($p < 0.05$) reduced hunger, increased fullness, decreased desire to eat, and decreased food quantity compared to wheat noodles.

Figure 3 shows that the curve of the fullness index of the sago noodles with Bambara groundnut flour exceeds the standard white bread food curve, indicating that it can provide a longer feeling of fullness than white bread. The regression equations showed that sago noodles with Bambara groundnut flour ($y = -0.344x + 96.354$) reached a 50% level of fullness at 135 min. In comparison, white bread ($y = -0.384x + 89.361$) and wheat noodles ($y = -0.296x + 72.911$) reached the same level of fullness significantly faster, i.e., at 102 min and 77 min, respectively. Therefore, it can be concluded that the consumption of noodles made from sago and Bambara groundnut flours in a 60:40 (w/w) ratio can prolong the feeling of fullness for up to 45 min longer than white bread, and even further up to 58 min longer than wheat noodles. These findings are consistent with observations made by Ridwan *et al.* [2024], who reported that tuber-bread with 60% Bambara groundnut flour could prolong the feeling of fullness up to 95 min longer than white bread. The increased feeling of fullness was due to the fact that F2 portion served had the highest content of total fiber (5.7 g) compared to F0 portion (2.4 g) and wheat noodle portion (1.1 g) (**Table 7**). According to Palupi *et al.* [2024], the consumption of foods high in fiber can

increase the feeling of fullness or satiety. Dietary fiber, particularly soluble fiber, plays a significant role in increasing satiety through several physiological mechanisms throughout the digestive tract [Hervik & Svihus, 2019]. Due to its water solubility and gel-forming ability, soluble dietary fiber may increase the viscosity of gastrointestinal digesta. Higher viscosity can slow down gastrointestinal transit, delay gastric emptying, and increase gastric distension [Ariyaratna *et al.*, 2025]. Increasing the gastric distension can inhibit the secretion of the ghrelin hormone, which sends signals to the hypothalamus to suppress hunger and stimulate satiety [Hervik & Svihus, 2019]. In addition, dietary fiber can increase oral processing time and effort needed for mastication. A longer oral processing time seems to stimulate cephalic phase responses, and these are deemed to contribute to satiety [Hervik & Svihus, 2019]. Thus, the sago with Bambara groundnut noodles can be used as an alternative high-fiber staple food that can increase the feeling of fullness and reduce the consumption of additional foods, which often contribute to excess calorie intake.

CONCLUSIONS

Instant noodles developed from sago starch enriched with Bambara groundnut flour exhibited superior nutritional quality and satiety effects compared with conventional wheat-based instant noodles. Their higher protein and dietary fiber contents contributed to enhanced fullness, highlighting the potential role of these noodles in promoting appetite control. Beyond their nutritional advantages, the use of sago and Bambara groundnut – both climate-resilient and locally available crops – supports diversification of raw materials and reduces dependence on imported wheat. Overall, this study provides evidence that climate-resilient crops can be successfully transformed into acceptable, nutrient-dense, and highly satiating instant foods. The sago–Bambara groundnut noodle represents a promising functional and sustainable alternative for future food innovation, particularly in regions vulnerable to climate change and food insecurity. Furthermore, future research is needed to optimize the physicochemical properties of the resulting flour, thereby improving the sensory and textural quality of the noodles.

RESEARCH FUNDING

The authors would like to express their sincere gratitude to the Directorate of Research, Technology, and Community Service (DRTPM), Ministry of Higher Education, Science, and Technology of the Republic of Indonesia (Kementerian Pendidikan Tinggi, Sains, dan Teknologi Republik Indonesia) for the financial support provided through the "Penelitian Tesis Magister (PTM)" or Master's Thesis Research scheme for the project entitled 'Development of Sago and Bambara Groundnut Noodles as a High-Fiber, Low Glycemic Index Alternative Food for Obesity and Diabetes' (Contract No. 006/C3/DT.05.00/PL/2025). The authors would also like to gratefully acknowledge IPB University for covering the Article Processing Charge.




Figure 3. Visual analogue scale for the fullness index at fasting intervals of 30, 60, 90, 120, 150, and 180 min after the consumption of white bread, wheat noodles and noodles made from sago and Bambara groundnut flours used in a 60:40 (w/w) ratio.

CONFLICT OF INTERESTS

The authors declare no conflict of interests.

INFORMED CONSENT

This research has obtained permission from the Commission on Research Ethics Involving Human Subjects, IPB University Number: 1633/IT3.KEPMSM-IPB/SK/2025.

ORCID IDs

N.M. Esa

Y.R. Indrining Tyas

Z. Nasution

E. Palupi

A. Pangestu

D.R. Tarigan

<https://orcid.org/0000-0003-1545-0306>

<https://orcid.org/0009-0007-7142-8142>

<https://orcid.org/0000-0001-8307-1012>

<https://orcid.org/0000-0003-2029-3106>

<https://orcid.org/0000-0002-6965-3988>

<https://orcid.org/0009-0009-5878-2387>

REFERENCES

- Agustia, F.C., Subardjo, Y.P., Sitasari, A. (2016). Formulation and characterization of free gluten high protein sago starch noodle substituted with beans flour. *Jurnal Gizi dan Pangan*, 11(3), 183–190 (in Indonesian, English abstract). <https://doi.org/10.25182/jgp.2016.11.3.183-190>
- Aremu, M.O., Aremu, S.O., Aliyu, I., Passali, D.B., Hussaini, M., Musa, B.Z., Salau, R.B. (2025). Amino acids profile and health attributes of Bambara groundnut (*Vigna subteranea* L.) and sesame (*Sesamum indicum* L.): a comparative study. *African Scientific Reports*, 4(1), art. no. 233. <https://doi.org/10.46481/asr.2025.4.1.233>
- Ariyathana, P., Mizera, P., Walkowiak, J., Dziedzic, K. (2025). Physicochemical and functional properties of soluble and insoluble dietary fibers in whole grains and their health benefits. *Foods*, 14, art. no. 2447. <https://doi.org/10.3390/foods14142447>
- Aussanasuwannakul, A., Teangpook, C., Treesuwan, W., Puntaburt, K., Butsuwan, P. (2022). Effect of the addition of soybean residue (okara) on the physicochemical, tribological, instrumental, and sensory texture properties of extruded snacks. *Foods*, 11(19), art. no. 2967. <https://doi.org/10.3390/foods11192967>
- AOAC (2005). *Official Methods of Analysis* (18th ed Association of Official Analytical Chemists International, Gaithersburg, MD, USA).
- Bayomy, H., Alamri, E. (2022). Technological and nutritional properties of instant noodles enriched with chickpea or lentil flour. *Journal of King Saud University Science*, 34(3), art. no. 101833. <https://doi.org/10.1016/j.jksus.2022.101833>
- Begna, T., Wakweya, R.B. (2026). Mitigating climate change impacts on food security via climate-smart agriculture. *Plant Stress*, 19, art. no. 101152. <https://doi.org/10.1016/j.plstress.2025.101152>
- Benitez-Alfonso, Y., Soanes, B.K., Zimba, S., Sinanaj, B., German, L., Sharma, V., Bohra, A., Kolesnikova, A., Dunn, J.A., Martin, A.C., Rahman, M.K., Saati-Santamaria, Z., Garcia-Fraile, P., Ferreira, E.A., Frazão, L.A., Cowling, W.A., Siddique, K.H.M., Pandey, M.K., Farooq, M., Varshney, R.K., Chapman, M.A., Boesch, C., Daszkowska-Golec, A., Foyer, C.H. (2023). Enhancing climate change resilience in agricultural crops. *Current Biology*, 33(23), R1246–R1261. <https://doi.org/10.1016/j.cub.2023.10.028>
- Bintoro, M.H., Nurulhaq, M.I., Pratama, A.J., Ahmad, F., Ayulia, L. (2018). Growing area of sago palm and its environment. In: H. Ehara, Y. Toyoda, D.V. Johnson (Eds.). *Sago Palm Multiple Contributions to Food Security and Sustainable Livelihoods*, Springer, Singapore, pp. 17–29. https://doi.org/10.1007/978-981-10-5269-9_2
- BSN (2018). SNI 3551:2018 Mi instan (Instant Noodle). Badan Standardisasi Nasional.
- Chakraborty, P., Witt, T., Harris, D., Ashton, J., Stokes, J.R., Smyth, H.E. (2019). Texture and mouthfeel perceptions of a model beverage system containing soluble and insoluble oat bran fibres. *Food Research International*, 120, 62–72. <https://doi.org/10.1016/j.foodres.2019.01.070>
- Chong, W.K., Mah, S.Y., Easa, A.M., Tan, T.C. (2019). Thermal inactivation of lipoxygenase in soya bean using superheated steam to produce low beany flavour soya milk. *Journal of Food Science and Technology*, 56(9), 4371–4379. <https://doi.org/10.1007/s13197-019-03905-4>
- Dewita, Desmelati, Edison, Syahrul (2019). The development of instant (dried) sago noodles fortified with functional components. *IOP Conference Series: Earth and Environmental Science*, 253, art. no. 012018. <https://doi.org/10.1088/1755-1315/253/1/012018>
- Estiasih, T., Hidayat, R., Putri, R.N., Ahmadi, K.A., Ali, D.Y., Laeliocattleya, R.A. (2025). *Moringa oleifera* leaf powder colour improvement and lipoxygenase inactivation by different pHs and water-blanching times. *CyTA – Journal of Food*, 23(1), art. no. 2586311. <https://doi.org/10.1080/19476337.2025.2586311>
- Farhan, I., Sarayrah, H., Hayek, W., Alkhasoneh, H., Almayouf, F. (2025). Analyzing wheat production in Jordan: The role of population dynamics, climate variability, and GIS-based projections. *Sustainability*, 17(8), art. no. 3493. <https://doi.org/10.3390/su17083493>
- Forde, C.G. (2018). Chapter 7: Measuring satiation and satiety. In: G. Ares, P. Varela (Eds.). *Methods in Consumer Research*, Volume 2, Woodhead Publishing, pp. 151–182. <https://doi.org/10.1016/B978-0-08-101743-2.00007-8>
- Gerstein, D.E., Woodward-Lopez, G., Evans, A.E., Kelsey, K., Drewnowski, A. (2004). Clarifying concepts about macronutrients' effects on satiation and satiety. *Journal of the American Dietetic Association*, 104(7), 1151–1153. <https://doi.org/10.1016/j.jada.2004.04.027>
- Hervik, A.K., Svihus, B. (2019). The role of fiber in energy balance. *Journal of Nutrition and Metabolism*, 2019, art. no. 4983657. <https://doi.org/10.1155/2019/4983657>
- Holt, S.H.A., Miller, J.C.B., Petocz, P., Farmakalidis, E. (1995). A satiety index of common foods. *European Journal of Clinical Nutrition*, 49(9), 675–690.
- Huh, I.S., Kim, H., Jo, H.K., Lim, C.S., Kim, J.S., Kim, S.J., Kwon, O., Oh, B., Chang, N. (2017). Instant noodle consumption is associated with cardiometabolic risk factors among college students in Seoul. *Nutrition Research and Practice*, 11(3), 232–239. <https://doi.org/10.4162/nrp.2017.11.3.232>
- Iju, S., Sarinah, Rejeki, S. (2025). Formulation of instant baby porridge from red rice flour substituted with moringa (*Moringa oleifera*) leaf flour and natural banana (*Musa paradisiaca*) flavor as an alternative complementary food to breast milk. *Jurnal Gizi dan Pangan*, 3(2), 241–260 (in Indonesian, English abstract).
- Indonesian Food and Drug Authority (2022). Regulation of Indonesian Food and Drug Authority Number 1 of 2022 About Monitoring Claims on Processed Food Label and Food Advertisement. Jakarta: Indonesian Food and Drug Authority.
- ISO 8586 (2014). Sensory analysis – General guidelines for the selection, training and monitoring of selected assessors and expert sensory assessors.
- Jahanshiri, E., Goh, E.V., Wimalasiri, E.M., Azam-Ali, S., Mayes, S., Suhairi, T.A.S.T.M., Nizar, N.M.M., Sinin, S.S.M. (2022). The potential of bambara groundnut: an analysis for the People's Republic of China. *Food and Energy Security*, 11(2), art. no. e358. <https://doi.org/10.1002/fes3.358>
- Jamayah, H., Geeta, A., Safiza, M.N., Khor, G.L., Wong, N.F., Kee, C.C., Rahmah, R., Ahmad, A.Z., Suzana, S., Chen, W.S., Rajaah, M., Adam, B. (2010). Reliability, technical error of measurements and validity of length and weight measurements for children under two years old in Malaysia. *Medical Journal of Malaysia*, 65(Suppl. A), 131–137.
- Kudre, T.G., Benjakul, S. (2013). Effects of binary organic solvents and heating on lipid removal and the reduction of beany odour in Bambara groundnut (*Vigna subterranea*) flour. *Food Chemistry*, 141(2), 1390–1397. <https://doi.org/10.1016/j.foodchem.2013.04.047>
- Makhuvha, M.C., Laurie, S., Mosala, M. (2024). Effect of orange-fleshed sweet potato (*Ipomoea batatas*)–Bambara groundnut (*Vigna subterranea*) composite flour on quality properties of pasta. *Journal of Food Science*, 89(11), 7348–7359. <https://doi.org/10.1111/1750-3841.17432>
- Maphosa, Y., Jideani, V.A., Maphosa, L. (2022). Bambara groundnut production, grain composition and nutritional value: Opportunities for improvements. *Journal of Agricultural Science*, 160(6), 448–458. <https://doi.org/10.1017/S0021859622000521>
- Mashau, M.E., Mukwevho, T.A., Ramashia, S.E., Siwela, M. (2022). The influence of Bambara groundnut (*Vigna subterranea*) flour on the nutritional, physical and antioxidant properties of steamed bread. *CyTA – Journal of Food*, 20(1), 259–270. <https://doi.org/10.1080/19476337.2022.2130435>
- Ministry of Health Indonesia (2019). National Basic Health Research Report 2018. Jakarta: Ministry of Health Indonesia. <https://repository.kemkes.go.id/book/1323>
- Mo, Y., Ma, X., Lu, J., Shen, Y., Wang, Y., Zhang, L., Lu, W., Zhu, W., Bao, Y., Zhou, J. (2021). Defining the target value of the coefficient of variation by continuous glucose monitoring in Chinese people with diabetes. *Journal of Diabetes Investigation*, 12(6), 1025–1034. <https://doi.org/10.1111/jdi.13453>
- Moshawih, S., Kong, J.P.T., Goh, B.H., Renganathan, E., Goh, H.P., Goh, K.W., Ming, L.C. (2025). Exploring the nutritional, cultural, and industrial significance of *Metroxylon sagu*. *Discovery Food*, 5, art. no. 337. <https://doi.org/10.1007/s44187-025-00638-6>

33. Nguyen, T.N., Tran, T.T.T., Le, V.V.M. (2025). Effects of blanching conditions on the enzyme inhibition and antioxidant loss in rambutan (*Nephelium lappaceum* L.) seeds. *Polish Journal of Food and Nutrition Sciences*, 75(1), 49–59. <https://doi.org/10.31883/pjfn/200589>
34. Nolan, S.J., Hambleton, I., Dwan, K. (2016). The use and reporting of the cross-over study design in clinical trials and systematic reviews: a systematic assessment. *PLoS One*, 11(7), art. no. e0159014. <https://doi.org/10.1371/journal.pone.0159014>
35. Palamae, S., Suyapoh, W., Boonrat, O., Zhang, B., Amin, M., Buatong, J., Benjakul, S. (2024). Activity of bambara groundnut seed coat extract against *Shewanella* species: efficacy and mechanisms of action. *Foods*, 13(21), art. no. 3516. <https://doi.org/10.3390/foods13213516>
36. Palupi, E., Nurdin, N.M., Mufida, G., Valentine, F.N., Pangestika, R., Rimba-wan, Sulaeman, A., Briawan, D., Filianty, F. (2024). High-fiber extruded purple sweet potato (*Ipomoea batatas*) and kidney bean (*Phaseolus vulgaris*) extends the feeling of fullness. *Polish Journal Food and Nutrition Sciences*, 74(1), 82–91. <https://doi.org/10.31883/pjfn/183995>
37. Pretorius, B., Otto, M., Schönfeldt, H.C. (2023). Antinutrients and metabolomic compounds of Bambara groundnut (*Vigna subterranean*) as affected by traditional processing by smallholder farmers. *Journal of Food Science*, 88(8), 3435–3444. <https://doi.org/10.1111/1750-3841.16698>
38. Ridwan, M., Palupi, E., Setiawan, B., Heksana, R.A.R. (2024). Bambara bean substitution improves the nutritional content and increases the satiety index of purple sweet potato bread. *Preventive Nutrition and Food Science*, 29(4), 512–521. <https://doi.org/10.3746/pnf.2024.29.4.512>
39. Salihu, S., Elezaj, B., Qorri, D., Gashi, N. (2025). Exploring instant noodle consumption patterns and consumer awareness in Kosovo. *Foods*, 14(24), art. no. 4245. <https://doi.org/10.3390/foods14244245>
40. Savastano, D.M., Hodge, R.J., Nunez, D.J., Walker, A., Kapikian, R. (2014). Effect of two dietary fibers on satiety and glycemic parameters: a randomized, double-blind, placebo-controlled, exploratory study. *Nutrition Journal*, 13, art. no. 45. <https://doi.org/10.1186/1475-2891-13-45>
41. Septiani, M., Sugiatmi, S. (2026). Relationship between ultra-processed food consumption patterns and physical activity with overnutrition in adolescents of SMK Kesehatan YHB Sukabumi. *Jurnal Ilmu Gizi dan Dietetik*, 5(1), 423–430 (in Indonesian, English abstract). <https://doi.org/10.25182/jigd.2026.5.1.423-430>
42. Singthong, J., Oonmetta-aree, J., Oonsivilai, R. (2026). Impact of red kidney bean flour substitution on the nutritional, physicochemical, and functional properties of gluten-free noodles. *Food Chemistry Advances*, 11, art. no. 101279. <https://doi.org/10.1016/j.focha.2026.101279>
43. Tan, X.L., Azam-Ali, S., Goh, E.V., Mustafa, M., Chai, H.H., Ho, W.K., Mayes, S., Mabhaudhi, T., Azam-Ali, S., Massawe, F. (2020). Bambara groundnut: An underutilized leguminous crop for global food security and nutrition. *Frontiers in Nutrition*, 7, art. no. 601496. <https://doi.org/10.3389/fnut.2020.601496>
44. Ulijaszek, S.J., Kerr, D.A. (1999). Anthropometric measurement error and the assessment of nutritional status. *British Journal of Nutrition*, 82(3), 165–177. <https://doi.org/10.1017/S0007114599001348>
45. WHO (2024). World Health Organization. Overweight and Obesity. UK. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>