

Review Article

Freshwater Snails in Malaysia: Diversity, Roles in Trematode Transmission, and Agricultural Impacts

Nazir Ahmad Tookhy^{1,2}, Md Isa Nur-Mahiza^{2*}, Emmanuel Busayo Ibitoye³,
Hakim Idris Lokman⁴ and Shoaib Ahmad Shakhes¹

¹Department of Paraclinic, Faculty of Veterinary Science, Herat University, Herat, Afghanistan

²Department of Veterinary Pathology and Microbiology, Faculty of Veterinary Medicine, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

³Department of Theriogenology and Animal Production, Faculty of Veterinary Medicine, Usmanu Danfodiyo University Sokoto, Nigeria

⁴Department of Veterinary Pre-Clinical Science, Faculty of Veterinary Medicine, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

ABSTRACT

Freshwater snails play a critical role as intermediate hosts in the transmission of trematode diseases, affecting human and animal health and posing challenges to agriculture. This review summarises studies on freshwater snails and trematode diseases in Malaysia from 1973 to 2024, based on comprehensive literature searches across multiple databases, including Web of Science, PubMed, Scopus, and Google Scholar. A total of 32 snail species were identified, with *Radix rubiginosa* hosting five types of cercariae and *Robertsia silvicola* acting as a host for furcocercous cercariae. Economically significant species, such as *Pomacea maculata* and *Pomacea canaliculata*, were highlighted for their agricultural impact. While treatments for trematode infections are available, prevention through controlling snail populations remains essential due to the complex transmission involving reservoir hosts. This study underscores the zoonotic potential of trematode parasites in Malaysia and identifies a significant research gap regarding intermediate host snails.

Keywords: Freshwater snails, trematode, Malaysia, overview

ARTICLE INFO

Article history:

Received: 23 September 2024

Accepted: 26 December 2024

Published: 30 June 2025

DOI: <https://doi.org/10.47836/pitas.48.4.09>

E-mail addresses:

tookhy@yahoo.com (Nazir Ahmad Tookhy) numahiza@upm.edu.my (Md Isa Nur-Mahiza) eb.ibitoye@gmail.com (Emmanuel Busayo Ibitoye) hakim_idris@upm.edu.my (Hakim Idris Lokman) Sshakhes@yahoo.com (Shoaib Ahmad Shakhes)

* Corresponding author

INTRODUCTION

Snails are prevalent in freshwater environments across tropical and subtropical regions, notably Southeast Asia (Böhm et al., 2021). Scholars suggest that more than 350 snail species may have medical or

veterinary significance (Madsen & Hung, 2014). Around 45 species from 26 genera and seven families in Southeast Asia are implicated in transmitting trematodes (Lu et al., 2018; Madsen & Hung, 2014).

Mud and freshwater snails act as obligatory intermediate hosts for approximately 71 trematode parasites (Lu et al., 2018; Phiri et al., 2007). These parasites include members of the families Clinostomidae, Echinostomatidae, Fasciolidae, Schistosomatidae, Paramphistomidae, and Pronocephalidae (Tookhy et al., 2023; Saijuntha et al., 2021). In humans and animals, snails, for example, the *Bulinus*, *Biomphalaria*, *Lymnaea* and *Oncomelania* (Tookhy et al., 2023; Martin & Cabera, 2018; Dida et al., 2014; Dung et al., 2013) serve as the common intermediate parasitic hosts. Furthermore, according to Yahaya et al. (2017) and Greene (2018), some particular species of snails, for example, apple snails, constitute a major challenge as they can destroy rice plantations.

As reported by Lu et al. (2018), Prasopdee et al. (2015), and Tookhy et al. (2023), infections by some trematodes are an example of parasitic diseases transmitted by snails, and they represent an important health challenge to people and livestock, which can lead to socioeconomic losses. According to Martin and Cabera (2018), the exchange of nutrients between aquatic and terrestrial systems is being expedited by the roles played by some freshwater snails in the aquatic environment.

Among the foregoing is a critical fascioliasis disease by *Fasciola hepatica* and *Fasciola gigantica*, from which man and livestock suffer (World Health Organization [WHO], 2020). Almost two million and four hundred thousand people in about 700 nations of the world, by estimation, are victims of fascioliasis (WHO, 2018). In the same vein, other zoonoses and neglected tropical diseases, such as schistosomiasis, kill about 200 million people yearly, affect around 230 million individuals, and more than 700 million people are at higher risk of being infected globally (Colley et al., 2014; WHO, 2016). The diagnosis of schistosomiasis has been made more difficult by the massive human population drift to the western part of the globe.

Although there are reports on human and livestock trematode infections in Malaysia, there is, however, grossly inadequate research that specifically addresses the role and impact of snails in the transmission of this parasite. Attwood et al. (2005), Greer et al. (1988), and Tookhy et al. (2023) have documented the significance of human medical and veterinary trematodes. In-depth findings into the roles played by snails in the transmission of this parasite in the environment are lacking. This is a knowledge gap that has hampered the proper understanding and implementation of appropriate snail control measures to manage and prevent disease transmission.

Therefore, this work aims to provide a comprehensive overview of Malaysian freshwater snails and their role as intermediate hosts in the transmission of trematode parasites. By a critical review of the ecology and epidemiology of freshwater snails, their importance in human and livestock health will be discussed, as well as specific control measures.

OVERVIEW OF TREMATODE INFECTIONS IN MALAYSIA

Different species of trematodes, such as *Fasciola* spp., have been documented in various locations and have been encountered in many states in Malaysia. Among the cattle in Terengganu, and according to studies by Khadijah et al. (2017), Rita et al. (2017), and Saad et al. (2019), *Fasciola* infection has been established and reported; the infection has also been established in buffalo along with cattle in Sabah (Kamaruddin et al., 2021). In Pahang, as documented by Rosilawati et al. (2017), *Fasciola* infections were predominant in buffalo, while the study of Diyana et al. (2020) has reported the infection in cattle and buffalo in Selangor and Perak. In Kuala Lumpur, *Fasciola* infections have been reported in humans by Naresh et al. (2006). Infection by the rumen fluke has been reported in cattle and buffalo from Terengganu (Kahdijah et al., 2021; Rita et al., 2017; Saad et al., 2019); Sarawak (Harizt et al., 2021) and Perak (Tookhy et al., 2024). Furthermore, in Perak, a 2% mortality rate was reported among local cattle (Debbra et al., 2018), and in Johor, among buffalo (Jamnah et al., 2013). Infection due to *Paragonimus westermani* was documented in Selangor when *Brotia costula* was established as an intermediate host (Kim, 1978); this infection has also been found in tigers in Selangor (Lee, 1965).

Schistosoma spp. infections have been documented across different regions and hosts, with *Schistosoma malayensis* infections in humans reported in Selangor and Pahang, the latter also showing infections with *Robertsia* spp. (Attwood et al., 2005; Greer et al., 1988; Latif et al., 2013). *Schistosoma japonicum* infections have been observed in humans in Peninsular Malaysia (Greer & Anuar, 1984), while a variety of *Schistosoma* spp. and other trematodes have been reported in wildlife, with no associated mortality. Notably, the intermediate host is unknown in Malaysia for most of the above trematodes.

The overview of the pattern of distribution and identification methods, comprising conventional morphological assessment and molecular analyses such as polymerase chain reaction (PCR), for different species of snails found in various aquatic ecosystems across Malaysia is shown in Table 1. Details such as species nomenclature, geographical location, preferred habitats (wetlands, rivers or streams), and identifiable types of cercaria, where possible. Notably, this study from Peninsular Malaysia highlighted a relatively small

Table 1
Snail species reported in different water resources based on different methods and their larval stage

Snail	Location	Water resources	Cercarial	Identification	References
<i>Radix rubiginosa</i>	Perak	Wetlands	<i>Echinostome, gymnocephalus, xiphidiocercariae, furcocercous</i>	Morphological ID and PCR (COI)	Tookhy et al. (2023)
<i>Robertsia silvicola</i>	Perak	Stream	<i>Furcocercous cercaria</i>	Morphological ID	Attwood et al. (2005)

Table 1 (continue)

Snail	Location	Water resources	Cercarial	Identification	References
<i>Brotia costula</i>	Kuala Lumpur	Stream		Morphological ID	Kim (1978)
<i>Melanoides tuberculata</i>	Perak	Stream			Attwood et al. (2005)
<i>Brotia costula</i>					
<i>Thiara scabra</i>					
<i>Pila ampullaceal</i>					
<i>Bithynia</i>					
<i>Pygmaea</i>					
<i>Stenothyra cambodiensis</i>					
<i>Gyraulus chinensis</i>					
<i>R. gismanni</i>					
<i>R. kaporensis</i>					
<i>Pomacea maculata</i>	Peninsular	Rivers, streams, lakes, ponds	-	PCR (COI)	Hah et al. (2022)
<i>Pomacea canaliculata</i>					
<i>Pomacea maculata</i>	Selangor	Wetlands	-	PCR (COI)	Phoong et al. (2018)
<i>Pomacea canaliculata</i>					
<i>P. canaliculate</i>	Peninsular	Pond, lakes, rice fields	-	SM and PCR (COI)	Rao et al. (2018)
<i>P. maculata</i>					
<i>Pomacea maculata</i>	Selangor,	Rice fields	-	Morphological ID	Arfan et al. (2014)
<i>Pomacea canaliculata</i>	Perak, Penang, Kedah and Kelantan				
<i>Physella acuta</i>	Sabah	Different water bodies	-	Morphological ID	Ng et al. (2017)
<i>Indoplanorbis exustus</i>					
<i>Melanoides tuberculata</i>					
<i>Sinotaia guangdungensis</i>					
<i>Pila ampullacea</i>					
<i>Pila scutate</i>					
<i>Clea banguyensis</i>					
<i>Neritina pulligera</i>					
<i>Septaria porcellana</i>					
<i>Vittina variegata</i>					
<i>Coromandeliana</i>					
<i>Sulcospira pageli</i>					
<i>Paludomus everetti</i>					
<i>Paludomus luteus</i>					
<i>Mieniphotia scabra</i>					
<i>Tarebia granifera</i>					

number of freshwater snail species, with only 15 species identified through a combination of molecular and morphological approaches. Through morphological evaluation, 17 species were identified in Sabah, a region in Malaysia rich in a variety of freshwater snails. It was observed that cercarial types were reported in only two species, *R. rubiginosa* and *R. silvicola*, suggesting that they play a role in Malaysian freshwater ecosystems. All the cercaria reported are zoonotic, suggesting they are of public importance.

THE IMPORTANT SNAIL FAMILIES PRESENT IN MALAYSIAN FRESHWATER ENVIRONMENTS

Family: Lymnaidae

Genus: Lymnea

Species: *Radix rubiginosa*

Earlier, the presence of *R. rubiginosa* and cercariae in the wetlands of Perak was established using both morphological and molecular (PCR, ITS2) methodologies by Tookhy et al. (2023). The identification of cercariae, such as *Echinostome*, *Gymnacephalous*, *Xiphidiocercariae*, and *Furocercous*, corroborates the ecological diversity within this environment (Tookhy et al., 2023). Additionally, Tookhy et al. (2023) have reported that *R. rubiginosa* can accommodate the larval stage of *Schistosoma* spp., which causes human cercarial dermatitis in Malaysia. The freshwater snail of the *Lymnaeidae* family has gained enormous interest because of its involvement in the life cycle of different trematodes, which have great biomedical and veterinary importance (Bargues et al., 2016; Saijuntha et al., 2021).

Family: Pomatiopsidae

Genus: Robertsiella

Species: *Robertsiella silvicola*

As reported by Attwood et al. (2005), the small size, lack of a defined spiral macrosculpture on the shell, and distinctive style characteristics are suggestive of *R. silvicola*, differentiating it from other species in the *Robertsiella* genus. This species of snail can be found in different aquatic environments ranging from diminutive, spring-fed, first-order streams to meagre rivulets originating from forest hillsides. The occurrence of this snail species is restricted to limestone regions surrounding the mountainous bases of the Perak and Kedah states in West Malaysia. It is important that *R. silvicola* display affinity to serve as an intermediate host for the parasitic blood fluke *S. malayensis* that infects humans and livestock (Attwood et al., 2005).

Family: Ampullariidae

Genus: Pomacea

Species: *Pomacea maculata* and *Pomacea canaliculate*

It has been reported that *Pomacea maculate* and *P. canaliculate* have been isolated in diverse bodies of water such as streams, rivers, lakes, wetlands and ponds within Peninsular

Malaysia (Hah et al., 2022; Rao et al., 2018) as well as in Selangor (Arfan et al., 2014; Phoong et al., 2018). In this study, PCR (COI) confirmed the presence of these two species of snails, suggesting that they can adapt and be found in various ecosystems.

The financial loss and damage caused by apple snails in rice farming are enormous; hence, their management has taken centre stage (Yahaya et al., 2017). The feeding patterns of these two snail species, which consume aquatic plants, thereby altering the food chain, may significantly impact the community structure, water quality, and functionality of aquatic ecosystems in wetlands (Horgan et al., 2014).

Family: **Thiaridae**

Genus: **Melanoides**

Species: ***Melanoides tuberculata***

The *M. tuberculata*, commonly known as red-rimmed Melania or the Malaysia Trumpet Snail (Müller, 1774), is a benthic freshwater gastropod of the *Thiaridae* family. According to a report by Krailas et al. (2014), *M. tuberculata* is indigenous to Asia and Africa. Naturally, this species is widely distributed, including Africa, the East Indies, southern Asia, northern Australia, Madagascar, the Mediterranean, and the Pacific Islands (Maciaszek et al., 2019; Facon et al., 2003).

The presence of *M. tuberculata* (Müller, 1774) has been documented by Piechocki et al. (2003) in Europe (Hungary, Germany, Malta, the Netherlands, Spain and Austria), where it has been identified as an intermediate host for many trematode species, some of these trematodes are of paramount importance in human and livestock health (Francis, 2012; Lopes et al., 2021; Post et al., 2022). In Malaysia, and as documented by Ng et al. (2017), the prevalence of *M. tuberculata* cut across various regions, such as offshore islands like Tenom, Kota Kinabalu, Tuaran, Pulau Gaya, Tawau, Pulau Tiga, Kota Belud, Pulau Bodgaya near Semporna, and Pulau Bohey Dulang. It inhabits rivers, paddy fields, and man-made concrete drains.

Family: **Planorbidae**

Genus: **Indoplanorbis**

Species: ***Indoplanorbis exustus***

Indoplanorbis exustus is widely found throughout Southeast Asia, especially in rice plantations, and it plays the role of an intermediate host for some trematodes (Dumidae et al., 2024; Bawm et al., 2022; Nguyen et al., 2022). Following harvest, farmers often introduce ducks into their rice fields to consume these snails, thereby facilitating the maintenance and completion of the parasitic trematode life cycle through a zoonotic route. Thailand has reported a notable prevalence of *Echinostoma* infections in free-ranging ducks, as documented by Saijuntha et al. (2013). *Indoplanorbis exustus* also act as intermediate host

of furcocercous cercaria (*Schistosoma cercariae*) (Dumidae et al., 2024). In Peninsular Malaysia, this species has been demonstrated to harbour *S. spindale*, which is responsible for causing cercarial dermatitis in humans who become infected (Chiew et al., 2009).

ROLE OF SNAILS IN AGRICULTURE

Freshwater snails exhibit a dual role in agriculture, beneficial and detrimental, depending on the context. Economically, they are valued in rural communities as a source of food and traditional medicine (Gupta & Khanal, 2024). Nonetheless, a species such as *Pomacea canaliculata*, commonly called the golden apple snail, is a stubborn pest and a threat to agricultural success, predominantly in rice cultivation, by extensively damaging the shoots of young rice plantations, which results in a gross reduction in crop yields (Joshi et al., 2020). The combination of these roles highlights the complication of their impacts on the environment and the economy.

Yusa et al. (2006) have highlighted that *Pomacea canaliculata* is a main pest in Southeast Asia; however, it is less encountered, with less associated damage in other regions with lower annual precipitation, like parts of South Asia. Variations in climate, crop management, or the presence of a predator could have been the reason for this. Therefore, this suggests that the environment and geographical locations might have meaningfully impacted the level of infestation and resultant damage occasioned by freshwater snails.

Considering the level of loss in high-yielding crops, as reported by Kumar (2020) and Yusa et al. (2006), this urgently calls for effective and operational control measures. However, the differences in loss levels across regions and studies could be attributed to variations in farming systems, snail population densities, and mitigation strategies. For instance, integrated pest management methods that incorporate natural predators and manual removal are promising in reducing infestations (Thakuri et al., 2019). The report states that snails contaminate agricultural by-products with zoonotic parasites after they have damaged the by-product (Kumar, 2020). In the area of food security and livestock health, this perspective could have broader implications, especially in resource-limited settings. It is believed that if this study is compared with other studies on the dynamics of parasite transmission in other snail species, there could be better clarity. Unexpectedly, some studies have reported increased snail populations during dry spells within the rainy season (Faiz, 2010). It could indicate adaptive behaviour or resilience in certain species, necessitating further research on the triggers and ecological factors that enable such growth.

One potential strategy for mitigating the negative impacts of freshwater snails while leveraging their benefits involves utilising them as food and animal feed sources. This approach reduces their population and provides economic incentives for farmers. There could be better insights if studies on the nutritional and economic feasibility of snail-derived feed were compared against those of traditional livestock feed.

THE ROLE OF SNAIL IN MEDICINE AND VETERINARY PRACTICE

From time immemorial, snails have been known for their potent therapeutic properties, principally in traditional medicine. For instance, the mucus of some snail species, such as *Helix aspersa* and *Eremina desertorum*, is rich in bioactive compounds that have anti-inflammatory and antimicrobial effects by regulating transforming growth factor beta 1 (TGF- β 1) and vascular endothelial growth factor (VEGF) gene expression that promotes wound healing (El-Zawawy & Mona, 2021). Also, Noothuan et al. (2021) have reported the antibacterial, antioxidant, and anti-tyrosinase properties demonstrated by mucus from *Hemiplecta distincta* and *Lissachatina fulica*, while available documents on *Helix aspersa* show it is effective in the treatment of gastric ulcers and reduction of oxidative stress, inflammation, and skin conditions (Gugliandolo et al., 2021).

Furthermore, lipids in *Bellamyia benghalensis*, comprising oleic acid and cyclopropane fatty acids, possess anti-inflammatory potential for managing immune-related diseases, such as rheumatism, through the modulation of macrophage activity (Bhattacharya et al., 2014). Antimicrobial properties against infection, such as whooping cough, have been demonstrated in snail meat (Cobbinah et al., 2008). Furthermore, cancer-combating potential has been raised in the mucus of *Achatina fulica* (EFSA Panel on Nutrition, Novel Foods and Food Allergens et al., 2024). However, concerns about the consistency and safety of *Helix aspersa* maxima mucus as a novel food, according to the European Food Safety Authority (Adeyeye et al., 2020). Despite these challenges, snails are promising for medicinal purposes and as a novel food or feed source.

In animal production, snails can serve as an unconventional protein source, as they can be raised on low-grade biowastes and bioresources (Pathak et al., 2024). Therefore, they are a valuable and sustainable option for animal feed, as they reduce reliance on conventional feed resources (International Feed Industry Federation, 2021). Conversely, these snails also serve as a reservoir for zoonotic parasites and other pathogenic microbes, constituting health threats such as Fascioliasis, Schistosomiasis, Angiostrongyliasis and microbial diseases like Gastroenteritis and Rat-bite fever (Tookhy et al., 2024; Pathak et al., 2024). Specifically, ruminants and humans are susceptible to liver fluke disease (Lee et al., 2017). While snail farming presents opportunities for sustainable food and feed production, these health risks will necessitate careful and effective management to ensure the safe commercialisation of snails.

GENETIC CHARACTERISATION OF SNAIL

The extinction of some species has been a continuous event in the long history of nature, driven by various natural processes and human activities (Ceballos et al., 2017; Barnosky et al., 2011). Currently, the five leading drivers of species extinction are well-documented: habitat loss, invasive species, exploitation, disease, and climate change (Sodhi et al., 2009).

The most diverse group of animals, next to arthropods, is the molluscs, which are principally vulnerable, with an estimated number of known species falling between 70,000 and 120,000. The International Union for Conservation of Nature (IUCN) Red List has documented about 83,125 mollusc species, out of which 8,934 have been evaluated, with about 2,340 species (11%) currently on the verge of extinction (Chapman, 2009; IUCN, 2021).

Freshwater snails are a critical component of marine bionetworks. However, their environment is highly fragile, and coupled with the rising pressures due to man's activities, they can easily become extinct (Johnson et al., 2013; Strayer & Dudgeon, 2010). For example, lymnaeid snails display considerable eco-phenotypic plasticity with highly varying shell morphology, which is intensely affected by the environment (Hurtrez-Bouss et al., 2005). This variability can confound the taxonomy using only shell characteristics, as morphological variations may lead to misclassification (Vinarski et al., 2020).

The standard tools for accurately identifying species, particularly when morphological characteristics are insufficient, are molecular phylogenetics and molecular taxonomy (Japa et al., 2021). The commonly employed genetic markers like the internal transcribed spacer 2 (ITS2), mitochondrial cytochrome c oxidase subunit I (COI) and 16s rRNA genes were known to be reliable in the study of lymnaeid snails (Tookhy et al., 2023; Japa et al., 2021). These are valuable molecular markers for species identification, investigating genetic divergence, and understanding the structural relationships within snail populations. This methodology offers a more accurate and reliable understanding of the taxonomy and genetics of freshwater snails (Mirfendereski et al., 2021; Vinarski et al., 2020).

SNAIL CONTROL

Methods for the control of snails include physical, chemical, and biological methods (Lu et al., 2018). Physical control measures target the reduction of snail populations by effective management of the environment (Garba Djirmay et al., 2024). For example, the removal of natural bodies of water, like the marshes and ponds, and controlling human settlements in high-risk zones are effective approaches. In other areas, the spread of *Schistosoma haematobium* and *Schistosoma japonicum* has been significantly brought low employing appropriate drainage systems and environmental engineering, such as upsetting epilithic snail habitats by the use of boat-mounted rototillers or tractors with rakes, which is also effective in eliminating a substantial portion of snail populations (Li et al., 2016). Other methods, such as removing bird roosting sites, introducing mechanised farming, and rotating between aquatic and xerophytic crops, help further reduce snail populations (Leighton et al., 2000).

In chemical control, molluscicides are a widely used method since they rapidly decimate snail populations, thereby reducing disease transmission (Xia et al., 2014). However, its long-lasting effect is a function of sustained application (Maes et al., 2021).

Chemical control measures typically involve the use of synthetic or natural molluscicides, and this is one of the most effective methods for managing snail populations (Xia et al., 2014). Since the 1950s through the 1970s, compounds such as sodium pentachlorophenate (NaPCP), copper sulfate, Ntritylmorpholine, and niclosamide (Bayluscide) were widely used, especially in the control of snails responsible for the transmission of schistosomiasis in regions of Asia, Africa, and South America (King et al., 2015).

The biological control of molluscs is a more sustainable approach, where natural predators, parasites, or competitors to subdue the number of snails are introduced (Younes et al., 2017). Predatory fish, ducks, or some kind of insects can be employed to control the snail population; nonetheless, success is a function of the availability of appropriate predators and the ability to maintain ecological balance (Sokolow et al., 2015). Predatory prawns, like *Macrobrachium vollehoveni*, have been used to reduce infected snails and schistosomiasis transmission in Senegal (Sokolow et al., 2014). The water bug *Sphaerodema urinator* and black carp (*Mylopharyngodon piceus*) have been reported to effectively control snails that host parasitic diseases (Younes et al., 2017). Even though these techniques display potential, careful management is required to prevent negative impacts on human health. Biological control benefits humans and the environment (Hung et al., 2013).

CONCLUSION

This literature review on freshwater snails and trematode diseases in Malaysia has highlighted the complex roles these snails play in human and animal health, as well as in agriculture. The findings of this study underscore the multiplicity of snail species in the region and their association with diverse trematode infections, including zoonotic parasites. Moreover, the economic importance of certain snail species in agriculture stresses the need for effective control procedures to prevent the spread of risks. Conversely, despite advancements in medical treatments for trematode diseases, preventive schemes that target intermediate hosts are of great importance, given the complexity of transmission dynamics involving the final hosts. The paucity of research centred on snails as intermediate hosts suggests an essential area for further exploration to improve the understanding of disease transmission, hence facilitating concerted control efforts. Employing integrated control methods, physical, chemical, and biological, can help avoid the negative impacts of snails while promoting sustainable practices. A well-adjusted approach is fundamental to maximising the benefits of freshwater snails while minimising their challenges.

ACKNOWLEDGEMENTS

This study received support from the Universiti Putra Malaysia (Grant number: 9559100). The lead author appreciates the Ministry of Higher Education, Afghanistan, for its involvement in this manuscript.

REFERENCES

- Adeyeye, S. A. O., Bolaji, O. T., Abegunde, T. A., & Adesina, T. O. (2020). Processing and utilization of snail meat in alleviating protein malnutrition in Africa: A review. *Nutrition & Food Science*, 50(6), 1085-1097. <https://doi.org/10.1108/NFS-10-2019-0303>
- Arfan, A., Muhamad, R., Omar, D., Azwady, A., & Manjeri, G. (2014). Distribution of two *Pomacea* spp. in Rice Fields of Peninsular Malaysia. *Annual Research and Review in Biology*, 4(24), 4123-4136. <https://doi.org/10.9734/arrb/2014/11398>
- Attwood, S. W., Lokman, H. S., & Ong, K. Y. (2005). *Robertsia silvicola*, a new species of triculine snail (Caenogastropoda: Pomatiopsidae) from peninsular Malaysia, intermediate host of *Schistosoma malayensis* (Trematoda: Digenea). *Journal of Molluscan Studies*, 71(4), 379-391. <https://doi.org/10.1093/mollus/eyi040>
- Bargues, M. D., Malandrini, J. B., Artigas, P., Soria, C. C., Velásquez, J. N., Carnevale, S., Mateo, L., Khoubbane, M., & Mas-Coma, S. (2016). Human fascioliasis endemic areas in Argentina: Multigene characterisation of the lymnaeid vectors and climatic-environmental assessment of the transmission pattern. *Parasites and Vectors*, 9(1), 1-20. <https://doi.org/10.1186/s13071-016-1589-z>
- Bawm, S., Khaing, N. H. E., Win, S. Y., Thein, S. S., Khaing, Y., Thaw, Y. N., Soe, N. C., Chel, H. M., Hmoon, M. M., Hayashi, N., Htun, L. L., Katakura, K., Nonaka, N., & Nakao, R. (2022). Morphological and molecular identification of trematode cercariae related with humans and animal health in freshwater snails from a lake and a dam in Myanmar. *Parasitology Research*, 121(2), 653-665. <https://doi.org/10.1007/s00436-022-07428-4>
- Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O., Swartz, B., Quental, T. B., & Ferrer, E. A. (2011). Has the Earth's sixth mass extinction already arrived? *Nature*, 471(7336), 51-57. <https://doi.org/10.1038/nature09678>
- Bhattacharya, S., Chakraborty, M., Bose, M., Mukherjee, D., Roychoudhury, A., Dhar, P., & Mishra, R. (2014). Indian freshwater edible snail *Bellamyia bengalensis* lipid extract prevents T cell-mediated hypersensitivity and inhibits LPS-induced macrophage activation. *Journal of Ethnopharmacology*, 157, 320-329. <https://doi.org/10.1016/j.jep.2014.09.012>
- Böhm, M., Dewhurst-Richman, N. I., Seddon, M., Ledger, S. E. H., Albrecht, C., Allen, D., Bogan, A. E., Cordeiro, J., Cummings, K. S., Cuttelod, A., Darrigran, G., Darwall, W., Fehér, Z., Gibson, C., Graf, D. L., Köhler, F., Lopes-Lima, M., Pastorino, G., Perez, K. E., Smith, K ... Collen, B. (2021). The conservation status of the world's freshwater molluscs. *Hydrobiologia*, 848(12-13), 3231-3254. <https://doi.org/10.1007/s10750-020-04385-w>
- Ceballos, G., Ehrlich, P. R., & Dirzo, R. (2017). Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines. *Proceedings of the National Academy of Sciences*, 114(30), E6089-E6096. <https://doi.org/10.1073/pnas.1704949114>
- Chapman, A. D. (2009). *Number of living species in Australia and the world* (2nd ed.). Department of the Environment, Water, Heritage and the Arts, Australian Government: Canberra, Australia. <https://www.dcceew.gov.au/science-research/abrs/publications/other/numbers-living-species>
- Cobbinah, J. R., Vink, A., & Onwuka, B. (2008). *Snail farming: Production, processing and marketing*. Agromisa. <https://publications.cta.int/en/publications/publication/1497/post-harvest-processing/index.html>

- Chiew, E. W., Lim, L. H. S., & Ambu, S. (2009). Cercarial dermatitis in Kelantan, Malaysia an occupation related health problem. *International e-Journal of Science, Medicine and Education*, 1(2), 69-73.
- Colley, D. G., Bustinduy, A. L., Secor, W. E., & King, C. H. (2014). Human schistosomiasis. *The Lancet*, 383(9936), 2253-2264. [https://doi.org/10.1016/S0140-6736\(13\)61949-2](https://doi.org/10.1016/S0140-6736(13)61949-2)
- Debbra, M., Premalatha, B., Nurulaini, R., Azima, L., Erwanas, A., & Sohayati, A. R. (2018). Rumen paramphistomosis in *Bos indicus* from a sample received by Veterinary Research Institute of Ipoh, Perak. *Malaysian Journal of Veterinary Research*, 9(2), 154-159. <https://doi.org/doi/pdf/10.5555/20193004793>
- Dida, G. O., Gelder, F. B., Anyona, D. N., Matano, A. S., Abuom, P. O., Adoka, S. O., Ouma, C., Kanangire, C. K., Owuor, P. O., & Ofulla, A. V. O. (2014). Distribution and abundance of schistosomiasis and fascioliasis host snails along the Mara River in Kenya and Tanzania. *Infection Ecology and Epidemiology*, 4(1), 24281. <https://doi.org/10.3402/iee.v4.24281>
- Diyana, J. N. A., Mahiza, M. I. N., Latiffah, H., Fazila, S. H. N., Lokman, I. H., Hazfalinda, H. N., Chandrawathani, P., Ibitoye, E. B., & Juriah, K. (2020). Occurrence, morphometric, and molecular investigation of cattle and buffalo liver adult fluke in Peninsular Malaysia main abattoirs. *Journal of Parasitology Research*, 9-14. <https://doi.org/10.1155/2020/5436846>
- Dumidae, A., Subkrasae, C., Ardpairin, J., Pansri, S., Polseela, R., Thanwisai, A., & Vitta, A. (2024). Population genetic structure of *Indoplanorbis exustus* (Gastropoda: Planorbidae) in Thailand and its infection with trematode cercariae. *PloS One*, 19(1), e0297761. <https://doi.org/10.1371/journal.pone.0297761>
- Dung, B. T., Doanh, P. N., The, D. T., Loan, H. T., Losson, B., & Caron, Y. (2013). Morphological and molecular characterization of lymnaeid snails and their potential role in transmission of *Fasciola* spp. in Vietnam. *Korean Journal of Parasitology*, 51(6), 657-662. <https://doi.org/10.3347/kjp.2013.51.6.657>
- EFSA Panel on Nutrition, Novel Foods and Food Allergens, Turck, D., Bohn, T., Castenmiller, J., De Henauw, S., Hirsch-Ernst, K. I., & Knutsen, H. K. (2024). Safety of *Helix*Complex snail mucus (HSM) as a novel food pursuant to Regulation (EU) 2015/2283. *EFSA Journal*, 22(1), e8492. <https://doi.org/10.2903/j.efsa.2024.8492>
- El-Zawawy, N. A., & Mona, M. M. (2021). Antimicrobial efficacy of Egyptian *Eremina desertorum* and *Helix aspersa* snail mucus with a novel approach to their anti-inflammatory and wound healing potencies. *Scientific Reports*, 11(1), 24317. <https://doi.org/10.1038/s41598-021-03751-8>
- Facon, B., Pointier, J. P., Glaubrecht, M., Poux, C., Jarne, P., & David, P. (2003). A molecular phylogeography approach to biological invasions of the New World by parthenogenetic Thiarid snails. *Molluscan Ecology*, 12, 3027-3039. <https://doi.org/10.1046/j.1365-294X.2003.01972.x>
- Faiz, L. Z. (2020). Diversity and damage assessment of snail in cultivated crops of Neelabut Bagh, Azad Jammu and Kashmir (Pakistan). *Journal of Bioresource Management*, 7(4), 11. <https://doi.org/10.35691/JBM.0202.0149>
- Francis, R. A. (Ed.). (2012). *A handbook of global freshwater invasive species*. Routledge. <https://doi.org/10.4324/9780203127230>
- Garba Djirmay, A., Yadav, R. S., Guo, J., Rollinson, D., & Madsen, H. (2024). Chemical control of snail vectors as an integrated part of a strategy for the elimination of Schistosomiasis review of the state of

- knowledge and future needs. *Tropical medicine and infectious disease*, 9(9), 222. <https://doi.org/10.3390/tropicalmed9090222>
- Greene, S. D. (2008). Extending integrated pest management to the golden apple snail: Examining a community centre approach in northeast Thailand. *International Journal of Pest Management*, 54(2), 95-102. <https://doi.org/10.1080/09670870701621282>
- Greer, G. J., Ow-Yang, C. K., & Yong, H. S. (1988). *Schistosoma malayensis* n.sp.: A *Schistosoma japonicum*-complex schistosome from Peninsular Malaysia. *Journal of Parasitology*, 74(3), 471-480. <https://doi.org/10.2307/3282058>
- Greer, G. J., & Anuar, H. (1984). Serological evidence of schistosomiasis among Orang Asli from three areas of Peninsular Malaysia. *Southeast Asian Journal of Tropical Medicine and Public Health*, 15, 303-12.
- Gugliandolo, E., Cordaro, M., Fusco, R., Peritore, A. F., Siracusa, R., Genovese, T., & Crupi, R. (2021). Protective effect of snail secretion filtrate against ethanol-induced gastric ulcer in mice. *Scientific Reports*, 11(1), 3638. <https://doi.org/10.1038/s41598-021-82921-4>
- Gupta, A., & Khanal, P. (2024). The potential of snails as a source of food and feed. *Journal of Agriculture and Food Research*, 18, 101330. <https://doi.org/10.1016/j.jafr.2024.101330>
- Hurtrez-Boussès, S., Pendino, A., Bernabé, C., Rondelaud, D., Durand, C., Meunier, C., Hurtrez-Boussès, J. E., & Renaud, F. (2005). Comparison between shell morphology and genetic diversity in two sympatric lymnaeid snails, vectors of fasciolosis. *Canadian Journal of Zoology*, 83(11), 1643-1648. <https://doi.org/10.1139/z05-150>
- Hung, N. M., Duc, N. V., Stauffer, J. R., & Madsen, H. (2013). Use of black carp (*Mylopharyngodon piceus*) in biological control of intermediate host snails of fish-borne zoonotic trematodes in nursery ponds in the Red River Delta, Vietnam. *Parasites and Vectors*, 6(1), 142. <https://doi.org/10.1186/1756-3305-6-142>
- Harizt, A. M., Malahubban, M., Syed-Hussain, S. S., Ramanoon, S. Z., Sadiq, M. B., Sarbini, S. R., Hassan, M. N., Mustafa, S., Aziz, N. A. A., Isa, N. M. M., Nating, W. R., Asut, R., & Kamaludeen, J. (2021). Gastrointestinal parasitic infections of buffaloes (*Bubalus bubalis*) in Sarawak Borneo: Prevalence, risk factors, and farming practices. *Tropical Biomedicine*, 38(3), 318-326. <https://doi.org/10.47665/TB.38.3.072>
- Hah, H. E., Liew, T. S., Rama Rao, S., Yow, Y. Y., & Ratnayeke, S. (2022). Distribution and environmental associations of invasive freshwater Pomacea snails in Peninsular Malaysia. *Biological Invasions*, 24(1), 189-204. <https://doi.org/10.1007/s10530-021-02634-0>
- Horgan, F. G., Stuart, A. M., & Kudavidanage, E. P. (2014). Impact of invasive apple snails on the functioning and services of natural and managed wetlands. *Acta Oecologica*, 54, 90-100. <https://doi.org/10.1016/j.actao.2012.10.002>
- International Union for Conservation of Nature and Natural Resources. (2021). *The IUCN Red List of Threatened Species* (Version 2021-2). <https://www.iucnredlist.org>
- International Feed Industry Federation. (2021). *Global feed statistics*. <https://ifif.org/global-feed/statistics>
- Japa, O., Suwanchaoen, C., Bunsong, T., & Phuangsri, C. (2021). Parasitological and molecular characterization of the avian schistosomatid cercariae infecting lymnaeidae snails in Phayao, Northern Thailand. *Veterinary World*, 14(11), 2655-2661. <https://doi.org/10.14202/vetworld.2021.2655-2661>

- Johnson, P. D., Bogan, A. E., Brown, K. M., Burkhead, N. M., Cordeiro, J. R., Garner, J. T., & Strong, E. E. (2013). Conservation status of freshwater gastropods of Canada and the United States. *Fisheries*, 38(6), 247-282. <https://doi.org/10.1080/03632415.2013.785396>
- Jamnah, O., Marliah, A., Norazura, A. H., Maizatul Azlina, A. M., Premaalatha, B., Erwanas, A. I., Ainani, A., Mohamad Bohari, J., Mohd Rostan, A. A., Norhafiza, A. H., Naheed, M., Chandrawathani, P., & Ramlan, M. (2013). Parasites in local buffaloes – incidental findings from mortalities in nili-ravi buffaloes. *Malaysian Journal of Veterinary Research*, 4(2), 47-49.
- Joshi, R. C., Cowie, R. H., & Sebastian, L. S. (Eds.). (2017). Biology and management of invasive apple snails. *Philippine Rice Research Institute (PhilRice)*. 406 pp. https://ri.conicet.gov.ar/bitstream/handle/11336/108894/CONICET_Digital_Nro.6add5657-f918-46ab-b63c-3146fd3c10e8_r.pdf.pdf?sequence=5
- Kamaruddin, N. C., Razali, M. A. I., Busayo, I. E., Hamzah, N. H., Idris, L. H., & Md Isa, N. M. (2021). The first report of ruminant fascioliasis in Sabah, East Malaysia. *Journal of Parasitology Research*, 1(2021), 6691483. <https://doi.org/10.1155/2021/6691483>
- Kim, J. S. (1978). An ecological study of Paragonimus in Malaysia. *Kisaengch'unghak chapchi. The Korean Journal of Parasitology*, 16(1), 47-53. <https://doi.org/10.3347/kjp.1978.16.1.47>
- Krailas, D., Namchote, S., Koonchornboon, T., Dechruksa, W., & Boonmekam, D. (2014). Trematodes obtained from the thiarid freshwater snail *Melanooides tuberculata* (Müller, 1774) as vector of human infections in Thailand. *Zoosystematics and Evolution*, 90, 57. <https://doi.org/10.3897/zse.90.7306>
- Khadijah, S., Ariff, Z., Nurlaili, M. R., Sakiinah, A., Izzudin, A. H., Mursyidah, A. K., Rita, N., & Nur Aida, H. (2017). Fasciola and Paramphistomum infection in large ruminants. *International Journal of Agronomy and Agricultural Research International*, 10, 19-26.
- King, C. H., Sutherland, L. J., & Bertsch, D. (2015). Systematic review and meta-analysis of the impact of chemical-based mollusciciding for control of *Schistosoma mansoni* and *S. haematobium* transmission. *PLoS Neglected Tropical Diseases*, 9(12), e0004290. <https://doi.org/10.1371/journal.pntd.0004290>
- Kumar, P. (2020). A review—on molluscs as an agricultural pest and their control. *International Journal of Food Science and Agriculture*, 4(4), 383-389. <https://doi.org/10.26855/ijfsa.2020.12.013>
- Lee, J. H., Quan, J. H., Choi, I. W., Park, G. M., Cha, G. H., Kim, H. J., & Lee, Y. H. (2017). Fasciola hepatica: Infection status of freshwater snails collected from Gangwon-do (Province), Korea. *The Korean Journal of Parasitology*, 55(1), 95-100. <https://doi.org/10.3347/kjp.2017.55.1.95>
- Leighton, B. J., Zervos, S., & Webster, J. M. (2000). Ecological factors in schistosome transmission, and an environmentally benign method for controlling snails in a recreational lake with a record of schistosome dermatitis. *Parasitology International*, 49(1), 9–17. [https://doi.org/10.1016/S1383-5769\(00\)00035-7](https://doi.org/10.1016/S1383-5769(00)00035-7)
- Li, Z. J., Ge, J., Dai, J. R., Wen, L. Y., Lin, D. D., & Madsen, H., et al. (2016). Biology and control of snail intermediate host of *Schistosoma japonicum* in the People's Republic of China. *Advances in Parasitology*, 92, 197–236. <https://doi.org/10.1016/bs.apar>
- Latif, B., Heo, C. C., Razuin, R., Shamalaa, D. V., & Tappe, D. (2013). Autochthonous human schistosomiasis, Malaysia. *Emerging Infectious Diseases*, 19(8), 1340-1341.

- Lee, H. (1965). *Paragonimus westermani* infection in wild mammals and crustacean host in Malaysia. *The American Journal of Tropical Medicine and Hygiene*, 14(4), 581.
- Lopes, A. S., Pulido-Murillo, E. A., López-Hernández, D., Melo, A. L. de, & Pinto, H. A. (2021). First report of *Melanoides tuberculata* (Mollusca: Thiaridae) harboring a xiphidiocercaria in Brazil: A new parasite introduced in the Americas? *Parasitology International*, 82(October 2020). <https://doi.org/10.1016/j.parint.2021.102284>
- Lu, X. T., Gu, Q. Y., Limpanont, Y., Song, L. G., Wu, Z. D., Okanurak, K., & Lv, Z. Y. (2018). Snail-borne parasitic diseases: An update on global epidemiological distribution, transmission interruption and control methods. *Infectious Diseases of Poverty*, 7(1), 1-16. <https://doi.org/10.1186/s40249-018-0414-7>
- Maes, T., Hammoud, C., Volckaert, F. A., & Huyse, T. (2021). A call for standardised snail ecological studies to support schistosomiasis risk assessment and snail control efforts. *Hydrobiologia*, 848(8), 1773-1793. <https://doi.org/10.1007/s10750-021-04547-4>
- Madsen, H., & Hung, N. M. (2014). An overview of freshwater snails in Asia with main focus on Vietnam. *Acta Tropica*, 140, 105-117. <https://doi.org/10.1016/j.actatropica.2014.08.005>
- Martin, G. L., & Cabrera, E. C. (2018). Morphological characterization of emerging cercariae among lymnaeid snails from Barangay Cawongan, Padre Garcia, Batangas, Philippines. *Journal of Parasitology Research*, 2018(1), 1-13. <https://doi.org/10.1155/2018/5241217>
- Maciaszek, R., Marcinek, D., & Sosnowski, W. (2019). Tropical snail, the red-rimmed melania *Melanoides tuberculata* Müller, 1774 (Thiaridae) from artificial lake in Warsaw, Central Poland. *World Science News*, 122, 285-290.
- Mirfendereski, R., Hashemi, S., Shirali, S., Shemshadi, B., & Lawton, S. P. (2021). DNA barcoding of Iranian radicine freshwater snails begins to untangle the taxonomy and phylogeography of intermediate hosts of schistosomiasis and fasciolosis from the Middle East and across Central Asia. *Infection, Genetics and Evolution*, 89, 104728. <https://doi.org/10.1016/j.meegid.2021.104728>
- Naresh, G., Gomez, P. A., Salmah, B., & Suryati, M. Y. (2006). Fasciolosis (liver fluke) of the breast in a male patient: A case report. *Breast*, 15(1), 103-105. <https://doi.org/10.1016/j.breast.2005.02.009>
- Ng, T. H., Dulipat, J., Foon, J. K., Lopes-Lima, M., Zieritz, A., & Liew, T. S. (2017). A preliminary checklist of the freshwater snails of Sabah (Malaysian Borneo) deposited in the BORNEENSIS collection, Universiti Malaysia Sabah. *ZooKeys*, 673, 105-123. <https://doi.org/10.3897/zookeys.673.12544>
- Nguyen, H. M., Greiman, S. E., Van Hoang, H., Ngoc, C. N., Van Nguyen, H., Pham, T. C., & Madsen, H. (2022). Morphological and molecular identification of trematode cercariae related with humans and animal health in freshwater snails from a lake and a dam in Myanmar. *The Journal of Parasitology*, 108(6), 601-612. <https://doi.org/10.1645/21-125>
- Noothuan, N., Apitanyasai, K., Panha, S., & Sutcharit, C. (2021). Snail mucus from the mantle and foot of two land snails, *Lissachatina fulica* and *Hemiplecta distincta*, exhibits different protein profile and biological activity. *BMC Research Notes*, 14, 138. <https://doi.org/10.1186/s13104-021-05557-0>
- Pathak, C. R., Luitel, H., Utaaker, K. S., & Khanal, P. (2024). One-health approach on the future application of snails: A focus on snail-transmitted parasitic diseases. *Parasitology Research*, 123(1), 28. <https://doi.org/10.1007/s00436-023-07975-y>

- Phiri, A. M., Phiri, I. K., Chota, A., & Monrad, J. (2007). Trematode infections in freshwater snails and cattle from the Kafue wetlands of Zambia during a period of highest cattle-water contact. *Journal of Helminthology*, 81(1), 85-92. <https://doi.org/10.1017/S0022149X07387786>
- Phoong, M. J. C., Hah, H. E., Rao, S. R., Yow, Y. Y., & Ratnayeke, S. (2018). Invasive apple snails in wetlands of Selangor, Malaysia: Species, Distribution, and ecological associations. *Journal of Tropical Biology and Conservation (JTBC)*, 15, 43-60. <https://doi.org/10.51200/jtbc.v15i0.1474>
- Piechocki, A., Wawrzyniak-Wydrowska, B., & Zdanowski, B. (2003). *Melanoides tuberculatus* (O. F. Müller, 1774) (Orthogastropoda: Thiaridae), a gastropod species new for the fauna of Poland. *Folia Malacologica*, 11, 39-41. <https://doi.org/10.12657/folmal.011.004>
- Post, J. M., Reasch, R. J., & Bailey, E. S. (2022). Detection of trematodes from the host exotic aquatic snail *Melanoides tuberculata* in an urban stormwater system. *Zoonotic Diseases*, 2(4), 258-266. <https://doi.org/10.3390/zoonoticdis2040021>
- Prasopdee, S., Tesana, S., Cantacessi, C., Laha, T., Mulvenna, J., Grams, R., Loukas, A., & Sotillo, J. (2015). Data set from the proteomic analysis of *Bithynia siamensis* goniomphalos snails upon infection with the carcinogenic liver fluke *Opisthorchis viverrini*. *Data in Brief*, 2, 16-20. <https://doi.org/10.1016/j.dib.2014.09.005>
- Rao, S. R., Liew, T. S., Yow, Y. Y., & Ratnayeke, S. (2018). Cryptic diversity: Two morphologically similar species of invasive apple snail in Peninsular Malaysia. *PLoS ONE*, 13(5), 1-16. <https://doi.org/10.1371/journal.pone.0196582>
- Rita, N., Mursyidah, A. K., & Khadijah, S. (2017). The prevalence of helminthiasis in cattle, Terengganu, Peninsular Malaysia. *Tropical Biomedicine*, 34(2), 324-331.
- Rosilawati, K., Ramli S., & Saipul, B. A. R. (2017). Fascioliasis in an adult draught buffalo in Malaysia: A case report. *Malaysian Journal of Veterinary Research*, 8(1), 169-172.
- Saad, K., Mohammad Faiz, M., Nadarasan, R., Mursyidah, A. K., & Hassan, Q. N. (2019). High prevalence fluke infection at four cattle farms located in Kuala Terengganu, Malaysia. *Tropical Life Sciences Research*, 30(3), 59-67. <https://doi.org/10.21315/tlsr2019.30.3.4>
- Saijuntha, W., Andrews, R. H., Sithithaworn, P., & Petney, T. (2021). *Biodiversity of Southeast Asian parasites and vectors causing human disease*. In T. N. Petney, W. Saijuntha, H. Mehlorn (Eds.), *Parasitology research monographs*. Springer. https://doi.org/10.1007/978-3-030-71161-0_1
- Sodhi, N. S., Brook, B. W., & Bradshaw, C. J. (2009). Causes and consequences of species extinctions. *The Princeton Guide to Ecology*, 1(1), 514-520.
- Sokolow, S. H., Huttinger, E., Jouanard, N., Hsieh, M. H., Lafferty, K. D., Kuris, A. M., Riveau, G., Senghor, S., Thiam, C., N'Diaye, A., Faye, D. S., & De Leo, G. A. (2015). Reduced transmission of human schistosomiasis after restoration of a native river prawn that preys on the snail intermediate host. *Proceedings of the National Academy of Sciences of the United States of America*, 112(31), 9650–9655. <https://doi.org/10.1073/pnas.1502651112>
- Sokolow, S. H., Lafferty, K. D., & Kuris, A. M. (2014). Regulation of laboratory populations of snails (*Biomphalaria* and *Bulinus* spp.) by river prawns, *Macrobrachium* spp. (Decapoda, Palaemonidae):

- Implications for control of schistosomiasis. *Acta Tropica*, 132, 64–74. <https://doi.org/10.1016/j.actatropica.2013.12.021>
- Strayer, D.L., & Dudgeon, D. (2010). Freshwater biodiversity conservation: recent progress and future challenges. *Journal of The North American Benthological Society*, 29(1), 344-359. <https://doi.org/10.1899/08-171.1>
- Thakuri, B., Acharya, B. K., & Sharma, G. (2019). Population density and damage of invasive giant African snail *Achatina fulica* in an organic farm in East Sikkim, India. *Indian Journal of Ecology*, 46(3), 631-635.
- Tookhy, N. A., Isa, N. M. M., Mansor, R., Rahaman, Y. A., Ahmad, N. I., Bui, D. T., Idris, L. H., Hamzah, N. H., & Zulkifli, N. (2023). Morphological and molecular identification of lymnaeid snail and trematodes cercariae in different water bodies in Perak, Malaysia. *Parasitology Research*, 122(7), 1475-1488. <https://doi.org/10.1007/s00436-023-07845-z>
- Tookhy, N. A., Mahiza, M. I. N., Mansor, R., Yasmin, A. R., Ahmad, N. I., Hamzah, N. H., & Idris, L. H. (2022). Rumen Fluke in Cattle and Buffaloes in Asia: A Review. *Pertanika Journal of Tropical Agricultural Science*, 45(3), 781-803. <https://doi.org/10.47836/pjtas.45.3.15>
- Tookhy, N. A., Isa, N-M. M., Rahaman, Y. A., Ahmad, N. I., Sharma, R. S. K., Idris, L. H., Mansor, R., Bui, D. T., Hamzah, N. H., & Shakhes, S. A. (2024). Epidemiology of rumen fluke infection in selected buffalo farms in Perak, Malaysia: Prevalence, molecular species identification, and associated risk factors. *Parasitology Research*, 123, 199. <https://doi.org/10.1007/s00436-024-08219-9>
- Vinarski, M., Bolotov, I. N., Aksenova, O. V., Babushkin, E. S., Bespalaya, Y. V., et al. (2020). Freshwater Mollusca of the Circumpolar Arctic: A review on their taxonomy, diversity, and biogeography. *Hydrobiologia*, 1–28. <https://doi.org/10.1007/s10750-020-04345-5>
- World Health Organization. (2016). *Schistosomiasis: WHO reports substantial treatment progress for school age children*. <https://www.who.int/news/item/08-12-2017-schistosomiasis-who-reports-substantial-treatment-progress-for-school-age-children>
- World Health Organization. (2018). *Human fascioliasis: Review provides fresh perspectives on infection and control*. <https://www.who.int/news/item/26-07-2018-human-fascioliasis-review-provides-fresh-perspectives-on-infection-and-control>
- World Health Organization. (2020). *Neglected tropical diseases*. <https://www.who.int/news-room/questions-and-answers/item/q-a-on-fascioliasis>
- Xia, J., Yuan, Y., Xu, X., Wei, F., Li, G., Liu, M., ... & Nie, S. (2014). Evaluating the effect of a novel molluscicide in the endemic schistosomiasis japonica area of China. *International Journal of Environmental Research and Public Health*, 11(10), 10406-10418.
- Yahaya, H., Amzah, B., Sivapragasam, A., Nordin, M., Muhamad Hisham, M. N., & Misrudin, H. (2017). Invasive apple snails in Malaysia. *Biology and Management of Invasive Apple Snails*, 1, 169195.
- Yi, X. L., Liu, J., Cao, M. L., Xiong, J., Deng, Y. P., Wang, H. M., Ma, P. P., Liu, G. H., & Yang, H. (2024). Population genetics and genetic variation of *Pomacea canaliculata* (Gastropoda: Ampullariidae) in China revealed by sequence analyses of three mitochondrial genes. *Ecology and evolution*, 14(1), e10836. <https://doi.org/10.1002/ece3.10836>

- Yusa, Y., Sugiura, N., & Wada, T. (2006). Predatory potential of freshwater animals on an invasive agricultural pest, the apple snail *Pomacea canaliculata* (Gastropoda: Ampullariidae), in southern Japan. *Biological Invasions*, 8, 137–147. <https://doi.org/10.1007/s10530-004-1790-4>
- Younes, A., El-Sherief, H., Gawish, F., & Mahmoud, M. (2017). Biological control of snail hosts transmitting schistosomiasis by the water bug, *Sphaerodema urinator*. *Parasitology Research*, 116(4), 1257–1264. <https://doi.org/10.1007/s00436-017-5402-5>