

Comparison of PCR fingerprinting techniques for the discrimination of *Salmonella enterica* subsp. *enterica* serovar Weltevreden isolated from indigenous vegetables in Malaysia.

ABSTRACT

Salmonella enterica subsp. *enterica* (S.) serovar Weltevreden has emerged as a public health problem in many countries. Genomic DNA of *S. Weltevreden* from indigenous vegetables namely 'selom' (*Oenanthe stolonifera*), 'pegaga' (*Centella asiatica*), 'kesum' (*Polygonum minus*) and 'kangkong' (*Ipomoea aquatica*) were characterized by duplex-polymerase chain reaction (duplex-PCR), multiplex-polymerase chain reaction (multiplex-PCR), random amplified polymorphic DNA (RAPD), enterobacterial repetitive intergenic consensus-polymerase chain reaction (ERIC-PCR) and polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP). The results demonstrated that a total of four clusters and three single isolates were generated from ERIC-PCR with primers ERIC-1 and ERIC-2 whereas RAPD with arbitrary primers OPAR2, OPAR17 and OPAR19 discriminated the *S. Weltevreden* into nine clusters and eight single isolates at a common 65% similarity level with discriminatory index (D) of 0.7443 and 0.9394 respectively. Composite analysis of banding profiles generated from RAPD-PCR and ERIC-PCR showed eight clusters and six single isolates at 65% similarity level with the highest D value that is 0.9508. On the other hand, PCR-RFLP and duplex PCR data exhibited a consistent profile for *S. Weltevreden*. Multiplex-PCR targeting three different antibiotic resistance genes and a common *Salmonella* specific gene segment produced two distinguishing profiles among the *S. Weltevreden* examined. These results demonstrated that the combined analysis of RAPD-PCR and ERIC-PCR is a better tool for characterizing *S. Weltevreden* than individual methods.

Keyword: Duplex-PCR; ERIC-PCR; Indigenous vegetables; Multiplex-PCR; PCR-RFLP; RAPD; *S. Weltevreden*.