

Development of an inhibitive enzyme assay for copper

ABSTRACT

In this work the development of an inhibitive assay for copper using the molybdenum-reducing enzyme assay is presented. The enzyme is assayed using 12-molybdophosphoric acid at pH 5.0 as an electron acceptor substrate and NADH as the electron donor substrate. The enzyme converts the yellowish solution into a deep blue solution. The assay is based on the ability of copper to inhibit the molybdenum-reducing enzyme from the molybdate-reducing *Serratia* sp. Strain DRY5. Other heavy metals tested did not inhibit the enzyme at 10 mg l⁻¹. The best model with high regression coefficient to measure copper inhibition is one-phase binding. The calculated IC₅₀ (concentration causing 50% inhibition) is 0.099 mg l⁻¹ and the regression coefficient is 0.98. The comparative LC₅₀, EC₅₀ and IC₅₀ data for copper in different toxicity tests show that the IC₅₀ value for copper in this study is lower than those for immobilized urease, bromelain, Rainbow trout, *R. meliloti*, Baker's Yeast dehydrogenase activity *Spirillum volutans*, *P. fluorescens*, *Aeromonas hydrophilia* and synthetic activated sludge assays. However the IC₅₀ value is higher than those for *Ulva pertusa* and papain assays, but within the reported range for *Daphnia magna* and Microtox assays.

Keyword: Inhibitive enzyme assay; Copper; Mo-reducing enzyme