



UNIVERSITI PUTRA MALAYSIA

**SECONDARY METABOLITES FROM STEM BARK OF
GERONGGANG (*CRATOXYLUM ARBORESCENS* (VAHL) BL.) AND
KETEMAU (*CRATOXYLUM GLAUCUM* K.) AND THEIR BIOLOGICAL
ACTIVITIES**

SIM WEI CHUNG

FS 2010 27



**SECONDARY METABOLITES FROM STEM BARK OF GERONGGANG
(*CRATOXYLUM ARBORESCENS* (VAHL) BL.) AND KETEMAU
(*CRATOXYLUM GLAUCUM* K.) AND THEIR BIOLOGICAL ACTIVITIES**

By

SIM WEI CHUNG

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

September 2010



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Science

**SECONDARY METABOLITES FROM STEM BARK OF GERONGGANG
(*CRATOXYLUM ARBORESCENS* (VAHL) BL.) AND KETEMAU
(*CRATOXYLUM GLAUCUM* K.) AND THEIR BIOLOGICAL ACTIVITIES**

By

SIM WEI CHUNG

September 2010

Chairman : Prof. Gwendoline Ee Cheng Lian, PhD

Faculty : Science

Phytochemical and biological studies were carried out on two species from the Guttiferae family, *Cratoxylum arborescens* and *Cratoxylum glaucum*. The chemical investigation discovered triterpenoids, quinones and xanthones. These compounds were isolated using common chromatographic techniques and were identified using modern spectroscopic techniques such as 1D and 2D NMR, MS, IR and UV.

Cratoxylum arborescens (stem bark) afforded friedelin, stigmasterol, fuscaxanthone C and two anthraquinone, vismiaquinone and 3-geranyloxy-6-methyl-1,8-dihydroxyanthraquinone. Meanwhile, studies on *Cratoxylum glaucum* (stem bark)



revealed the presence of three xanthones, α -mangostin, β -mangostin and fuscaxanthone C, along with an anthraquinone, 1,8-dihydroxy-3-methoxy-6-methylanthraquinone, friedelin, β -sitosterol and betulinic acid. This is the first report of α -mangostin, β -mangostin and betulinic acid from *Cratoxylum glaucum*. The acetylation of fuscaxanthone C has successfully yielded dimethylmangostin acetate.

Antioxidant properties of the crude extracts of these two species were tested using 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging method and compared to a well known antioxidant, ascorbic acid. The crude methanol extracts of both *Cratoxylum arborescens* and *C. glaucum* showed high antioxidant activities with low EC₅₀ values of 10.40 and 7.48 ppm, respectively, and are comparable to that of ascorbic acid, 5.17 ppm. The crude chloroform extract of *C. arborescens* also gave good activity, with an EC₅₀ value of 31.04 ppm. The total phenolic content of the methanol extracts of both species were measured using Folin-Ciocalteu method.

The hexane and ethyl acetate extracts of *Cratoxylum glaucum* and the hexane and chloroform extracts of *Cratoxylum arborescens* demonstrated low IC₅₀ values against the MCF7 cancer cell line. Meanwhile, the ethyl acetate and methanol extracts of *Cratoxylum glaucum* and all the extracts of *Cratoxylum arborescens* were active against HL-60 cancer cell line.



The antibacterial activity test was also carried out using eight bacteria, namely, *Bacillus cereus*, *Micrococcus luteus*, Methicillin-Sensitive *Staphylococcus Aureus* (MSSA), Methicillin-resistant *Staphylococcus aureus* (MRSA), *Escherichia coli*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa* and *Enterobacter aerogenes*. However, all the crude extracts gave only weak or no inhibition to these bacteria.

The larvicidal tests were performed against the larvae of *Aedes aegypti* using the WHO (1981) standard procedures with slight modifications. All the extracts of *Cratoxylum arborescens* and *C. glaucum* were found to be inactive against the larvae of *Aedes aegypti*.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**METABOLIT SEKUNDER DARIPADA KULIT GERONGGANG
(*CRATOXYLUM ARBORESCENS* (VAHL) BL.) DAN KETEMAU
(*CRATOXYLUM GLAUCUM* K.) DAN AKTIVITI BIOLOGI**

Oleh

SIM WEI CHUNG

September 2010

Pengerusi : Prof. Gwendoline Ee Cheng Lian, PhD

Fakulti : Sains

Kajian fitokimia dan biologi telah dilakukan terhadap dua spesies daripada famili Guttiferae, iaitu *Cratoxylum arborescens* dan *Cratoxylum glaucum*. Penyelidikan kimia ini telah menemukan triterpenoid, kuinon dan xanton. Sebatian ini diasingkan dengan menggunakan teknik kromatografi biasa dan dikenal pasti dengan menggunakan teknik spektroskopi seperti 1D dan 2D NMR, MS, IR dan UV.

Daripada *Cratoxylum arborescens* (kulit batang), diperolehi friedelin, stigmasterol, fuskaxanton C dan dua antrakuinon, vismiakuinon dan 3-geraniloksi-6-metil-

1,8-dihidroksiantrakuinon. Manakala, kajian terhadap *Cratoxylum glaucum* (kulit batang) telah menemukan tiga xanton, α -mangostin, β -mangostin dan fuskaxanton C, bersama dengan satu antrakuinon, 1,8-dihidroksi-3-metoksi-6-metilantrakuinon, friedelin, β -sitosterol and asid betulinik. Ini merupakan laporan pertama bagi α -mangostin, β -mangostin dan asid betulinik daripada *Cratoxylum glaucum*. Asetilasi fuskaxanton C telah berjaya menghasilkan dimetilmangostin asetat.

Keupayaan antipengoksidaan ekstrak dua spesis ini telah diuji dengan menggunakan cara penghapusan radikal 2,2-difenil-1-pikrilhidrazil (DPPH) dan dibandingkan dengan agen antipengoksidaan yang umum, iaitu asid askorbik. Ekstrak metanol bagi kedua-dua *Cratoxylum arborescens* dan *C. glaucum* menunjukkan aktiviti anti-pengoksidaan yang tinggi dengan EC₅₀ yang rendah, iaitu 10.40 dan 7.48 ppm, berbanding dengan asid askorbik, iaitu 5.17 ppm. Ekstrak kloroform *C. arborescens* juga menunjukkan aktiviti yang baik dengan nilai EC₅₀ pada 31.04 ppm. Jumlah kandungan fenolik bagi ekstrak methanol bagi kedua-dua spesis telah disukat dengan menggunakan kaedah Folin-Ciocalteu.

Ekstrak heksana dan etil asetat *Cratoxylum glaucum* dan ekstrak heksana dan kloroform *Cratoxylum arborescens* menunjukkan nilai IC₅₀ yang rendah terhadap sel barah MCF7. Manakala, ekstrak etil asetat dan metanol daripada *Cratoxylum glaucum* dan semua ekstrak daripada *Cratoxylum arborescens* adalah aktif terhadap sel barah HL-60.

Ujian aktiviti antibakteria dikaji terhadap lapan jenis bakteria, iaitu *Bacillus cereus*, *Micrococcus luteus*, Methicillin-Sensitive *Staphylococcus aureus* (MSSA), Methicillin-resistant *Staphylococcus aureus* (MRSA), *Escherichia coli*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa* dan *Enterobacter aerogenes*. Akan tetapi, kesemua ekstrak tidak menunjukkan aktiviti ataupun hanya menunjukkan aktiviti lemah terhadap bakteria tersebut.

Ujian ke atas larva telah dilakukan dengan larva *Aedes aegypti* dengan mengikuti prosedur piawai WHO (1981) dengan sedikit pengubahsuaihan. Kesemua ekstrak *Cratoxylum arborescens* dan *C. glaucum* didapati tidak aktif terhadap larva *Aedes aegypti*.



ACKNOWLEDGEMENTS

It is a pleasure for me to express my greatest gratitude to the following groups and individuals who made this thesis possible. First of all, I would also like to express my greatest gratitude to Prof. Dr Gwendoline Ee Cheng Lian for her dedicated supervision. She was always available for consultation with invaluable comments and constructive suggestions in conducting my project. I would like to thank my co-supervisor Prof. Dr Mohd. Aspollah Hj. Sukari who provided me a lot of help in my research. Heartiest thanks are extended to all the laboratory assistants and science officers in Chemistry Department for assisting me in characterising the natural products I have successfully isolated. Especially, Mr. Johadi Iskandar for NMR analysis, Mr. Zainal Abidin Kassim for recording the mass spectrum and Mrs. Rusnani Amirudin for IR analysis.

A deepest acknowledgment is extended to Lim Chyi Meei, Wen Yin Ping and Yaw Chong Wey for their willingness to help, listen and share my problems over the past two years. Without their assistance and help, this thesis would not materialise. And thanks also to all my friends and juniors, who would be pages long for me to list down here, but have filled my life with colours and joy.

Last but most importantly, I would like to express my deepest gratitude and my appreciation to my family members, especially my parents, who have supported me emotionally, which means a tremendous lot to me.



I certify that a Thesis Examination Committee has met on 21 September 2010 to conduct the final examination of Sim Wei Chung on his thesis entitled “Secondary Metabolites from the Stem Bark of Geronggang (*Cratoxylum arborescens* (Vahl) Bl.) and Ketemau (*Cratoxylum glaucum* K.) and Their Biological Activities” in accordance with the Universities and University College Act 1971 and the Constitution of Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the candidate be awarded the Master of Science.

Members of the Examination Committee are as follows:

Intan Safinar Bt Ismail, PhD

Lecturer

Faculty of Science

Universiti Putra Malaysia

(Chairman)

Siti Mariam Bt Md. Noor

Lecturer

Faculty of Science

Universiti Putra Malaysia

(Internal Examiner)

Mawardi Rahmani

Professor

Faculty of Science

Universiti Putra Malaysia

(Internal Examiner)

Nor Hadiani Ismail, PhD

Professor

Universiti Teknologi Mara

(External Examiner)

SHAMSUDDIN SULAIMAN, PhD

Professor and Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 26 November 2010



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Gwendoline Ee Cheng Lian, PhD

Professor

Faculty of Science

Universiti Putra Malaysia

(Chairman)

Mohd. Aspollah Hj. Sukari, PhD

Professor

Faculty of Science

Universiti Putra Malaysia

(Member)

HASAMAH MOHD GHAZALI, PhD

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: December 2010



DECLARATION

I declare that the thesis is my original work except for quotation and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently submitted, for any degree at Universiti Putra Malaysia or other institutions.

SIM WEI CHUNG

Date: 21 September 2010



TABLE OF CONTENT

	Page
ABSTRACT	ii
ABSTRAK	v
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xv
LIST OF FIGURES	xvii
LIST OF ABBREVIATIONS	xxv
 CHAPTER	
1 INTRODUCTION	1
1.1 The Value of Natural Products	1
1.2 Botanical Information of the Genus <i>Cratoxylum</i> Blume	5
1.3 <i>Cratoxylum arborescens</i> (Vahl) Blume	6
1.4 <i>Cratoxylum glaucum</i> Korth	7
1.5 Objectives of Study	9
2 LITERATURE REVIEW	10
2.1 <i>Cratoxylum arborescens</i> (Vahl) Blume & <i>Cratoxylum glaucum</i> Korth	10
2.1.1 Xanthones	10
2.1.2 Quinones	11
2.1.3 Flavonoids	12
2.2 The Chemistry of <i>Cratoxylum</i>	13
3 EXPERIMENTAL	41
3.1 Plant Material	41
3.2 Instruments	41
3.2.1 Infrared Spectroscopy (IR)	41
3.2.2 Mass Spectrometry (MS)	41
3.2.3 Melting Point	41
3.2.4 Nuclear Magnetic Resonance (NMR)	42
3.2.5 Ultra Violet Spectroscopy (UV)	42
3.3 Chromatographic Methods	42
3.3.1 Column Chromatography	42
3.3.2 Thin Layer Chromatography (TLC)	43
3.4 Dyeing Reagents for TLC	43
3.4.1 Vanillin-Sulphuric Acid Solution	43



3.4.2 Iron(III) Chloride Solution	44
3.4.3 2,2-diphenyl-1-picrylhydrazyl (DPPH)	44
3.5 Extraction and Isolation of Natural Products from <i>Cratoxylum arborescens</i> and <i>Cratoxylum glaucum</i> .	45
3.5.1 <i>Cratoxylum arborescens</i>	46
3.5.2 <i>Cratoxylum glaucum</i>	47
3.6 Natural Products from <i>Cratoxylum arborescens</i> and <i>Cratoxylum glaucum</i>	48
3.6.1 Isolation of Friedelin (2)	48
3.6.2 Isolation of Stigmasterol (102)	49
3.6.3 Isolation of Fuscaxanthone C (44)	50
3.6.4 Isolation of Vismiaquinone (51)	52
3.6.5 Isolation of 3-geranyloxy-6-methyl-1,8-dihydroxy anthraquinone (30)	53
3.6.6 Isolation of β -sitosterol (115)	54
3.6.7 Isolation of β -mangostin (4)	56
3.6.8 Isolation of 1,8-dihydroxy-3-methoxy-6-methyl anthraquinone (101)	57
3.6.9 Isolation of Betulinic Acid (42)	59
3.6.10 Isolation of α -mangostin (3)	60
3.7 Synthesis of Dimethylmangostin Acetate (116) from Fuscaxanthone C (44)	62
3.8 Bioassay	64
3.8.1 Larvicidal Assay	64
3.8.2 Cytotoxic Assay	65
3.8.3 Antibacterial Activity	66
3.8.4 2,2-diphenyl-1-picrylhydrazyl (DPPH) Radical Scavenging Assay	67
3.8.5 Total Phenolic Content (TPC)	67
4 RESULTS AND DISCUSSION	68
4.1 Isolation of Chemical Constituents from <i>Cratoxylum arborescens</i>	68
4.2 Isolation of Chemical Constituents from <i>Cratoxylum glaucum</i>	70
4.3 Natural Products from <i>Cratoxylum arborescens</i> and <i>Cratoxylum glaucum</i>	72
4.3.1 Characterisation of Friedelin (2)	72
4.3.2 Characterisation of Stigmasterol (102)	80
4.3.3 Characterisation of Fuscaxanthone C (44)	85



4.3.4 Characterisation of Vismiaquinone (51)	108
4.3.5 Characterisation of 3-geranyloxy-6-methyl-1,8-dihydroxyanthraquinone (30)	126
4.3.6 Characterisation of β -sitosterol (115)	144
4.3.7 Characterisation of β -mangostin (4)	150
4.3.8 Characterisation of 1,8-dihydroxy-3-methoxy-6-methylantraquinone (101)	163
4.3.9 Characterisation of Betulinic Acid (42)	176
4.3.10 Characterisation of α -mangostin (3)	184
4.4 Acetylation of Fuscaxanthone C	200
4.4.1 Characterisation of Dimethylmangostin Acetate (116)	201
4.5 Bioassay Analysis	215
4.5.1 Larvicidal Assay	215
4.5.2 2,2-diphenyl-1-picrylhydrazyl (DPPH) Radical Scavenging Assay	215
4.5.3 Total Phenolic Content (TPC)	217
4.5.4 Cytotoxic Assay	217
4.5.5 Antibacterial Assay	221
5 CONCLUSIONS	222
BIBLIOGRAPHY	225
APPENDIX	230
BIODATA OF STUDENT	231

