

# **UNIVERSITI PUTRA MALAYSIA**

# DIVERSITY OF MIGRATORY SHOREBIRDS AND THEIR HABITAT CHARACTERISTICS IN KUALA GULA BIRD SANCTUARY, PERAK, MALAYSIA

**ROMEO M. LOMOLJO** 

# DIVERSITY OF MIGRATORY SHOREBIRDS AND THEIR HABITAT CHARACTERISTICS IN KUALA GULA BIRD SANCTUARY, PERAK, MALAYSIA

By

ROMEO M. LOMOLJO

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia in Fulfilment of the Requirements for the Degree of Doctor of Philosophy

January 2011

## **DEDICATION**

To my daughters En-En, Barbie and my wife Evangeline for the love and understanding

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

# DIVERSITY OF MIGRATORY SHOREBIRDS AND THEIR HABITAT CHARACTERISTICS IN KUALA GULA BIRD SANCTUARY, PERAK, MALAYSIA

By

#### **ROMEO LOMOLJO**

January 2011

Chairman: Prof. Dr. Ahmad Ismail, PhD

**Faculty: Science** 

The research study was conducted in one of the primary corridors and stopover sites of the migratory shorebirds during annual migrations in Kuala Gula Bird Sanctuary, Perak in the west coast of Peninsular Malaysia. The objective were to determine the diversity of migratory shorebirds and to assess the habitat quality, prey availability, environmental conditions, heavy metal concentrations in the sediments, macrobenthic organisms, fish and shrimps from October 2006-September 2007. Two sampling stations each covering an area of 100 x 100 m, were established to represent the whole Kuala Gula Bird Sanctuary between 04°55.185'N, 100°27.8'E and 100°27.761'E. Survey of shorebirds was carried out through direct count and identification to species level. Sixty-one migratory shorebird species from 12 families were identified in both stations. The overall

total of shorebirds density was found to be higher in northward migration (7913 individual/ha) than southward migration (4396 individual/ha). Similarly, the species diversity index (H' = 3.20; p<0.05) and the species richness (d' = 6.19; p<0.05) were also higher in northward migration. However, no significant (p>0.05) differences were found in species evenness, between southward and northward migration throughout the study period. Meanwhile, 28 species of macrobenthos in both stations were recorded and identified representing 15 families throughout the study period. The species diversity index of macrobenthos were higher (p<0.05) in station 2 (H'= 3.24) compared to station 1 (H'= 3.13). Similarly, the species richness was higher (p<0.05) in station 2 (d'= 6.22) compared to station 1 (d' = 6.11). Among the macrobenthos, polychaetes were found to be the most abundant (204.3  $\pm$  6.84 individual/m<sup>2</sup>), followed by molluscs (156.8  $\pm$  6.3 individual/m<sup>2</sup>), fishes (132.8  $\pm$  6.6 individual/m<sup>3</sup>), and shrimps (135.3  $\pm$  4.1 individual/m<sup>3</sup>). The overall total macrobenthos density and biomass revealed a positive (r = 0.920; p<0.05) correlation with bird density, suggesting the importance of food resources to the distribution of migratory shorebirds in the area.

The environmental quality in the area in terms of water quality and nutrients were maintained in healthy levels. No significant variations in the *in situ* water quality parameters (salinity, pH, temperature and dissolved oxygen) were observed in both sampling stations throughout the study period. In addition, nitrate nitrogen (NO<sub>3</sub>-N), total ammonia nitrogen (TAN) and soluble reactive phosphorus (SRP) in the water surface were all in safe levels and maintained at minimum (SRP =  $55.92 \pm 7.88 \,\mu g/L$ ; nitrate-N =  $85.68 \pm 24.33 \,\mu g/L$ ; TAN =  $85.91 \pm 6.54 \,\mu g/L$ ) throughout the study period. The total

geochemical fractionation analysis of sediments in Station 1 and Station 2 ranged from 80.4 - 91.3%, 61.9 - 86.4%, 49.4 - 72.4%, 61.3 - 96% and 63 - 95% of the total Cd, Cu, Ni, Pb and Zn were accumulated in the 'resistant' fractions throughout the sampling period which indicated that the mobility of these metals in Kuala Gula coastal water were low in terms of contribution or contamination from anthropogenic sources. Significant (p<0.05) and positive correlation were detected between (r = 0.578) birds density and Pb metal concentration of sediments in station 2. However, the total fraction of Cu (r = 0.591 and Ni (r = -0.513) in sediments had significant p<0.05) negative correlation with crabs density in station 1. Similarly, the Pb (r = 0.657) metal concentrations in sediments was significantly p<0.05) correlated to shrimp density in station 1. Meanwhile, positive correlation, with lesser confidence (p<0.05) was also detected between mollusc density and Cd (r = 0.492) metal concentrations in station 2.

Overall, this study revealed that Kuala Gula ecosystems are still pristine with low anthropogenic sources, abundant food supply and ideal habitat for migratory shorebirds. Therefore, Kuala Gula coastal water is still in healthy condition but continuous environmental monitoring program should be implemented before it reaches alarming level as the ever increasing human activities and rapid developments in many infrastructures and industry sectors are continuously expanding. At present, the results of the study will provide important baseline data as reference for future studies on migratory shorebirds.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

KEPELBAGAIAN BURUNG HIJRAH PANTAI DAN CIRI-CIRI HABITAT DI SANTUARI BURUNG, KUALA GULA PERAK, MALAYSIA

Oleh

ROMEO M. LOMOLJO

January 2011

Pengerusi: Prof. Dr. Ahmad Ismail, PhD

**Fakulti: Sains** 

Kajian ini telah dijalankan di salah sebuah laluan persinggahan utama bagi burung-burung hijrah pantai semasa migrasi tahunan di Sanctuary Burung Kuala Gula, Perak di pantai barat Semenanjung Malaysia bagi mengenalpasti kepelbagaian burung hijrah pantai di Kuala Gula, menguji kualiti habitat, kebolehdapatan sumber makanan, keadaan persekitaran, kepekatan logam berat di dalam sedimen, organisma makrobentik, ikan dan udang bermula dari Oktober 2006-September 2007. Dua stesen persampelan telah dipilih yang merangkumi kawasan seluas 100 x 100 meter untuk setiap stesen di santuari burung Kuala Gula yang terletak di antara 04° 55.185'N, 100° 27.840'E dan 04° 55.00'N, 100° 27.761'E. Burung- burung telah dienumerasi dan dikenalpasti sehingga ke peringkat spesis secara langsung sewaktu tinjauan dijalankan. Sejumlah 61 spesis burung hijrah pantai yang terdiri daripada 12 famili telah dikenalpasti dikedua-dua stesen kajian. Jumlah keseluruhan kepadatan burung hijrah pantai semasa penghijrahan ke arah utara (7913 individu/ha) didapati lebih tinggi berbanding penghijrahan ke arah selatan (4396

individu/ha). Begitu juga, indeks kepelbagaian (H = 3.20; p<0.05) dan kekayaan spesis (d = 6.19; p<0.05) semasa penghijrahan ke arah utara didapati ternyata lebih tinggi. Walaubagaimanapun, tiada perbezaan signifikan bagi kesamarataan sebaran spesis (p>0.05), di antara penghijrahan ke arah selatan dan utara di sepanjang tempoh pengajian. Sementara itu, sepanjang kajian dijalankan, sejumlah 28 spesis makrobentos daripada 15 famili telah direkodkan dan dikenalpasti di kedua-dua stesen kajian. Index kepelbagaian spesis di stesen 2 (H'= 3.24) didapati lebih tinggi (p<0.05) berbanding stesen 1 (H'=3.13). Begitu juga dengan kekayaan spesis dimana stesen 2 (d' = 6.22) didapati mempunyai nilai yang lebih tinggi (p<0.05) berbanding stesen 1 (d'=6.11). Polichaete  $(204.3 \pm 6.84 \text{ individu/m}^3)$  merupakan kumpulan makrobentos yang paling kerap ditemui dikalangan makrobenthos, diikuti oleh molluska (156.8  $\pm$  6.3 individu/m<sup>3</sup>), ikan (132.8  $\pm$ 6.6 individu/m<sup>3</sup>) dan udang (135.3  $\pm$  4.1 individu/m<sup>3</sup>). Secara keseluruhan, didapati bahawa kepadatan serta biomass makrobentos adalah berkolerasi secara positif (r = 0.920; p<0.05) dengan kepadatan burung yang ditemui. Oleh itu, sumber makanan boleh dikatakan sebagai faktor penting yang mempengaruhi sebaran burung hijrah pantai di kawasan persisiran Kuala Gula.

Kualiti air dan nutrien di Kuala Gula didapati masih kekal di paras yang sihat. Parameter kualiti air *in situ* ( suhu, saliniti, oksigen terlarut, pH dan konduktiviti) didapati tidak menunjukkan sebarang variasi signifikan dari segi bulanan sepanjang tempoh kajian dijalankan di kedua-dua stesen. Kepekatan nitrat nitrogen (NO<sub>3</sub>-N), ammonia nitrogen total (TAN), dan fosforus reaktif terlarut (SRP) didalam air di kedua-dua stesen didapati berada pada paras yang selamat dan kekal pada paras yang minimum (SRP = 55.92 ±

 $7.88~\mu g/L$ ,  $NO_3$ -N=  $85.68 \pm 24.33~\mu g/L$ ,  $TAN = 85.91 \pm 6.54~\mu g/L$ ) di sepanjang tempoh kajian.

Bagi analisis keseluruhan pecahan geokimia bagi logam berat Cd, Cu, Pb and Zn di dalam sedimen di stesen 1 dan 2, pecahan 'resistant' telah didapati mendominasi nilai keseluruhan logam berat di sepanjang tempoh kajian dijalankan dengan julat di antara 80.4 - 91.3%, 61.9 - 86.4%, 49.4 - 72.4%, 61.3 - 96% dan 63 - 95% telah menggambarkan bahawa mobiliti metal yang berpunca daripada sumber anthropogenik di perairan Kuala Gula adalah masih rendah. Terdapat kolerasi secara positif (r = 0.578; p<0.05) di antara kepadatan burung dengan kepekatan Pb di dalam sedimen di stesen 2. Walaubagaimanapun, terdapat kolerasi secara negatif (p<0.05) diantara pecahan keseluruhan logam berat bagi Cu (r = -0.591 dan Ni (r = -0.513) di dalam sedimen dengan kepadatan populasi ketam di stesen 1. Begitu juga dengan kepekatan Pb (r = 0.657) di dalam sedimen di stesen 1 dimana terdapat kolerasi secara positive dengan kepadatan udang telah diperhatikan. Sementara itu, kolerasi positif (p<0.05) yang lemah juga dapat diperhatikan di antara kepadatan moluska dengan kepekatan Cd (r = 0.492) di stesen 2.

Secara keseluruhannya, kajian ini menunjukkan bahawa ekosistem Kuala Gula masih berada ditahap yang bersih dengan sumber antropogenik yang minimal, kebolehdapatan sumber makanan yang banyak dan habitat yang ideal untuk burung-burung hijrah pantai. Walaubagaimanapun, pemantauan yang berterusan di kawasan ini perlu dilaksanakan bagi memastikan sebarang gangguan atau perubahan ekosistem akibat aktiviti manusia

dan pembangunan pesat dalam sektor infrastruktur dan industri yang dijalankan dapat dikesan diperingkat awal lagi bagi mengelakkan kesan yang lebih buruk terhadap kepelbagaian serta habitat hidupan liar dikawasan ini. Maka, diharapkan hasil kajian ini dapat dijadikan sebagai rujukan dan memberi panduan asas untuk kajian di masa hadapan.



#### **ACKNOWLEDGEMENTS**

I would like to express my deepest gratitude to my main supervisor Professor Dr. Ahmad Bin Ismail for his expertise and consistent constructive advice, ideas intelligent counsel, motivation, suggestion encouragement and guidance for the successful completion of my study.

My sincere thanks also goes to my co-supervisors Associate Professor Dr. Yap Chee Kong and Dr. Abdul Rahim Ismail for their professional advice and comments. My appreciation and thanks to the Malaysian Wildlife Department for providing the opportunity to conduct research in Kuala Gula Bird Sanctuary and for giving me staff to assist during the study period. A special thanks to MOSTI-Science Fund No. 06-01-04-SF0715.

A word of thanks goes to my loving and understanding wife Evangeline, my daughters En-En and Barbie for their love, understanding and motivation which gave me strength throughout the whole period of my study. I would also like to thank goes to Pakmin, Ikram, Franklin, Wanhee, Hatta, Norhaida, Safura, Zulika, for their friendship and technical assistance. My special appreciation goes to those whose names are not mentioned but definitely have contributed so much in my pursuit for knowledge and scientific advancement.

Lastly, I am very grateful to all my Filipino friends, and for the helpful comments provided by Ivy Esteban and Hazel Peralta, Orly and family, Ruben and family, Elvie, Elsa and Amelia. Their names will always be encrypted in my heart, for their smiles, hugs, embraces, laughs, problems, nice foods, parties, games, intellectual discussions, but most especially for the love and prayers, the best and foremost thing I could ever receive from them in times of great needs. They are my family away from home. Mere words can not describe my appreciation to all of them.

And most of all to the Almighty God, in whom nothing is impossible.

I certify that an Examination committee has met 6<sup>th</sup> January 2011 to conduct the final examination of Romeo M. Lomoljo on his Doctor of Philosophy thesis entitled "Diversity of Migratory Shorebirds and Their Habitat Characteristics in Kuala Gula Bird Sanctuary Perak, Malaysia" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980. The Committee recommends that the student be awarded the doctor of Philosophy Degree.

Members of the Examination Committee were as follows:

#### Dr. Hishamuddin Omar, PhD

Lecturer
Faculty of Science
Universiti Putra Malaysia
(Chairman)

**Dr. Mohamed Zakaria Hussin, PhD**Professor Madya
(Internal Examiner)

**Dr. Nor Azwady Abd. Aziz, PhD**Professor Madya
(Internal Examiner)

Masanori Sato, PhD Professor (External Examiner)

#### NORITAH OMAR, PhD

Associate Professor and Deputy Dean School of Graduate Studies Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The Members of the Supervisory Committee were as follows.

#### Ahmad Ismail, PhD

Professor Faculty of Science Universiti Putra Malaysia (Chairman)

#### Yap Chee Kong, PhD

Associate Professor Faculty of Science Universiti Putra Malaysia (Member)

#### **Abdul Rahim Ismail, PhD**

Lecturer
Faculty of Science
Universiti Putra Malaysia
(Member)


### HASANAH MOHD GHAZALI, PhD

Professor and Dean School of Graduate Studies Universiti Putra Malaysia

Date:

#### **DECLARATION**

I declare that the thesis is my original work except for quotation and citations which have been duly acknowledge. I also declare that it has not been previously, or concurrently, submitted for any other degree at Universiti Putra Malaysia at any institutions.



## TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENT	X
APPROVAL	xii
DECLARATION	xiv
LIST OF TABLES	xix
LIST OF FIGURES	xxi
LIST OF ABBREVIATION	XXV
CHAPTER	
1 INTRODUCTION	1
1.1 General	1
1.2 Statement of the problem	1 3
1.3 Significance of the study	5
1.4 Objectives	6
1.5 Scope and content of the thesis	7
2 LITERATURE REVIEW	9
2.1 Coastal areas of Peninsular Malaysia	9
2.2 Birds composition of migratory and non-migratory species	10
2.3 Shorebirds' characteristics	12
2.4 Shorebirds migration	13
2.5 Effects of food availability on shorebird distribution	14
2.6 Macrobenthos as sources of food	15
2.7 Food availability and feeding habits	16
2.8 Causes of the declining number of migratory shorebirds	18
2.8.1 Habitat destruction	18
2.8.2 Hunting and poaching of shorebirds	20
2.8.3 Disturbances	21
2.9 Heavy metal pollution in the coastal environment	22
2.10 Heavy metals in intertidal sediment	26
2.11 Speciation of heavy metals in the sediment	27
2.12 Effects of heavy metals to organisms	29
2.13 Assessment of habitat quality	32
2.14 The effects of hazardous chemicals to shorebirds habitat	34
2.15 Water quality	38

3 GENERAL MATERIALS AND METHODS	40
3.1 Sampling location	40
3.1.1 Description of the sampling site	41
3.2 Data collection	42
3.2.1 Shorebird survey	42
3.2.2 Sampling of macrobenthic fauna	43
3.2.3 Water	44
3.2.4 Sediment sampling	45
3.3 Laboratory work	45
3.3.1 Nutrients analysis	45
3.3.2 Sorting and identification of macrobenthic fauna	48
3.3.3 Heavy metal in sediments	49
3.3.4 Heavy metal in macrobenthos	55
3.3.5 Bird and macrobenthos density	56
3.3.6 Biomass determination	56
3.4 Data analysis	57
4124000 42000	
4 RESULTS	59
4.1 Birds diversity	59
4.1.1 Birds density and diversity	58
4.1.2 Bird species relative abundance	62
4.2 Temporal and spatial distribution of bird density relationship	65
4.3 Macrobenthos density	70
4.4 Monthly variation of macrobenthos, fish and shrimp density	74
4.4.1 Polychaetes	74
4.4.2 Crabs	75
4.4.3 Shrimp	76
4.4.4 Fish	77
4.4.5 Molluscs	78
4.5 Macrobenthos, fish and shrimps biomass	79
4.6 Macrobenthos diversity	80
4.6.1 Species diversity index	80
4.7 Temporal and spatial distribution of macrobenthos relationship	80
4.8 Variations in environmental parameters	85
4.8.1 Rainfall	85
4.8.2 Temperature	86
4.8.3 Salinity	86
4.8.4 Dissolved oxygen	86
4.8.5 pH	87
4.8.6 Total ammonia nitrogen (TAN)	87
4.8.7 Soluble reactive phosphorous (SRP)	88
4.8.8 Nitrate + Nitrite- N	88
4.9 Heavy metal concentrations of the intertidal sediments in Kuala Gula	91
4.9.1 Total metal concentrations	91

	4.9.2 Metal speciation in station 1	94
	4.9.3 Metal speciation in station 2	95
	4.10 Percentages of geochemical fractions of sediment between station 1 and station 2	98
	4.11 Comparison of non-resistant and resistant fractions between station 1	104
	and station 2	
	4.12 Correlation between bird density and environmental parameters	107
	4.13 Heavy metal concentrations of macrobenthos, fish and shrimps in station 1	108
	4.14 Heavy metal concentrations of macrobenthos, fish and shrimps in station 2	108
	4.15 Correlation between heavy metals of macrobenthos, fish and shrimps and geochemical fraction of metals in the intertidal sediments	111
	4.15.1 Correlation of Cd between geochemical fractions of sediments and macrobenthos	112
	4.15.2 Correlation of Cu between geochemical fractions of sediments and macrobenthos	112
	4.15.3 Correlation of Ni between geochemical fractions of sediments and macrobenthos	113
	4.15.4 Correlation of Pb between geochemical fractions of sediments and macrobenthos	114
	4.15.5 Correlation of Zn between geochemical fractions of sediments and macrobenthos	115
	4.16 Correlation between physico-chemical parameters and macrobenthos density in station 1	115
	4.17 Correlation between physico-chemical parameters and macrobenthos density in station 2	116
	4.18 Relationship between bird density, macrobenthos, fish and shrimps density in station 1 and station 2	117
	4.19 Relationship between bird density and the biomass of macrobenthos in station 1 and station 2	117
	4.20 Relationship between macrobenthos, fish and shrimps density and biomass in station1 and station 2	118
5	DISCUSSION	125
	5.1 Bird density and diversity	125
	5.2 Temporal and spatial distribution of migration shorebirds	123
	5.3 Macrobenthos density	127
	5.4 Diversity, species richness and evenness of macrobenthos	135
	5.5 Variation in environmental parameters	136
	5.5.1 Physico-chemical parameters	136
	5.5.2 Nutrients in seawater	140
	5.6 Heavy metals in sediments	143
	5.7 Sediment Quality Guidelines	148

5.7.1 Effective range low (ERL)/Effective range median (ERM)	148
5.8 Heavy metal concentrations of macrobenthos, fish and shrimps in station 1 and 2	150
5.9 Correlation of Cd between macrobenthos and sediments	156
5.10 Correlation of Cu between macrobenthos and sediments	158
5.11 Correlation of Ni between macrobenthos and sediments	159
5.12 Correlation of Pb between macrobenthos and sediments	160
5.13 Correlation of Zn between macrobenthos and sediments	161
5.14 Correlation between physicochemical parameter and macrobenth in station 1	nos 162
5.15 Correlation between physicochemical parameter and macrobenth in station 2	nos 163
5.16 Relationship between different environmental parameters	164
5.17 Correlation between bird density and macrobenthos density	165
5.18 Correlation between bird density and macrobenthic biomass	166
5.19 Correlation between macrobenthos density and biomass	168
6 SUMMARY, CONCLUSION AND RECOMMENDATION FO FUTURE RESEARCH	<b>OR</b> 170
REFERENCES	176
BIODATA OF STUDENT	208
LIST OF PUBLICATION	209