



UNIVERSITI PUTRA MALAYSIA

**SCREENING OF MALAYSIAN MEDICINAL/HERBS AND AQUATIC
PLANTS FOR PANCREATIC LIPASE INHIBITORY ACTIVITIES AND
IDENTIFICATION OF ACTIVE CONSTITUENT**

MUHAMMAD ABUBAKAR ADO

FSTM 2010 11



**SCREENING OF MALAYSIAN MEDICINAL/HERBS AND AQUATIC PLANTS
FOR PANCREATIC LIPASE INHIBITORY ACTIVITIES AND
IDENTIFICATION OF ACTIVE CONSTITUENT**

By

MUHAMMAD ABUBAKAR ADO

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

June 2010



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Masters of Science

**SCREENING OF MALAYSIAN MEDICINAL/HERBS AND AQUATIC PLANTS
FOR PANCREATIC LIPASE INHIBITORY ACTIVITIES AND
IDENTIFICATION OF ACTIVE CONSTITUENT**

By

MUHAMMAD ABUBAKAR ADO

June 2010

Chairperson : Professor Hasanah Mohd Ghazali, PhD

Faculty : Food Science and Technology

The compounds that help to slow down the digestion of triacylglycerols inside pancreatic and small intestine of human play an important role in the control of obesity. In this study the effects of 80% methanol extracts of different parts (seeds, fruits, leaves, flowers, roots) of some medicinal/herbals and aquatic plants for their anti-lipase activity was determined. The effect of each plant extract on porcine pancreatic lipase was measured based on titrimetric method. The results revealed of the one hundred (100) plant samples screened, eighty eight (88) of the extracts were observed to inhibit while twelve (12) were found to promote the activity and only two extracts did not show any activity on the enzyme. Four of the plant extracts *Archidendron jiringa* (Jack) I.C Nielsen, *Averrhoa carambola* L. *Cynometra cauliflora* and *Alevrites moluccana* (L.) Willd have shown the highest anti-lipase activity of 100%, and twenty one (21) extracts observed to have high



activity above 70%. Eighteen (18) out of these twenty one (21) have anti-lipase activity of greater than 80%. Thirteen (13) have presented moderate inhibition with a high proportion of 62% exhibited lower inhibition. Twenty nine (29) extracts out of these (86) can be regarded as very poor inhibitors, as their percent inhibition was less than 20% and only 1.74% was shown to have no inhibitory activity. In the process of isolating the active compounds, liquid-liquid partitioning of the crude methanolic extract of *Cynometra cauliflora* leaves was carried out. Five different fractions hexane, DCM, EtOAc, n-BuOH and aqueous was obtained. All the fractions were tested for anti-lipase activity and the active components reside mainly in ethyl acetate fraction. The isolation and identification of active compounds from the most active fraction (ethyl acetate) was further fractionated using silica gel column chromatography (normal phase). The active fraction was further purified by the use of Sephadex LH-20 chromatography. The structure of the active compound from ethyl acetate fraction of *Cynometra cauliflora* was found to be kaempferol-3-*O*-rhamnoside. The structure of this active compound was identified from spectroscopic data analysis of IR, MS, ¹H NMR, ¹³C NMR and 2D-NMR.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master of Science

PENYARINGAN TUMBUHAN UBATAN/HERBA DAN TUMBUHAN YANG TERDAPAT DI MALAYSIA UNTUK AKTIVITI PERENCATAN LIPASE PANKREAS DAN PENGENALPASTIAN KOMPONEN AKTIF

Oleh

MUHAMMAD ABUBAKAR ADO

Jun 2010

Pengerusi : Profesor Hasanah Mohd Ghazali, PhD

Fakulti : Sains dan Teknologi Makanan

Komponen yang melambatkan penghadaman triasilgliserol di dalam pankreas dan usus kecil manusia, memainkan peranan yang penting dalam mengawal kegemukan. Kesan metanol ekstrak daripada pelbagai bahagian yang berlainan (biji, buah, daun, bunga dan akar) daripada tumbuhan ubatan/herba dan akuatik telah dikaji bagi aktiviti anti-lipase. Kesan setiap ekstrak tumbuhan ini terhadap lipase porcine pankreas diukur berdasarkan kaedah pentitratan. Keputusan menunjukkan daripada seratus (100) sampel tumbuhan yang di periksa, lapan puluh lapan (88) daripadanya didapati menunjukkan perencatan manakala dua belas (12) lagi didapati menggalakkan aktiviti enzim tersebut. Empat ekstrak tumbuhan iaitu *jering*, *belimbing besi*, *nam-nam* dan *buah keras* telah menunjukkan aktiviti anti-lipase yang paling tinggi iaitu 100%, dengan dua puluh satu

(21) ekstrak didapati mempunyai aktiviti yang tinggi melebihi 70%. Lapan belas (18) daripada dua puluh satu (21) mempunyai aktiviti anti-lipase yang melebihi 80%. Tiga belas (13) telah mempamerkan perencatan secara sederhana dengan sebahagian besar iaitu 62% menunjukkan perencatan yang lebih rendah. Dua puluh sembilan (29) ekstrak daripadanya (88) boleh dipertimbangkan sebagai perencat yang lemah, disebabkan oleh peratusan perencatannya yang kurang daripada 20% dan hanya 1.74% menunjukkan tiada aktiviti. Proses penulenan komponen aktif daripada pokok nam-nam dimulakan dengan proses pempartisian cecair-cecair terhadap ekstrak metanol memberikan 5 fraksi iaitu heksana, dikloromethana, etil asetat, butanol dan air. kesemua fraksi ini di uji aktiviti anti-lipase dan di dapati komponen bioaktif kebanyakannya terdapat dalam fraksi etil asetat. Pengasingan komponen aktif ini daripada fraksi yang paling aktif (etil asetat) kemudiannya di teruskan lagi dengan menggunakan kaedah kromatografi turus gel silika (fasa normal). Pecahan yang aktif daripada kromatografi fasa normal ini kemudiannya di tulenkan dengan menggunakan kromatografi pembezaan saiz, Sephadex LH-20. Komponen aktif daripada pokok nam-nam dihenalpasti sebagai kaempferol-3-*O*-rhamnosida. Struktur ini komponen dikenalpasti menggunakan analisis data spektroskopik IR, MS, ^1H NMR, ^{13}C NMR, dan 2D-NMR.



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Masters of Science. The members of the Supervisory were as follows:

Hasanah Mohd Ghazali, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairperson)

Abdulkarim Sabo Mohammed, PhD

Associate Professor.
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Faridah Abas, PhD

Doctor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of graduate Studies
Universiti Putra Malaysia

Date:



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

MUHAMMAD ABUBAKAR ADO

Date: 29th June 2010



ACKNOWLEDGEMENTS

First of all, all praise due to Allah, lord of the world. Only by his Grace and Mercy this work has been finished.

I would like to express my intensive and earnest gratitude to my supervisor, Prof. Dr. Hasanah Mohd Ghazali, for her sincere, constructive suggestions, invaluable guidance, and continuous discussion throughout this project. Also, I wish to express my very much thankful to her for the financial support, without this support it is impossible for me to pursue the success of this project.

I would like to extend my sincere appreciations and gratitude to my Co-supervisors: Dr. Abdulkarim Sabo Mohammed and Dr. Faridah Abas for their guidance, suggestions, honestly encouragements throughout this research.

Special thanks are extended to my Lab mates for their encouragements and helps throughout these years.

My genuine appreciation and gratitude goes to my family members: my parents, my brothers and sisters for their prayers, encouragements and support through good and bad times.



TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	iv
APPROVAL	vi
DECLARATION	viii
ACKNOWLEDGMENTS	ix
LIST OF TABLES	xii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xvii
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	
2.1 Obesity	8
2.1.1 Types of Obesity	8
2.1.2 Causes of Obesity	10
2.2 Fat digestion	12
2.3 Lipase (triacylglycerol hydrolase)	15
2.3.1 Method for determining lipase activity	16
2.3.2 Lipase inhibitors	18
2.4 Anti-obesity drugs	19
Synthetic anti-obesity drug	21
Orlistat	22
Mechanism of action of Orlistat	23
Sibutramine	25
2.5 Natural anti-obesity agents	26
	29
3 SCREENING OF MEDICINAL/HERBS AND AQUATIC PLANTS FOR ANTI-LIPASE ACTIVITY	
3.1 Introduction	
3.2 Materials and methods	67
3.2.1 Materials	69
3.2.2 Methods	69
3.2.2.1 Extraction of plant materials	73
3.2.2.2 Assay for pancreatic lipase activity	73
3.3 Results and Discussion	74
3.4 Conclusion	76



4 ISOLATION AND STRUCTURE ELUCIDATION OF LIPASE INHIBITOR FROM <i>Cynometra cauliflora</i> (Nam Nam)	93
4.1 Introduction	
4.2 Materials and methods	95
4.2.1 Plant extraction	96
4.2.2 General	97
4.2.3 Instrumentation	97
4.2.3.1 Infrared Spectroscopy (IR)	97
4.2.3.2 Nuclear magnetic resonance	97
4.2.3.3 Mass Spectra (MS)	98
4.2.3.4 High Performance Liquid Chromatography (HPLC)	98
4.2.3.5 Ultra violet	99
4.3 Methods	100
4.3.1 Chromatography	100
4.3.2 Thin Layer Chromatography	100
4.3.3 Dyeing Reagent for TLC	100
4.3.4 Vanillin-Sulfuric Acid Solution	101
4.4 Isolation of Lipase Inhibitor from the leaves of <i>Cynometra cauliflora</i> (Nam Nam)	
4.4.1 Solvent-solvent fractionation	102
4.4.2 Column chromatography of the EtOAc fraction of <i>Cynometra cauliflora</i> leaves	102
4.4.3 Isolation of Compound 1	103
4.5 Results and Discussion	104
4.5.1 Isolation of Compound 1 from <i>Cynometra cauliflora</i> leaves	108
4.5.2 Characterization of Compound 1	108
4.5.3 Identification of Compound 1	108
4.5.4 Conclusion	112
	125
5 SUMMARY, GENERAL CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCH	129
REFERENCES	
APPENDICES	131
BIODATA OF AUTHOR	144
	157

