



**UNIVERSITI PUTRA MALAYSIA**

**PHYSICO - CHEMICAL, THERMAL AND RHEOLOGICAL  
PROPERTIES OF VARIOUS FISH PROTEIN - SAGO STARCH  
FORMULAS IN KEROPOK LEKOR**

**MOHAMED ABD ELGADIR MOHAMED SAEED**

**FSTM 2010 14**



**PHYSICO - CHEMICAL, THERMAL AND RHEOLOGICAL PROPERTIES OF  
VARIOUS FISH PROTEIN - SAGO STARCH FORMULAS IN *KEROPOK LEKOR***

**By**

**MOHAMED ABD ELGADIR MOHAMED SAEED**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

**September 2010**



**DEDICATED TO MY PARENTS,  
MY BROTHERS,  
WIFE, KIDS AND FRIENDS**



Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in  
Fulfillment of the requirement for the degree of Doctor of Philosophy

**PHYSICO - CHEMICAL, THERMAL AND RHEOLOGICAL PROPERTIES OF  
VARIOUS FISH PROTEIN - SAGO STARCH FORMULAS IN *KEROPOK LEKOR***

By

**MOHAMED ABD ELGADIR MOHAMED SAEED**

**September 2010**

**Chairman: Professor Jamilah Bakar, PhD**

**Faculty: Food Science and Technology**

This study was conducted to investigate the interaction between fish protein and sago starch using *keropok lekor* as a model with the following objectives: to study the thermal and physical properties of the model, to optimize the contents used in the model based on their thermal and rheological properties, to study the effects of sodium chloride and sucrose on transition temperatures of gelatinization and to investigate the effect of sodium chloride and sucrose on cooking time of the model. The ingredients used in preparing the formulations were red tilapia (*Oreochromis spp.*) minced fish, sago starch (*Metroxylon sagu*), and water in the ranges of 20 - 50, 10 - 40 and 10 - 35, respectively. A two-level factorial design was carried out to obtain the experimental formulations of the model. Differential scanning



calorimetry (DSC), texture profile analysis (TPA), sensory evaluation and the changes in thermophysical properties such as thermal conductivity ( $k$ ), specific heat capacity ( $C_p$ ), density ( $\rho$ ), thermal diffusivity ( $\alpha$ ) and the effects of sodium chloride and sucrose concentrations on the cooking time of the optimized fish protein-sago starch model were conducted. The onset temperatures ( $T_o$ ) of the different formulations of the fish protein - sago starch model varied from  $60.9 \pm 0.5$  -  $80.8 \pm 0.4^\circ\text{C}$  with an average value of  $70.9 \pm 0.5^\circ\text{C}$ . The formulations had peak temperatures of gelatinization that varied from  $65.5 \pm 0.1$  -  $86.0 \pm 0.4^\circ\text{C}$  with average value of  $75.8 \pm 0.3^\circ\text{C}$ . Increasing the ratios of the minced fish in the formulations was found to be the most significant factor affecting the values of  $T_o$  and  $T_p$ . The highest hardness value 50.9 N was obtained in the sample formulated with 5: 4: 1 minced fish: sago starch: water, respectively followed by 45.0 N which was obtained in the sample formulated with the ratios of 3.5 minced fish 4 sago starch: 2.25 water. The sensory evaluation study showed that the panelists preferred the model formulated with 2: 1: 1 minced fish: sago starch: water followed by 3.5: 2.5: 2.25 of the components, respectively. The optimum formulation of the model was highly dependant on the ratios of fish protein: sago starch. The addition of NaCl to the optimized formulation resulted in the shifting of the  $T_o$  from  $82.2 \pm 0.9^\circ\text{C}$  to  $98.2 \pm 0.1^\circ\text{C}$ ,  $T_p$  from  $94.8 \pm 0.2^\circ\text{C}$  to  $106.2 \pm 0.3^\circ\text{C}$  and  $T_c$  from  $108.0 \pm 0.9^\circ\text{C}$  to  $116.3 \pm 0.2^\circ\text{C}$ . However, the addition of sucrose shifted  $T_o$  from  $82.2 \pm 0.9^\circ\text{C}$  to  $111.4 \pm 0.2^\circ\text{C}$ ,  $T_p$  from  $94.8 \pm 0.2^\circ\text{C}$  to  $118.2 \pm 0.2^\circ\text{C}$  and  $T_c$  from  $108.0 \pm 0.9^\circ\text{C}$  to  $119.3 \pm 0.3^\circ\text{C}$ . During frying an increase in the thermal conductivity ( $k$ ), specific heat capacity ( $c_p$ ) and thermal diffusivity ( $\alpha$ ) values was observed. The  $k$  value of the model increased dramatically from 0.418 to 0.584  $\text{W}/\text{m}^2\cdot^\circ\text{C}$  in the control. The calculated  $k$  values confirmed that the NaCl and sucrose caused an increase in  $k$  values of the model from 0.419

to 0.585 W/m<sup>2</sup>.°C for NaCl and from 0.423 to 0.588 W/m<sup>2</sup>.°C for sucrose treated ones. The calculated values of thermal diffusivities of the model varied from 1.19 x 10<sup>-7</sup> to 1.51 x 10<sup>-7</sup> m<sup>2</sup> s<sup>-1</sup> during deep - fat frying. In summary, it can be concluded that the thermal properties of the model were highly dependant on the ratios of fish protein, sago starch and water used. The optimum formulation was obtained in the model formulated with 2: 1.01: 1 minced fish: sago starch: water, respectively. The shifts in the transition temperatures values of  $T_o$ ,  $T_p$  and  $T_c$  caused by NaCl were 16, 11.4 and 8.3°C, respectively. However, sucrose shifted the transition temperatures values of  $T_o$ ,  $T_p$  and  $T_c$  to 29.2, 23.4 and 11.3°C, respectively. Sucrose was found to be more effective in increasing the onset and conclusion temperature of gelation of the model. The actual cooking time of the control sample of the model was 180 s and the predicted cooking time of the model according to Ansari's equation was 195 s. The model with either added sucrose or NaCl needed more time to be cooked.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**SIFAT- SIFAT FISIKA - KIMIA, TERMA DAN RHEOLOGI BAGI BERBAGAI FORMULASI PROTEIN IKAN - KANJI SAGU DI DALAM *KEROPOK LEKOR***

Oleh

**MOHAMED ABD ELGADIR MOHAMED SAEED**

**September 2010**

**Pengerusi: Professor Jamilah Bakar, PhD**

**Fakulti: Sains dan Teknologi Makanan**

Kajian ini dijalankan untuk menyiasat interaksi antara protein ikan dan kanji sagu dengan menggunakan keropok lekor sebagai model dan mempunyai objektif - objektif berikut: sifat-sifat terma dan fizikal sistem, pengoptimuman kandungan yang digunakan dalam model berdasarkan sifat reologi, kesan penambahan bahan (natrium klorida dan sukrosa) terhadap suhu peralihan pengelatinan, dan jangkaan masa memasak akibat dipengaruhi oleh penambahan bahan (NaCl dan sukrosa) dikaji. Nisbah bahan - bahan yang digunakan untuk menyediakan sampel adalah ikan tilapia merah (*Oreochromis spp.*) yang dicincang (MF), kanji sagu (*Metroxylon sagu*) (SS) dan air (W), masing-masing di dalam lingkungan 20-50, 10 - 40 dan 10.35. Rekabentuk faktorial dua - peringkat telah digunakan untuk mendapatkan formulasi ujikaji. Teknik kalorimetri pergimbasan pembeza (DSC), analisis tekstur profil



(TPA), penilaian sensori dan perubahan sifat termofizikal seperti terma konduktivi ( $k$ ), kapasiti haba spesifik ( $C_p$ ), ketumpatan ( $\rho$ ), difusiviti terma ( $\alpha$ ) dan kesan kepekatan natrium klorida dan sukrosa ke atas jangkaan masa memasak model protein ikan - kanji sago yang dioptimumkan telah dijalankan. Suhu awal gelatinasi ( $T_o$ ) bervariasi dalam lingkungan  $60.9 \pm 0.5 - 80.8 \pm 0.4^\circ\text{C}$  dalam formulasi protein ikan - kanji sago yang berlainan dengan nilai purata  $70.9 \pm 0.5^\circ\text{C}$ . Formulasi - formulasi tersebut mempunyai suhu puncak gelatinasi yang bervariasi dalam lingkungan  $65.5 \pm 0.1 - 86.0 \pm 0.4^\circ\text{C}$  dengan nilai purata  $75.8 \pm 0.3^\circ\text{C}$ . Peningkatan nisbah ikan yang dicincang dalam formulasi ditemui sebagai faktor signifikansi yang menjejaskan nilai  $T_o$  dan  $T_p$ . Nilai kekerasan tertinggi iaitu 50.9 N dicapai dalam sampel berformulasi 5:4:1, masing-masing ialah ikan dicincang: kanji sago: air, diikuti 45.0 N di mana ia dicapai oleh sampel berformulasi dengan nisbah 3.5 ikan yang dicincang: 4 kanji sago: 2.25 air. Kajian penilaian sensori menunjukkan bahawa ahli panel lebih suka model yang diformulakan dengan 2: 1: 1 ikan yang dicincang: kanji sago: air diikuti 3.5: 2.5: 2.25. Formulasi optima model sangat bergantung kepada nisbah campuran protein ikan - kanji sago dan diperolehi dalam model berformula 2: 1.01 : 1. Penambahan NaCl ke atas formula optima menyebabkan pengalihan  $T_o$  daripada  $82.2 \pm 0.9$  kepada  $98.2 \pm 0.1$ ,  $T_p$  daripada  $94.8 \pm 0.2$  kepada  $106.2 \pm 0.3$  dan  $T_c$  daripada  $108.0 \pm 0.9$  kepada  $116.3 \pm 0.2$ . Namun, penambahan sukrosa mengalihkan  $T_o$  daripada  $82.2 \pm 0.9$  kepada  $111.4 \pm 0.2$ ,  $T_p$  daripada  $94.8 \pm 0.2$  kepada  $118.2 \pm 0.2$  dan  $T_c$  daripada  $108.0 \pm 0.9$  kepada  $119.3 \pm 0.3$ . Sukrosa ditemui lebih berkesan dalam meningkatkan suhu awal dan akhir gelatinasi model. Semasa penggorengan, peningkatan terma konduktivi ( $k$ ), kapasiti haba spesifik ( $C_p$ ), ketumpatan ( $\rho$ ), dan difusiviti terma ( $\alpha$ ) diperhatikan. Nilai  $k$  model meningkat dengan dramatik daripada 0.418 kepada  $0.584 \text{ W/m}^2 \text{ }^\circ\text{C}$  dalam kawalan. Nilai  $k$  yang dikira



menyakinkan bahawa NaCl dan sukrosa menyebabkan peningkatan dalam nilai  $k$  daripada 0.419 kepada 0.585  $\text{W/m}^2 \text{ } ^\circ\text{C}$  untuk model yang diperlakukan oleh NaCl dan daripada 0.423 kepada 0.588  $\text{W/m}^2 \text{ } ^\circ\text{C}$  untuk model yang diperlakukan oleh sukrosa. Nilai difusiviti terma model yang dikira bervariasi daripada  $1.19 \times 10^{-7}$  kepada  $1.51 \times 10^{-7} \text{ m}^2 \text{ s}^{-1}$  semasa penggorengan lemak dalam. Sebagai rumusan, dapat disimpulkan sifat-sifat terma model sangat bergantung kepada nisbah protein ikan, kanji sagu dan air yang digunakan. Formulasi optima telah didapati daripada model berformulasi 2: 1.01: 1 ikan yang dicincang: kanji sagu: air. Pertukaran nilai suhu peralihan  $T_o$ ,  $T_p$  dan  $T_c$  disebabkan oleh NaCl adalah 16, 11.4 dan  $8.3^\circ\text{C}$ . Bagaimanapun, sukrosa telah menukar nilai-nilai suhu peralihan masing - masing,  $T_o$ ,  $T_p$  dan  $T_c$  kepada 29.2, 23.4 dan  $11.3^\circ\text{C}$ . Sukrosa didapati lebih efektif dalam meningkatkan suhu awal dan kesimpulan gelatinasi model tersebut. Jangka masa memasak sebenar untuk sampel kawalan model adalah 180 saat dan jangka masa jangkaan model mengikuti persamaan Ansari adalah 195 saat. Model yang ditambah dengan sama ada sukrosa atau NaCl memerlukan lebih masa untuk dimasak.

## ACKNOWLEDGEMENTS

First of all, I would like to thank the Almighty Allah for keeping me healthy to complete this study successfully. I would also like to express my unlimited and deepest thanks to the Chairman of my supervisory committee, Prof. Dr. Jamilah Bakar for her continuous assistance, excellent guidance, patience and forgiveness. She had strongly encouraged and supported me every moment from the beginning until the end of this study. My appreciations and gratefulness are also extended to the members of supervisory committee; Prof. Dr. Russly Abdul Rahman, Dr. Roselina Karim and Assoc. Prof. Dr. Kassim Ali Abbas for their time, kindness and constructive comments during all the committee meetings. I am very grateful to the Malaysian Technical Co-operative Programme (MTCP) for providing me with great financial support. I also owe great thanks to Mr. Dzulkifly Mat Hashim for his good suggestions and guidance especially with regards rheological studies. During the course of this study, I have come to recognize many individual who have kindly supported me throughout. I am indebted to Prof. Dr. Yaakob Che Man for his kind permission for allowing me to use Differential Scanning Calorimeter in Faculty of Food Science and Technology, UPM. My appreciations and thanks also go to the technicians who had contributed to this study either by giving their time or knowledge. Among those who deserve special notice and thanks in Faculty of Food science and Technology, UPM are: Mr. Razali, MS. Jamaliah in the Food Processing Laboratory, Mr. Azman in the Biochemistry Laboratory, Mr. Soib, Mr. Amran, Mr. Mohammad Noh in the Food Engineering Laboratory, Mr. Mohammad Kidin and Mr. Azhar in the Pilot Plant, Mr. Syed and his team in the IT unit. Last but not least I would like to express my appreciations to all my friends in the Fish Laboratory for their



friendly and caring assistance and discussions. All of the above people have contributed to a certain extent in this study. Therefore, the merits of the study are due to them; however, on the other hand whatever shortfalls that occur are due to my negligence.



I certify that an Examination Committee has met on 30/09/2010 to conduct the final examination of Mohamed Abd Elgadir Mohamed Saeed on his Ph.D. thesis entitled “PHYSICO - CHEMICAL, THERMAL AND RHEOLOGICAL PROPERTIES OF VARIOUS FISH PROTEIN - SAGO STARCH FORMULAS IN *KEROPOK LEKOR*” in accordance with Universiti Pertanian Malaysia Doctor of philosophy Act 1980 and Universiti Pertanian Malaysia Doctor of philosophy Regulations 1981. The committee recommends that the student be awarded the Doctor of philosophy.

Members of the Examination Committee were as follows:

**Name of Chairman, PhD**

Associate Professor Dr. Badlishah Sham Baharin  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Name of Examiner 1, PhD**

Associate Professor Dr. Sharifah Kharidah Syed Muhamad  
Faculty of Food Science and Technology  
(Internal Examiner)

**Name of Examiner 2, PhD**

Associate Professor Dr. Md Zaidul Islam Sarker  
Faculty of Food Science and Technology  
(Internal Examiner)

**Name of External Examiner**

Professor Dr. Nazlin K. Howell  
Faculty of Health and Medical Sciences  
University of Surrey  
United Kingdom  
(External Examiner)

---

**SHAMSUDDIN SULAIMAN, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 18 January 2011



This thesis was submitted to the Senate of Univesiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Jamilah Bakar, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Russly Abdul Rahman, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Member)

**Roselina Karim, Lecturer**

Faculty of Food Science and Technology  
Univesiti Putra Malaysia  
(Member)

**Kassim Ali Abbas, Associate Professor**

Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Member)

---

**HASANAH MOHD. GHAZALI, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:



## **DECLARATION**

I declare that the thesis is my original work except for the quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and it not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

---

**MOHAMED ABD EGADIR MOHAMED SAEED**

Date: 30 September



## LIST OF TABLES

Table		Page
1	Values of constants in Ansari's equation	54
2	Proximate analysis of the minced fish and sago starch used in fish protein - sago starch food system	66
3	Central composite data obtained for : independent and response variables (mean $\pm$ SD) measured by differential scanning calorimeter (DSC)	69
4	Estimated regression coefficient $R^2$ , adjusted $R^2$ and lack of fit values for the response surface model	70
5	Analysis of variance (ANOVA) showing regression coefficient of the first and second order polynomial regression model	71
6	Texture profile analysis parameters of the protein-starch <i>model</i> formulated with different starch - fish ratios	82
7	Scores attributed to texture and overall acceptance of fish protein-sago starch model in the sensory evaluation	85
8	Central composite design: independent ( $X_i$ ) and response ( $Y_i$ ) variables (mean $\pm$ SD) measured by rheometer	90
9	Regression coefficient, $R^2$ , adjusted $R^2$ probability values and lack of fit for the variables	91
10	ANOVA and regression coefficient of the first and second order polynomial regression model	92
11	Texture profile analysis parameters of the local fish protein-starch model as compared to the optimized sample of the fish protein - sago starch model	102
12	Effect of sodium chloride concentration on the gelation of the optimized starch - protein model based on DSC measurement	114
13	Effect of sodium chloride concentration on the rheological properties of the optimized starch - protein model based on rheometer measurement	114
14	Effect of sucrose concentration on the gelation of the optimized starch - protein model based on D SC measurement	120



15	Effect of sucrose concentration on rheological properties of the optimized starch - protein model	120
----	---	-----





## LIST OF FIGURES

<b>Figure</b>	<b>Page</b>
1 The different types of structure of proteins in solution	4
2 Structure of (A) amylose and (B) amylopectin	32
3 Division of supplied thermal energy $Q$ into streams of thermal energy ( $Q_1 - Q_6$ ) during food frying	49
4 Response surface plots for onset temperature of gelatinization of fish protein - sago starch model dough as a function of (a) minced fish and sago starch ratios, (b) minced fish and water ratios and (c) sago starch and water ratios based on Differential Scanning Calorimetry (DSC) measurement	72
5 Response surface plots for peak temperature of gelatinization of fish protein-sago starch model as a function of (a) minced fish and sago starch ratios, (b) minced fish and water ratios and (c) sago starch and water ratios based on Differential Scanning Calorimetry (DSC) measurement	75
6 Response surface plots for conclusion temperature of fish protein-sago starch model as a function of (a) minced fish and sago starch ratios, (b) minced fish and water ratios and (c) sago starch and water ratios based on Differential Scanning Calorimetry (DSC) measurement	77
7 Response surface plots for enthalpy of gelatinization of fish protein-sago starch model as a function of (a) minced fish and sago starch (b) minced fish and water ratios and (c) sago starch and water ratios based on Differential Scanning Calorimetry (DSC) measurement	79
8 Figure 8 (a, b and c): Effect of the ratios of the minced fish on $T_o$ , $T_p$ and $T_c$ temperatures of the fish protein - sago starch model	86
9 Figure 9 (a, b and c): Effect of the ratios of sago starch on $T_o$ , $T_p$ and $T_c$ temperatures of the fish protein - sago starch model	87
10 Response surface plot for onset temperature of gelatinization of fish protein - sago starch model as a function of (a) minced fish and sago starch ratios, (b) minced fish and water ratios and (c) sago starch and water ratios based on rheometer measurements	93



11	Response surface plot for peak temperature of gelatinization of fish protein - sago starch model as a function of (a) minced fish and sago starch ratios, (b) minced fish and water ratios and (c) sago starch and water ratios based on rheometer measurements	95
12	Response surface plots for storage modulus of fish protein - sago starch model as a function of (a) minced fish and sago starch ratios, (b) minced fish and water ratios and (c) sago starch and water ratios based on rheometer measurements	97
13	Response surface plot for loss modulus of fish protein-sago starch model as a function of (a) minced fish and sago starch ratios, (b) minced fish and water ratios and (c) sago starch and water ratios based on rheometer measurements	99
14	Overall optimization plot for the independents and responses	101
15	Scores distribution attributed to the texture of the local and optimized fish protein - starch models	104
16	Scores distribution attributed to the overall acceptability of the local and optimized fish protein - starch models	104
17	Thermograms of the effects of selected sodium chloride concentrations (0 - 3.0%) on gelation temperature of fish protein - sago starch model measured using Differential Scanning Calorimetry (DSC)	115
18	Thermograms of the effects of selected sucrose concentrations (0 - 3.0%) on gelation temperature of fish protein-sago starch model measured using Differential Scanning Calorimetry (DSC)	121
19	Geometry of the fish protein - sago starch model used for simulated heat cooking time	127
20	Effect of sodium chloride and sucrose concentrations on thermal conductivity of the fish protein-sago starch model during deep - fat frying	132
21	Effect of sodium chloride and sucrose concentrations on specific heat capacity of the fish protein - sago starch model during deep fat frying	133
22	Effect of sodium chloride and sucrose concentrations on density of the fish protein - sago starch model during deep - fat frying	135
23	Effect of sodium chloride and sucrose concentrations on thermal diffusivity of the fish protein - sago starch model during deep - fat frying	137



24	Temperature histories of the center and superficial points of control sample of the fish protein - sago starch model during deep - fat frying	139
----	---	-----



## LIST OF ABBREVIATIONS

MF	Minced fish
SS	Sago starch
W	Water
$T_o$	Onset temperature of gelatinization
$T_p$	Peak temperature of gelatinization
$T_c$	Conclusion temperature of gelatinization
$G'$	Storage modulus
$G''$	Loss modulus
RSM	Response surface methodology
DSC	Differential scanning calorimetry
$\Delta H$	Enthalpy
J/g	Joule per gram
Pa	Pascal
kPa	Kilo Pascal
NaCl	Sodium chloride
3D	Three dimensional plots
$R^2$	Regression coefficient
$\alpha$	Thermal diffusivity
$c_p$	Specific heat capacity
$\rho$	density
$k$	thermal conductivity



$t$	time
$T$	temperature
s	second



## TABLE OF CONTENTS

	<b>Page</b>
<b>DEDICATION</b>	ii
<b>ABSTRACT</b>	iii
<b>ABSTRAK</b>	vi
<b>ACKNOWLEDGMENTS</b>	ix
<b>APPROVAL</b>	xi
<b>DECLARATION</b>	xiii
<b>LIST OF TABLES</b>	xiv
<b>LIST OF FIGURES</b>	xvi
<b>LIST OF ABBREVIATIONS</b>	xix
<b>CHAPTER</b>	
<b>I      GENERAL INTRODUCTION</b>	<b>1</b>
<b>II     LITERATURE REVIEW</b>	<b>3</b>
Thermal and rheological properties of different food systems	3
Effect of protein	3
Effect of starch	12
Effect of protein - starch mixtures	21
Gelatinization of starch in food system	31
<i>Keropok lekor</i> formulation in Malaysia	34
Optimization of food components using response surface methodology	35
Effect of some food ingredients on thermal and rheological behaviour of food systems	40
Effect of sugars	40
Effect of salts	44
Effect of different cooking methods on physicochemical and thermal Properties of food models	47
Heat conduction equation	52
Empirical equation (Ansari's equation)	53
<b>III    PHYSICOCHEMICAL, SENSORY PROPERTIES AND OPTIMIZATION OF THE FISH PROTEIN - SAGO STARCH MODEL</b>	
Introduction	55
Materials and methods	57
Raw materials	57
Preparation of the minced fish	57
Compositional analysis of the minced fish and sago starch	
Experimental design	58



Preparation of dough for the fish protein - sago starch system	59
Preparation of samples of the fish protein - sago starch system for sensory evaluation	60
Thermal measurements	60
Texture profile analysis of the formulated fish protein - sago starch system	61
Sensory evaluation of the formulated fish protein - sago starch system	62
Rheological measurements	62
Optimization procedure	63
Texture profile analysis of the optimized fish protein - sago starch model	64
Sensory evaluation of the optimized fish protein - sago starch model	64
Statistical analysis	64
Results and discussion	66
Compositional analysis	66
Thermal measurement	67
Texture profile analysis of the formulated fish protein - sago starch system	80
Sensory evaluation of the formulated fish protein - sago starch system	83
Rheological measurements	88
Optimization procedure	100
Texture profile analysis of the optimized fish protein - sago starch model	101
Sensory evaluation of the optimized and local fish protein - starch models	103
Conclusion	105
<b>IV</b>	
<b>EFFECT OF ADDING SODIUM CHLORIDE AND SUCROSE ON GELATION PROPERTIES OF THE PROTEIN-STARCH MODEL</b>	<b>107</b>
Introduction	107
Materials and methods	108
Materials	108
Methods	108
Samples preparation	109
Thermal measurements	109
Rheological measurements	110
Statistical analysis	110
Results and discussion	111
Effect of sodium chloride concentrations on transition temperatures and rheological properties of the systeml	111
Effect of sucrose concentrations on transition temperatures and rheological properties of the system	117
Conclusion	122



<b>V</b>	<b>THE EFFECT OF SODIUM CHLORIDE AND SUCROSE CONCENTRATIONS USED IN FISH PROTEIN - SAGO MODEL ON COOKING TIME ESTIMATION</b>	124
	Introduction	124
	Materials and methods	125
	Materials	125
	Sample preparation	125
	Measurement of center temperature of the samples	126
	Calculation of thermal conductivity, specific heat capacity, density, thermal diffusivity and cooking time of the samples	127
	Thermal conductivity determination	127
	Specific heat capacity determination	128
	Density determination	128
	Thermal diffusivity determination	129
	Cooking time calculation	129
	Results and discussion	130
	Thermal conductivity determination	130
	Specific heat capacity determination	132
	Density determination	134
	Thermal diffusivity determination	135
	Cooking time calculation	137
	Conclusion	139
<b>VI</b>	<b>GENERAL CONCLUSION AND RECOMMENDATIONS</b>	141
	<b>BIBLIOGRAPHY</b>	144
	<b>APPENDICES</b>	164
	<b>BIODATA OF STUDENT</b>	182

