




**UNIVERSITI PUTRA MALAYSIA**

**BIOASSAY-GUIDED IDENTIFICATION OF ANTI-CHOLINESTERASE AND  
ICHTHYOTOXIC COMPOUNDS FROM *WALSURA CHRYSOGYNE* (MIQ.)  
BAKH.F.APUD VAN STEENIS**

**ILYA IRYANI MAHMUD**

**FS 2012 9**

**BIOASSAY-GUIDED IDENTIFICATION OF ANTI-CHOLINESTERASE  
AND ICHTHYOTOXIC COMPOUNDS FROM *WALSURA CHRYSOGYNE*  
(MIQ.) BAKH.F.APUD VAN STEENIS**




**ILYA IRYANI MAHMUD**

**MASTER OF SCIENCE  
UNIVERSITI PUTRA MALAYSIA**

**2012**

**BIOASSAY-GUIDED IDENTIFICATION OF ANTI-CHOLINESTERASE  
AND ICHTHYOTOXIC COMPOUNDS FROM *WALSURA  
CHRYSOGYNE* (MIQ.) BAKH.F.APUD VAN STEENIS**



By

**ILYA IRYANI MAHMUD**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfillment of the Requirements for the Degree of Master of Science**

**September 2012**

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment  
of the requirement for the degree of Master of Science

**BIOASSAY-GUIDED IDENTIFICATION OF ANTI-CHOLINESTERASE  
AND ICHTHYOTOXIC COMPOUNDS FROM *WALSURA CHRYSOGYNE*  
(MIQ.) BAKH.F.APUD VAN STEENIS**

By

**ILYA IRYANI MAHMUD**

**September 2012**

**Chairman : Intan Safinar Ismail, PhD**

**Faculty : Faculty of Science**

Phytochemical studies on *Walsura chryso gyne* leaves and barks have resulted in the isolation of ten compounds including two new compounds which namely chrysur a (39) and 20*R*,24*S*-epoxy-25-hydroxydammaran-3-one (43). The other eight compounds are cycloart-23-ene-3 $\alpha$ , 25-diol (35), dymalol (36), 25-methoxycycloart-23-en-3 $\beta$ -ol (37), eichlerianic acid (38), foveolin A (40), viridiflorol (41), cabraleadiol (42) and methyl eichlerianate (44). The structures of the isolated compounds were elucidated using spectroscopic techniques including Nuclear Magnetic Resonance (NMR), Mass Spectrometry (MS), Fourier Transform Infrared (FT-IR), *X-ray* crystallography and by comparison with the reported data from literatures.

Based on a bioautographic Thin Layer Chromatography (TLC) profile of anti-acetylcholinesterase activity and ichthyotoxic assay, ethyl acetate fraction from leaves and hexane fraction from bark were further fractionated by several chromatographic techniques. The EtOAc-soluble part from leaves, which gave the strongest activity in ichthyotoxicity and TLC bioautography of acetylcholinesterase enzyme assay, was further fractionated and resulted in purification of six triterpenes (**35**, **36**, **37**, **38**, **39** and **40**). Repeated chromatography of the hexane bark fraction on normal phase silica gel led to the purification of a sesquiterpene (**41**) and three triterpenes (**42**, **43**, and **44**).

All of the isolated compounds from leaves and barks, except for **35** and **37** which amounts are too small, were then subjected to ichthyotoxic assay at a concentration of 40 ppm each. Only eichlerianic acid (**38**) and viridiflorol (**41**) exhibited strong toxicity against zebrafish (*Danio rerio*). The median tolerance limit ( $TL_M = IC_{50}$ ) for **39** and **41** was evaluated by straight line graphical interpolation and resulted in the values of 6.7 and 15 ppm respectively. Compound **36**, **38**, **39** and **41** showed positive result on TLC-bioautography with fast blue B salt. However, the obtained  $IC_{50}$  for them by using Ellman's method was not significant ( $>200 \mu\text{g/ml}$ ).

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai  
memenuhi keperluan untuk ijazah Master Sains

**PENGENALPASTIAN SEBATIAN BERPANDUKAN BIO CERAKINAN  
BERDASARKAN ANTI-CHOLINESTERASE DAN ICHTHYOTOXIC  
DARIPADA *WALSURA CHRYSOGYNE* (MIQ.) BAKH.F.APUD VAN  
STEENIS**

Oleh

**ILYA IRYANI MAHMUD**

**September 2012**

**Pengerusi : Intan Safinar Ismail, PhD**

**Fakulti : Fakulti Sains**

Kajian kimia terhadap daun dan kulit batang *Walsura chrysogyne* telah menghasilkan sepuluh sebatian termasuk dua sebatian baru yang dinamakan chrysura (**39**) 20*R*,24*S*-epoxy-25-hydroxydammaran-3-one (**43**). Lapan sebatian lain ialah cycloart-23-ene-3 $\beta$ , 25-diol (**35**), dymalol (**36**), 25-methoxycycloart-23-en-3 $\beta$ -ol (**37**), eichlerianic acid (**38**), foveolin A (**40**), viridiflorol (**41**), cabraleadiol (**42**) dan methyl eichlerianate (**44**). Struktur sebatian yang diasingkan telah dikenalpasti dengan menggunakan teknik spektroskopik termasuk, spektroskopi resonans magnet nukleus (NMR), spektrometri jisim (MS), spektroskopi inframerah (FT-IR), *X-ray* kristalografi dan berdasarkan kepada perbandingan dengan penemuan data yang terdahulu.

Berdasarkan profil bioautografik kromatografi lapisan nipis (TLC) bagi aktiviti perencatan acetylcholinesterase dan ichthyotoxic, pecahan etil asetat daripada daun dan pecahan heksana daripada kulit batang dipisahkan seterusnya oleh beberapa teknik kromatografi. Bahagian etil asetat daripada daun yang memberikan aktiviti tertinggi dalam ichthyotoxic dan bioautografi TLC, enzim acetylcholinesterase telah dipisahkan dan telah menghasilkan enam triterpena (**39**, **40**, **41**, **42**, **43** and **44**). Kromatografi diulangi dengan menggunakan fasa gel silica normal pada pecahan heksana kulit batang, membawa kepada penulenan sesquiterpena (**41**) dan tiga triterpena (**42**, **43**, and **44**).

Semua sebatian yang diasingkan daripada daun dan kulit batang, kecuali sebatian **35** dan **37** yang jumlahnya terlalu sedikit, telah diuji dengan ichthyotoxic pada kepekatan 40 ppm setiap satu. Hanya eichlerianic acid (**38**) dan viridiflorol (**41**) menunjukkan ketoksikan yang tinggi terhadap ikan zebra. Had toleransi median ( $TL_M = IC_{50}$ ) untuk **38** dan **41** telah dinilai dengan *Danio rerio* interpolasi garis lurus grafik dan masing-masing memberikan nilai  $TL_M$  6.7 dan 15 ppm. Sebatian **36**, **38**, **39** dan **41** menunjukkan keputusan positif pada bioautografi-TLC kromatografi lapisan dengan garam Fast Blue B. Walau bagaimanapun,  $IC_{50}$  yang diperolehi dengan cara Ellman adalah tidak ketara ( $>200 \mu\text{g/ml}$ ).

## ACKNOWLEDGEMENTS

Alhamdulillah, thanks to Allah for the completion of my study and this thesis.

First and foremost, I would like to convey my sincere and deepest appreciation to my supervisor, Dr. Intan Safinar Ismail who has enlighten me to appreciate the pursuit of knowledge as a reward in itself. She was always available for consultation with her invaluable comments and constructive criticism. I have learned so much from her. Also appreciations to Dr. Emilia who taught me structural modification and for making arrangement for me to use the chemical and also instrument in her laboratory. A special thanks also goes to Prof. Dr. Hoshi Morita for kindly being in my supervisory committee and also for the NMR and MS analysis. Thank you to all of you.

I must also thank the students and staff in the Department of Chemistry, Faculty of Science and also all the members of the Natural Products Laboratory, Institute of Bioscience for their advice, help and moral support. I am particularly grateful to Dr. Ibrahim and Wong for helping in preparation of *X-ray* crystallography data and paper.


My special thanks also goes to my only lab mate in Faculty of Science, Nawwar and my research group members, Izwan, Kok Wah and Wei Har for their guidance,



concern, understanding and support. To my best friends; Mie, Wani and Ngoh, thank you very much for always being my pillars of strength throughout my study.

Last but not least, to my beloved parents (ma and abah) and sisters (Na and Adik), for their prayers, unremitting love, encouragement, inspiration and continuous support which inspired me to accomplish this research.

‘MAY GOD BLESS YOU ALL’



I certify that an Examination Committee has meet on 27<sup>th</sup> September 2012 to conduct the final examination of Ilya Iryani Mahmud on her Master of Science thesis entitled 'Bioassay-Guided Identification of Anti-Cholinesterase and Ichthyotoxicity of *Walsura chrysogyne*' in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the student be awarded the degree of Master of Science.

Members of the Examination Committee were as follows:

**GWENDOLINE EE CHANG LIAN, PhD**

Professor  
Department of Chemistry  
Faculty of Science,  
Universiti Putra Malaysia  
(Chairman)

**KHOZIRAH SHAARI, PhD**

Professor  
Department of Chemistry  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

**SITI MARIAM MOHD NOR, PhD**

Lecturer  
Department of Chemistry  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

**JALIFAH LATIP, PhD**

Associate Professor  
Department of Chemistry and Food Technology  
Faculty of Science and Technology  
Universiti Kebangsaan Malaysia  
(External Examiner)

---

**ZULKARNAIN ZAINAL, PhD**

Professor and Deputy Dean  
School of Graduate Studies

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Intan Safinar Ismail, PhD**

Lecturer  
Institute of Bioscience  
Universiti Putra Malaysia  
(Chairman)

**Emilia Abd Malek, PhD**

Lecturer  
Faculty of Science,  
Universiti Putra Malaysia  
(Member)

**Hiroshi Morita, PhD**

Professor  
Faculty of Pharmaceutical Science  
Hoshi University, Japan  
(Member)

---


**BUJANG BIN KIM HUAT, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## DECLARATION

I declare that the thesis is on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.



---

**ILYA IRYANI MAHMUD**

Date: 27 September 2012



## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	I
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	vi
<b>APPROVAL</b>	viii
<b>DECLARATION</b>	Ix
<b>LIST OF TABLES</b>	xii
<b>LIST OF CHARTS</b>	xiii
<b>LIST OF FIGURES</b>	xiv
<b>LIST OF ABBREVIATIONS</b>	xviii
<b>CHAPTER</b>	
<b>1</b>	
<b>INTRODUCTION</b>	
1.1 General Introduction	1
1.2 Ichthyotoxic Properties of Plants	1
1.3 Cholinesterase Inhibitor from Plants	2
1.4 Objective of Research	3
<b>2</b>	
<b>LITERATURE REVIEW</b>	
2.1 Meliaceae Family	4
2.2 <i>Walsura</i> Genus	4
2.3 <i>Walsura chrysogyne</i> species	8
2.4 Ichthyotoxic Compounds in Plants	10
2.4.1 Zebrafish ( <i>Danio rerio</i> )	12
2.5 Anti-cholinesterase inhibitors	14
2.5.1 Acetylcholinesterase inhibitors	14
<b>3</b>	
<b>EXPERIMENTAL</b>	
3.1 General Instrumentation	17
3.2 Chromatographic Methods	17
3.3 Solvents	19
3.4 Plant Material	19
3.5 Extraction and Fractionation of <i>W. chrysogyne</i>	19
3.5.1 Isolation and Identification of Compounds from the Leaves of <i>W. chrysogyne</i>	21
3.5.2 Isolation and Identification of Compounds from the	25

	Barks of <i>W. chrysogyne</i>	
3.6	Ichthyotoxic Assay	31
3.7	Anti-acetylcholinesterase Assay	32
3.7.1	TLC Bioautography with Fast Blue B Salt	32
3.7.2	96 well plate (Ellman's Method)–Ellman <i>et al.</i> , 1976	34
3.8	Modification of Chrysurin ( <b>39</b> )	36
3.8.1	Reaction with bromoacetylchloride	36
3.8.2	Base hydrolysis of chrysurin ( <b>39</b> )	37
<b>4</b>	<b>RESULTS AND DISCUSSION</b>	
4.1	Phytochemical studies of <i>W. chrysogyne</i> leaves	43
4.2	Phytochemical studies of <i>W. chrysogyne</i> barks	44
4.3	Ichthyotoxic fraction and substance	44
4.4	Anti-acetylcholinesterase (AChE) assay	47
4.5	Isolated compounds from <i>W. chrysogyne</i> leaves and barks	49
4.5.1	Dammarane type	49
4.5.1.1	Modification of chrysurin	64
4.5.2	Aromadendrane type	99
4.5.3	Cycloartane type	103
4.6	Essential oils component of <i>W. chrysogyne</i> leaves	111
<b>5</b>	<b>CONCLUSION AND SUGGESTIONS</b>	113
	<b>BIBLIOGRAPHY</b>	115
	<b>BIODATA OF STUDENT</b>	119
	<b>PUBLICATION/ PROCEEDING</b>	120