

Natural Product Compounds from *Calophyllum depressinervosum*

Nor Hisam Zamakshshari, Gwendoline Cheng Lian Ee*, Soek Sin Teh,
Shaari Daud, Thiruventhan Karunakaran and Intan Safinar

Department of Chemistry, Faculty of Science, Universiti Putra Malaysia, 43400 Serdang, Selangor,
Malaysia

ABSTRACT

Our detailed study on the phytochemistry of the stem bark of *Calophyllum depressinervosum* resulted in the isolation of four xanthenes and one coumarin. The xanthenes are trapezifolixanthone (1), macluraxanthone (2), ananixanthone (3), caloxanthone C (4) and the coumarin calanolide E2 (5). The structures of these compounds were elucidated using spectroscopic analysis such as 1D and 2D-NMR, GCMS, IR and UV .

Keywords: Natural product compounds, *Calophyllum depressinervosum*

INTRODUCTION

The genus *Calophyllum* comprises 180-200 tree species, which are distributed in the tropical rain forest with some occurring in Malaysia (Cechinel et al, 2009). *Calophyllum depressinervosum* species is one species from this genus which grows abundantly in Malaysia. This species is also known as *Bintagor lekok* by local Malaysians (Whitmore

et al., 1973). Previous phytochemical studies have shown that the *Calophyllum* genus is a valuable source of secondary metabolites such as xanthenes, coumarins chromenes and flavonoids (Ee et al., 2006). These secondary metabolites have also been shown to give good bioactivities against the HIV virus and they possess good anti-proliferative activities against cancer cell lines (Mah et al., 2015). Many *Calophyllum* species are economically important for the timber industry especially for housing, shipbuilding, furniture, etc. (Sarangwood et al., 2009). This paper reports detailed structural elucidation of trapezifolixanthone (1) and spectroscopic data for macluraxanthone (2), anixanthone (3), caloxanthone C (4) and calanolide E2 (5).

ARTICLE INFO

Article history:

Received: 20 July 2015

Accepted: 28 September 2015

E-mail addresses:

shamcute87@gmail.com (Nor Hisam Zamakshshari),

gwen@upm.edu.my (Gwendoline Cheng Lian Ee),

judith_teh@hotmail.com (Soek Sin Teh),

shaari111@pahang.uitm.edu.my (Shaari Daud),

thiruventhan_90@yahoo.com (Thiruventhan Karunakaran),

safinar@upm.edu.my (Intan Safinar)

* Corresponding author

EXPERIMENTAL

Plant Material

The stem bark of *Calophyllum depressinervosum* was collected from the Sri Aman district in Sarawak, Malaysia, and identified by Associate Professor Dr. Rusea Go from the Biology Department, Universiti Putra Malaysia. A voucher specimen was deposited in the herbarium of Biology Department, Faculty of Science, Universiti Putra Malaysia.

General

Infrared spectra were measured using universal attenuated total reflection (UATR) technique on Perkin-Elmer 100 Series FT-IR spectrometer. EIMS were recorded on a Shimadzu GCMS-QP 5050A spectrometer (column, SGE BPX5 30meter x 0.25 mm I.D x 0.25 μ m film thickness, temperature, 200°C). The NMR spectra were obtained using a JEOL 500MHz FTNMR spectrometer using CDCl₃ as a solvent and tetramethylsilane (TMS) as internal standard. The UV spectra were recorded in EtOH on a shimadzu UV-160A, UV-Visible Recording Spectrophotometer. The melting points were obtained on a Leica Galen III instrument.

Extraction and Isolation

The dry and powdered stem bark of *Calophyllum depressinervosum* (2.1kg) was extracted three times by soaking in hexane at room temperature for 72 hours. The same procedure was repeated for another solvent, i.e. dichloromethane. Each extract was

dried under reduced pressure using a rotary evaporator to obtain the hexane (27.7g) and dichloromethane (26.8g) extracts. These extracts were chromatographed in a silica gel glass column under vacuum using a stepwise gradient system (hexane/dichloromethane, dichloromethane/ethyl acetate and ethyl acetate/methanol). Further purifications of the hexane extract using another silica gel column (gravity) gave trapezifolixanthone (1) (98.9mg), macluraxanthone (2) (157.6mg), ananixanthone (3) (90.7mg) and calanolide E2 (5) (19.4mg). Meanwhile, further purifications on the dichloromethane extract using gravity silica gel column chromatography gave caloxanthone C (4) (16mg).

Trapezifolixanthone (1). Yellow needle crystal; m.p 161-163°C (literature 140-142°C, Daud et al., 2014). UV λ_{max} : 3196, 1629, 1575, 1127 (EtOH) nm : 287, 260, 241, 210 EI-MS m/z: 378, 363, 154. ¹H NMR (500MHz, CDCl₃) and ¹³C NMR (125MHz, CDCl₃) (Table 1).

Macluraxanthone (2). Yellow needle crystal; m.p 170-174°C (literature 170-172°C, Iinuma et al., 1994). UV λ_{max} : 3417, 2971, 1582, 1193 (EtOH) nm : 346, 295, 251, 204 EI-MS m/z: 394, 379, 339, 162. ¹H NMR (500MHz, CDCl₃): δ 13.89 (s, 1-OH), δ 7.57 (d, 1H, *J*=8.05Hz, H-8), δ 6.97 (d, 1H, *J*=8.05Hz, H-7), δ 6.67 (d, 1H, *J*=9.10Hz, H-10), δ 6.50 (dd, 1H, *J*=10.8&17.2Hz, H-2'), δ 5.69 (d, 1H, *J*=9.10Hz, H-11), δ 5.02 (d, 1H, *J*=17.2Hz, H-3b'), δ 4.86 (d, 1H, *J*=10.8Hz, H-3a'), δ 1.71 (s, 6H, H-4'&H-5'), δ 1.46 (s, 6H, H-13&H-14). ¹³C NMR (125 MHz, CDCl₃) δ 180.9 (C-9),

δ 158.6 (C-1), δ 156.1 (C-4a), δ 155.2 (C-3), δ 151.7 (C-2'), δ 151.1 (C-5), δ 145.9 (C-5a), δ 132.9 (C-6), δ 127.5 (C-11), δ 116.2 (C-8), δ 115.5 (C-10), δ 113.5 (C-8a), δ 113.4 (C-4), δ 112.9 (C-7), δ 106.6 (C-3'), δ 104.8 (C-2), δ 102.8 (C-9a), δ 78.2 (C-12), δ 41.0 (C-1'), δ 29.2 (C-4'&C-5'), δ 27.2 (C-13&14).

Anixanthone (3). Yellow needle crystal; m.p 168-170°C (literature 170-171°C, Joaquim et al., 1998). UV λ_{\max} : 3217, 2918, 1573 (EtOH) nm : 302, 283, 261 EI-MS m/z: 378, 363, 335, 154. ^1H NMR (500MHz, CDCl_3): δ 13.22 (s, 1-OH), δ 7.78 (d, 1H, $J=8.02$ Hz, H-8), δ 7.30 (d, 1H, $J=8.02$ Hz, H-6), δ 7.23 (t, 1H, $J=8.02$, H-7), δ 6.79 (d, 1H, $J=10.3$ Hz, H-10), δ 5.65 (d, 1H, $J=10.3$ Hz, H-11), δ 5.25 (t, 1H, $J=6.87$ Hz, H-2'), δ 3.36 (d, 2H, $J=6.87$ Hz, H-1'), δ 1.81 (s, 3H, H-4'), δ 1.68 (s, 3H, H-5'), δ 1.49 (s, 6H, H-13&H-14). ^{13}C NMR (125 MHz, CDCl_3) δ 180.8 (C-9), δ 160.6 (C-1), δ 158.6 (C-3), δ 149.3 (C-4a), δ 144.3 (C-8a), δ 144.1 (C-10a), δ 131.7 (C-3'), δ 127.5 (C-11), δ 124.0 (C-7), δ 121.9 (C-2'), δ 121.2 (C-5), δ 120.1 (C-6), δ 117.2 (C-8), δ 115.0 (C-10), δ 112.3 (C-2), δ 103.2 (C-9a), δ 100.7 (C-4), δ 78.1 (C-12), δ 28.2 (C-14&C-13), δ 25.8 (C-5'), δ 21.3 (C-1'), δ 18.0 (C-4').

Caloxanthone C (4). Yellow needle crystal; m.p 211-213°C (literature 217°C, Iinuma et al., 1994). UV λ_{\max} : 3436, 2935, 1600, 1594, (EtOH) nm : 387, 294, 284, 233 EI-MS m/z: 378, 363, 335, 154. ^1H NMR (500MHz, CDCl_3): δ 13.42 (s, 1-OH), δ 7.72 (d, 1H, $J=9.15$ Hz, H-8), δ 7.23 (t, 1H, $J=9.15$ Hz, H-7), δ 7.20 (d, 1H, $J=9.15$ Hz, H-6), δ 6.78 (d, 1H, $J=10.3$ Hz, H-10), δ 6.77 (dd, 1H, $J=10.3$ & 17.2 Hz, H-2'), δ 5.63

(d, 1H, $J=10.3$ Hz, H-11), δ 5.24 (d, 1H, $J=17.2$ Hz, H-3b'), δ 5.07 (d, 1H, $J=10.3$ Hz, H-3a'), δ 1.64 (s, 6H, H-4'&H-5'), δ 1.51 (s, 6H, H-13&H-14). ^{13}C NMR (125 MHz, CDCl_3) δ 181.4 (C-9), δ 159.4 (C-3), δ 156.7 (C-1), δ 155.8 (C-2'), δ 154.0 (C-4a), δ 145.4 (C-5), δ 144.2 (C-5a), δ 127.4 (C-11), δ 124.2 (C-7), δ 120.5 (C-8a), δ 119.7 (C-6), δ 116.1 (C-8), δ 116.0 (C-10), δ 113.2 (C-4), δ 105.6 (C-2), δ 104.1 (C-3'), δ 103.6 (C-9a), δ 78.4 (C-12), δ 41.4 (C-1'), δ 28.3 (C-4'&C-5'), δ 28.0 (C-13&C-14).

Calanolide E2 (5). Yellowish oil. UV λ_{\max} : 2952, 1624, 1128 (EtOH) nm : 311, 297, 281, 240 EI-MS m/z: 388, 329, 271, 215, 107. ^1H NMR (500MHz, CDCl_3): δ 12.37 (s, 7-OH), δ 6.59 (d, 1H, $J=10.31$ Hz, H-9), δ 5.45 (d, 1H, $J=10.31$ Hz, H-10), δ 4.50 (m, 1H, H-3'), δ 3.68 (m, 1H, H-4), δ 2.84 (dd, 1H, $J=9.16$ & 14.89 Hz, H-3a), δ 2.67 (dd, 1H, $J=9.16$ & 14.89 Hz, H-3b), δ 2.53 (m, 1H, H-2'), δ 1.81 (m, 1H, H-14a), δ 1.46 (m, 1H, H-14b), δ 1.43 (s, 3H, H-12), δ 1.35 (s, 3H, H-13), δ 1.34 (d, 3H, $J=5.73$ Hz, H-4'), δ 1.15 (m, 2H, H-15), δ 1.12 (d, 3H, $J=6.87$ Hz, H-5'), δ 0.85 (t, 3H, $J=6.87$ Hz, H-16). ^{13}C NMR (125 MHz, CDCl_3) δ 201.1 (C-1'), δ 179.3 (C-2), δ 160.1 (C-8a), δ 159.9 (C-5), δ 157.3 (C-7), δ 125.7 (C-10), δ 115 (C-9), δ 108 (C-4a), δ 102.6 (C-6), δ 101.0 (C-8), δ 78.2 (C-11), δ 76.1 (C-3'), δ 44.2 (C-2'), δ 38.6 (C-3), δ 35.5 (C-14), δ 30.5 (C-4), δ 28.5 (C-13), δ 28.1 (C-12), δ 20.8 (C-15), δ 16.3 (C-4'), δ 14.0 (C-16), δ 9.3 (C-5').

RESULTS AND DISCUSSION

Trapezifolixanthone (1) (98.9 mg) was isolated as yellow needle crystals with

a melting point of 145-148°C (literature 140-142°C, Daud *et al.*, 2014). The EIMS spectrum showed a molecular ion peak at 378, which is consistent with the molecular formula $C_{23}H_{22}O_5$. The ion fragment peak at m/z 363 was due to the loss of a methyl group. The fragment ion peak at m/z 154 was due to the loss of a pyrano ring and the prenyl group that are attached to the xanthone skeleton.

The FTIR spectrum for compound 1 gives typical IR absorptions for xanthenes at 3196 cm^{-1} , 1629 cm^{-1} , 1575 cm^{-1} , and 1127 cm^{-1} . The strong absorptions at 3196 cm^{-1} and 1629 cm^{-1} were representative for hydroxyl and conjugated carbonyl stretching. Meanwhile, the absorption at 1575 cm^{-1} was due to the stretching of an aromatic group.

The ^1H NMR spectrum for compound 1 exhibited the presence of one chelated hydroxyl group at $\delta 13.05$ (OH-1). The presence of a 3-methylbut-2-enyl substituent in compound 1 was indicated by the ^1H NMR signals at $\delta 5.23$ (t, 1H, $J=5.73\text{Hz}$, H-2'), $\delta 3.49$ (d, 2H, $J=5.73\text{Hz}$, H-1'), $\delta 1.85$ (s, 3H, H-5') and $\delta 1.71$ (s, 3H, H-4'). In the COSY experiment, the nature of the allylic coupling systems within the prenyl moiety was clearly demonstrated. It showed COSY couplings between the olefinic proton C-2' and the benzylic proton of C-1'. The signal at $\delta 3.49$ (H-1') in the proton NMR showed a long range coupling with the carbon signal at $\delta 107.1$ (C-4), $\delta 153.8$ (C-4a) and $\delta 158.3$ (C-3) in the HMBC spectrum hence confirming the prenyl unit to be positioned at C-4.

The ^1H NMR spectrum for compound 1 also revealed the presence of a pyrano ring attached to the xanthone skeleton. The ring ^1H NMR spectrum shows a pair of ortho-coupled proton with a coupling constant value of 10.3Hz at $\delta 6.74$ and $\delta 5.61$ for H-11 and H-10. The HMBC spectrum shows a cross peak for the proton signal at $\delta 1.47$ (H-13 and H-14) with the carbon signal at $\delta 78.4$ (C-12). This indicates the two methyls to be directly bonded to the carbon at $\delta 78.4$ (C-12). The 2J correlation of the proton at $\delta 5.61$ (H-10) with the carbon signal at $\delta 104.8$ (C-2) suggests the pyrano ring to be attached to the carbons at $\delta 104.8$ (C-2) and $\delta 158.3$ (C-3).

The ^{13}C NMR and DEPT spectra for (1) exhibited 23 carbon signals, which consist of four methyls, one methylene, six methines and eleven quaternary carbons including one carbonyl signal at $\delta 181.1$ (C-9). The carbons at $\delta 181.1$ (C-9), $\delta 158.3$ (C-3), $\delta 156.1$ (C-1), $\delta 153.8$ (C-4a), $\delta 144.5$ (C-5a) and $\delta 144.3$ (C-5) were shifted downfield due to the electronegative element oxygen, making them more deshielded. The carbon at $\delta 181.1$ (C-9) peak is generally weak due to slow relaxation of the quaternary carbon and is highly deshielded.

The COSY spectrum for (1) shows protons coupled to each other, indicating the positions of adjacent protons. One doublet and triplet signals with coupling constant 8.02Hz were observed at $\delta 7.73$ (H-8) and $\delta 7.23$ (H-7) indicating the ortho-coupled proton in a benzene ring. The meta-coupled protons in the left benzene ring at $\delta 7.73$ (H-8) and $\delta 7.28$ (H-6) signal were assigned to




Figure 1. Structures of xanthenes and coumarin

their respective carbons at signals δ 116.9 (C-8) and δ 119.9 (C-6) via the HMQC spectrum.

Based on the information given by the 1D and 2D NMR data, compound 1 was identified as trapezifolixanthone previously isolated from *Calophyllum hosei* (Daud et al., 2014).

ACKNOWLEDGEMENTS

The authors would like to acknowledge financial support from UPM, under the RUGS research fund. The Sarawak Biodiversity Centre (SBC) is also acknowledged.

Table 1
 ^1H NMR (500 MHz, CDCl_3) and ^{13}C NMR (125 MHz, CDCl_3)

Position	δ_{H}	δ_{C}	HMBC, δ
1	13.05 (s, OH)	156.1	156.1 (C-1), 104.8 (C-2)
2		104.8	
3		158.3	
4		107.1	
4a		153.8	
5	5.74 (s, OH)	144.3	144.3 (C-5), 119.9 (-6)
5a		144.5	
6	7.28 (d, 1H, $J = 8.02\text{Hz}$)	119.9	144.3 (C-5), 116.9 (C-8)
7	7.23 (t, 1H, $J = 8.02\text{Hz}$)	124.0	144.3 (C-5), 120.9 (C-8a)
8	7.73 (d, 1H, $J = 8.02\text{Hz}$)	116.9	144.5 (C-5a), 119.9 (C-6)
8a		120.9	
9		181.1	
9a		103.1	
10	5.61 (d, 1H, $J = 10.3\text{Hz}$)	127.6	104.8 (C-2), 78.4 (C-12)
11	6.74 (d, 1H, $J = 10.3\text{Hz}$)	115.8	158.3 (C-3), 78.4 (C-12)
12		78.4	
13	1.47 (s, 6H)	28.4	127.6 (C-10), 78.4 (C-12)
14			28.4 (C-13&C-14)
1'	3.49 (d, 2H, $J = 5.73\text{Hz}$)	21.8	158.3 (C-3), 153.8 (C-4a), 131.7 (C-3'), 122.7 (C-2'), 107.1 (C-4)
2'	5.23 (t, 1H, $J = 5.73\text{Hz}$)	122.7	25.7 (C-4'), 18.0 (C-5')
3'		131.7	
4'	1.71 (s, 3H)	25.7	131.7(C-3'),122.7(C-2'), 18.0(C-5')
5'	1.85 (s, 3H)	18.0	131.7(C-3'),122.7(C-2'), 25.7(C-4')

REFERENCES

- Cechinel, F. V., Megre, S. C., & Niero, R. (2009). Chemical and pharmacological aspects of the genus *Calophyllum*, *Chemistry and Biodiversity*, 6(03), 313-327.
- Ee, G. C. L., Kua, A. S. M., Lim, C. K., Jong, V., & Lee, H. L. (2006). Inophyllin A, a new pyranoxanthone from *Calophyllum inophyllum* (Guttiferae), *Natural Product Research*, 20(5), 485-491.
- Daud, S., Ee, G. C. L., Malik, E., Teh, S. S., & See, I. (2014). A new coumarine from *Calophyllum hosei*. *Natural Product Research*, 49(19), 1534-1538.
- Joaquim, C. B., Mara, S. P. A., & Marcelino, S. N. (1998). A prenylated xanthone from the bark of *Symphonia globulifera*. *Phytochemistry*, 49(4), 1159-1160.
- Linuma, M., Tosa, H., Tanaka, T., & Yonemori, S. (1994a). Two new xanthone in the underground part of *Calophyllum inophyllum*. *Heterocycles*, 37(2), 833-838.
- Linuma M., Tosa H., Tanaka, T., & Yonemori, S. (1994). Two xanthones from root bark of *Calophyllum inophyllum*. *Phytochemistry*, 35(2), 527-532.

- Mah, S. H., Ee, G. C. L., Teh, S. S., & Mohd, A. S. (2015). *Calophyllum inophyllum* and *Calophyllum soulattri* source of anti-proliferative xanthone and their structure-activity relationships, *Natural Product Research*, 29(1), 98-101.
- Sarangwood, S., Somnuek, P., Duangchai, S., & Thawatchi, S. (2009). The genus *Calophyllum* (Guttiferae) in Thailand, *Journal of Forest Management*, 3(5), 1- 24.
- Whitmore, T. C. (1973). *Tree flora of Malaya: A manual for foresters (Vol 2)*. London: Logman Group Limited.

