

**OPTIMUM GROWTH PARAMETERS FOR BOTH  
INDOOR AND OUTDOOR PROPAGATION OF MICROALGAE,  
*CHLORELLA VULGARIS* AND *ISOCHRYSIS GALBANA***

**SITTI RAEHANAH BINTI MUHAMAD SHALEH**

**DOCTOR OF PHILOSOPHY  
UNIVERSITI PUTRA MALAYSIA**

**2004**

**OPTIMUM GROWTH PARAMETERS FOR BOTH  
INDOOR AND OUTDOOR PROPAGATION OF MICROALGAE,  
*CHLORELLA VULGARIS* AND *ISOCHRYSIS GALBANA***

**By**

**SITTI RAEHANAH BINTI MUHAMAD SHALEH**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy.**

**June 2004**

**Dedicated to**

Ayah, Mummy, Sharudin, Husna, Nadia & Sofea

Thank you for their true love, support, understanding and their non-stopped  
devotions for making everything come true.

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Doctor of Philosophy

**Optimum Growth Parameters for Both Indoor and Outdoor Propagation of Microalgae, *Chlorella vulgaris* and *Isochrysis galbana***

By

**SITTI RAEHANAH BINTI MUHAMAD SHALEH**

**June 2004**

**Chairmain : Hishamuddin Bin Omar, Ph.D**

**Faculty : Science**

*Chlorella vulgaris* and *Isochrysis galbana* were grown under different growth conditions both in indoor and outdoor cultures. Best culture conditions such as light intensities, photoperiods, salinities and nutrient concentrations were determined as well as the biochemical compositions. Indoor cultures of *C. vulgaris* and *I. galbana*, when continuously illuminated, grew optimally at  $100 \mu\text{molm}^{-2}\text{s}^{-1}$  and  $40 \mu\text{molm}^{-2}\text{s}^{-1}$ , respectively. However, under 12 hour light, *C. vulgaris* grew optimally at  $200 \mu\text{molm}^{-2}\text{s}^{-1}$ . The chlorophyll<sub>a</sub> concentrations in both algae were lower at higher light intensity and higher at lower light intensity. Total protein and total lipid contents of both species were also significantly higher ( $p < 0.05$ ) at lower light intensity. *I. galbana* contained more lipid than *C. vulgaris* in those conditions. In contrast to protein and lipid, carbohydrate content increased with increasing light intensity. Similar to *C. vulgaris*, the lowest carbohydrate content of *I. galbana* was obtained at  $10 \mu\text{molm}^{-2}\text{s}^{-1}$  conditions for both photoperiods. Although the protein contents of *C. vulgaris* varied under different light intensity and photoperiod, the relative

proportions of constituent amino acids were found to be almost similar. Total essential amino acids under continuous illumination was significantly higher ( $p < 0.05$ ) at the lowest and the highest intensities while in *I. galbana*, it was significantly higher ( $p < 0.05$ ) only at the lowest intensity. It was surprising that *C. vulgaris* contained high proportions of  $\omega$ -3 highly unsaturated fatty acids at continuous illuminations although both eicosapentaenoic acid (20:5 $\omega$ -3) and docosahexaenoic acid (22:6 $\omega$ -3) were absent at 12 hours light. EPA in *I. galbana*, was absent at 12 hours light, while the DHA was present in relatively smaller quantities than in continuous light.

In indoor conditions, at full media concentration, the best salinity for *C. vulgaris* was at 30 ppt and 25 ppt for *I. galbana*. *C. vulgaris* grown in full media concentration contained more chlorophyll<sub>a</sub> relative to that in half media concentration. For *I. galbana* grown in full media concentration, chlorophyll<sub>a</sub> increased as salinity increased. Protein contents decreased as salinity increase both at full and half strength media for *C. vulgaris* and *I. galbana*. The cellular lipid contents of both algae were significantly higher ( $p < 0.05$ ) in half media concentration. In full media concentration, total lipid contents in *C. vulgaris* decreased at higher salinity but in *I. galbana* no clear trend in lipid contents in relation to salinity observed. As salinity increased, higher carbohydrate contents were found in *C. vulgaris* in full media concentration, while in half media concentration an inverse trend was detected. *I. galbana* also revealed higher proportions of total carbohydrates in half media concentration compared to full media concentration. The amino acid profile of both algae exhibited minor differences at different salinity and media

concentrations. Most of the essential amino acid in *C. vulgaris* and *I. galbana* were significantly higher ( $p < 0.05$ ) at lower salinity. In *C. vulgaris*, the EPA was absent while only trace amount of DHA were detected in cells grown under full media concentration. However in *I. galbana* the amount of EPA was very high at higher salinity and in contrary the, DHA was higher at lower salinity.

The growth performances of *C. vulgaris* outdoor cultures were relatively better than *I. galbana* under all growth conditions studied. *C. vulgaris* grown both in unsheltered and sheltered areas demonstrated rapid increased in cell density after four days of culture with no lag phases observed. Unlike *C. vulgaris*, cultures of *I. galbana* grown under unshaded conditions demonstrated increase in cell numbers for 2 days but cells under sheltered place demonstrated no increased at all. The highest specific growth rate in *C. vulgaris* was in lower salinity at half media concentration while the maximum cell density was achieved by cells in full media concentration of the same salinity. For *I. galbana* cells density of half media concentration was higher then in full media concentration. The highest total protein content of *C. vulgaris* was obtained in culture under shaded conditions and generally total lipid and carbohydrate were found to be higher in unshaded conditions. However, it was unexplainable why less carbohydrate was found in older cells at both conditions. *I. galbana* was a fast growing algae and the protein as well as lipid contents were significantly higher ( $p < 0.05$ ) at sheltered condition while no significant changes to carbohydrate content was recorded. The carbohydrate content in *C. vulgaris* was higher in older cultures in full

and half media concentration while at early culture age, the lipid content was higher. In the early culture age of *I. galbana*, protein content was insignificantly affected ( $p>0.05$ ) by salinity and media concentration. However, total lipid and carbohydrate were found higher at 15 ppt compared to higher salinity in half media concentration.

In the bioeconomic study the production cost of *C. vulgaris* was found to be directly influenced by labour cost. From the production forecast, it was discovered that the high production cost was due to underutilized labour force which can be reduced by having bigger scale production. In order to break-even for one cycle of 100 L cultures, the minimum culture volume should be 1,902 litres.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**Parameter Pertumbuhan Optima untuk Propagasi Mikroalga  
*Chlorella vulgaris* dan *Isochrysis galbana* dalam Keadaan  
Terlindung dan Terdedah.**

Oleh

**SITTI RAEHANAH BINTI MUHAMAD SHALEH**

**Jun 2004**

**Pengerusi : Hishamuddin Bin Omar, Ph.D**

**Fakulti : Sains**

Pengkulturan di dalam makmal dan di luar makmal telah dijalankan ke atas *Chlorella vulgaris* dan *Isochrysis galbana*. Tindakbalas pertumbuhan unggul alga terhadap faktor cahaya, fotokala, saliniti air dan kepekatan nutrien dikenalpasti di samping komposisi biokimia. Dalam keadaan makmal iaitu pada cahaya berterusan, didapati intensiti cahaya optimum bagi *C. vulgaris* dan *I. galbana* adalah masing-masing pada paras  $100 \mu\text{molm}^{-2}\text{s}^{-1}$  dan  $40 \mu\text{molm}^{-2}\text{s}^{-1}$ . Manakala dalam keadaan 12 jam bergelap, kedua-dua mikroalga tumbuh secara optimum pada paras  $200 \mu\text{molm}^{-2}\text{s}^{-1}$ . Kepekatan klorofil<sub>a</sub> dalam kedua-dua alga juga serupa, iaitu rendah pada keadaan cahaya tinggi. Kandungan protien dan lipid kedua-duanya juga berbeza dengan ketara ( $p < 0.05$ ) iaitu tinggi dalam keadaan cahaya terhad. Namun *I. galbana* mengandungi lebih lipid berbanding *C. vulgaris*. Berbeza dengan protien dan lipid, karbohidrat meningkat dengan peningkatan intensiti cahaya. Seperti *C. vulgaris*, kandungan karbohidrat dalam *I. galbana* rendah pada cahaya terhad dalam kedua-dua fotokala. Sungguhpun kandungan



protien *C. vulgaris* berbeza pada intensiti cahaya dan fotokata berbeza, profil asid amino adalah hampir serupa. Dalam cahaya berterusan, jumlah asid amino perlu ketara lebih tinggi ( $p < 0.05$ ) pada cahaya terendah dan tertinggi, manakala perlakuan sama dalam *I. galbana* adalah pada cahaya yang rendah. *C. vulgaris* mengandungi asid lemak  $\omega$ -3 yang tinggi pada cahaya berterusan walaupun EPA dan DHA tiada pada keadaan 12 jam bergelap. Dalam keadaan 12 jam bergelap, EPA juga tidak didapati dalam *I. galbana* manakala DHA pula tinggi berbanding dalam cahaya berterusan.

Dalam kepekatan media penuh iaitu dalam kultur terkawal, saliniti unggul untuk *C. vulgaris* adalah 30 ppt dan bagi *I. galbana* pada 25 ppt. *C. vulgaris* dalam media berkepekatan penuh didapati mengandungi lebih klorofil<sub>a</sub> dan kepekatan klorofil<sub>a</sub> dalam *I. galbana* meningkat dengan peningkatan saliniti. Protien dalam *C. vulgaris* dan *I. galbana* menurun apabila saliniti air meningkat dalam kedua-dua media. Kandungan lipid kedua-dua alga ketara lebih tinggi ( $p < 0.05$ ) dalam media berkepekatan separuh. Dalam media berkepekatan penuh pula, lipid *C. vulgaris* menurun apabila saliniti meningkat manakala dalam *I. galbana* tiada corak perlakuan yang jelas terhadap kesan saliniti. Apabila saliniti meningkat, kandungan karbohidrat dalam *C. vulgaris* meningkat dalam media berkepekatan penuh manakala keadaan sebaliknya ditunjukkan dalam media berkepekatan separuh. *I. galbana* juga mengandungi karbohidrat yang tinggi dalam media berkepekatan separuh berbanding media berkepekatan penuh. Profil asid amino bagi kedua-dua mikroalga mengalami sedikit perbezaan terhadap kesan saliniti dan kepekatan media. Kebanyakan asid amino perlu ketara

lebih tinggi ( $p < 0.05$ ) dalam *C. vulgaris* dan *I. galbana* pada saliniti rendah. Dalam *C. vulgaris*, tiada EPA ditemui manakala DHA amat sedikit pada sel yang tumbuh dalam media berkepekatan penuh. Walau bagaimanapun, dalam *I. galbana*, EPA didapati sangat tinggi pada saliniti tinggi dan DHA pula secara songsang tinggi pada saliniti rendah.

Perlakuan pertumbuhan *C. vulgaris* di luar makmal yang tidak terkawal secara perbandingan dengan *I. galbana* adalah sangat memuaskan dalam semua keadaan kultur. *C. vulgaris* dalam cahaya semulajadi dan terlindung telah menunjukkan peningkatan kepadatan sel selepas empat hari pengkulturan. Manakala *I. galbana* pula menunjukkan peningkatan selama dua hari dan tiada sebarang peningkatan dalam kultur terlindung. Pertumbuhan spesifik tertinggi *C. vulgaris* adalah pada saliniti rendah iaitu dalam media separuh dan kepadatan sel maksimum direkodkan dalam media penuh. Sebaliknya kepadatan sel *I. galbana*, tinggi dalam media berkepekatan separuh. Kandungan protin tertinggi bagi *C. vulgaris* ditemui dalam kultur terlindung manakala lipid dan karbohidrat umumnya tinggi dalam keadaan cahaya semulajadi. Sesuatu yang tidak dapat dijelaskan mengenai karbohidrat yang berkurangan dengan peningkatan masa pengkulturan iaitu dalam sel-sel tua telah ditemui dalam *C. vulgaris*. Kandungan protin dan lipid dalam *I. galbana* direkodkan ketara lebih tinggi ( $p < 0.05$ ) dalam kultur terlindung. Dalam keadaan media berkepekatan penuh dan separuh, kandungan karbohidrat dalam sel-sel tua *C. vulgaris* juga lebih tinggi manakala sel-sel mudanya mengandungi lebih lipid. Protien pula dalam sel-sel muda *I. galbana* tidak menunjukkan perubahan ketara ( $p > 0.05$ )

terhadap kesan saliniti dan kepekatan media. Tetapi lipid dan karbohidrat didapati tinggi pada saliniti rendah dan lebih banyak tersintesis dalam keadaan media berkepekatan separuh.

Dalam kajian bioekonomi, kos pengeluaran *C. vulgaris* dipengaruhi secara langsung oleh kos buruh. Melalui unjuran pengeluaran, didapati bahawa kos pengeluaran tinggi yang disumbangkan oleh kos buruh dapat dikurangkan melalui penghasilan yang lebih tinggi dalam skala kultur yang lebih besar. Untuk mendapatkan semula modal pengeluaran untuk satu pusingan bagi setiap 100 L alga, isipadu minimum yang diperlukan ialah sebanyak 1,902 liter.

## ACKNOWLEDGEMENTS

BISMILLAHIRRAHMANIRAHIM, all thankful belong to ALLAH who makes everything possible.

First of all, I would like to express my very deep appreciation to Dr. Hishamuddin Bin Omar for his precious guidance, criticism, support and encouragement throughout the period of this study. My sincere appreciation is also extended to all the members of the Advisory Committee, Professor Dr. Abdul Manan Mat Jais, Ass. Professor Dr. Mohd. Salleh Kamarudin and Ass. Professor Dr. Mohd. Arief Syed for their constructive ideas and suggestions besides providing facilities to make the works running smoother. Not forgotten to extend my sincere gratefulness to Dr. Misri Kusnan and Prof. Dr. Hj. Fatimah Md. Yusoff for their valuable idea and allowing me to use all the facilities in Plant Physiology Laboratory and JICA-MASDEC Laboratory. Thousand of thanks to the anchorman of Biology Dept. Mr. Hidir, all COMAS staff especially Mr. Abdul Rahman, Mr. Sahari, Mr. Razali, Mr. Anuar and Mr. Kimon for helping me with no tired while running my algal cultures. Always in mind Mr. Zakaria from Nutrition Laboratory and all staff members of former Faculty of Fisheries and Marine Science for their valuable efforts in my works. Also my special appreciation to Mr. Ismail, Mr Hong Kock Seng and Mr. Mat Noor of Veterinary Dept., Petaling Jaya, for helping me in amino acid analysis and finally glade to record my thanks for everything to my fellow friends Annie C. (Ph.D.), Muta Harah (Ph.D.), Paymon (Ph.D.), Mr. Ahsan Habib (Ph.D.), Mr. Kennedy and all under-graduated students supervised by Dr. Hishamuddin Omar from 1995 to 2001.

I certify that an Examination Committee met on 15<sup>th</sup> June 2004 to conduct the final examination of Sitti Raehanah binti Muhamad Shaleh on her Doctor of Philosophy thesis entitled “Optimum Growth Parameters for Both Indoor and Outdoor Propagation of Microalgae, *Chlorella vulgaris* and *Isochrysis galbana*” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Siti Shapor Siraj, Ph.D.**

Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

**Fatimah Md. Yusoff, Ph.D.**

Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Member)

**Misri Kusnan, Ph.D.**

Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Member)

**Datin Ann Anton, Ph.D.**

Professor/Director  
Biotechnology Research Institute  
Universiti Malaysia Sabah  
(Independent Examiner)

---

**GULAM RUSUL RAHMAT ALI, Ph.D.**

Professor / Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

This Thesis submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfilment of the requirements for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

**Hishamuddin Bin Omar, Ph.D.**

Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

**Mohd. Arif Bin Syed, Ph.D.**

Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Member)

**Mohd. Salleh Bin Kamarudin, Ph.D.**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Abdul Manan Bin Mat. Jais, Ph.D.**

Professor, Dean  
School of Integrative Science  
Universiti College of Technology and Management Malaysia (KUTPM)  
(Member)

---

**AINI IDERIS, Ph.D.**

Prof.essor/ Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## **DECLARATION**

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

---

**SITTI RAEHANAH BINTI MUHAMAD SHALEH**

Date:

## TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vii
ACKNOWLEDGMENTS	xi
APPROVAL	xiii
DECLARATION	xiv
LIST OF FIGURES	xviii
LIST OF TABLES	xxiii
LIST OF ABBREVIATIONS	xxv
CHAPTER	
<b>1 GENERAL INTRODUCTION</b>	
1.1 Background of Study	1
1.2 Statement of Problems	7
1.3 Significance of the Study	10
1.4 Study Objectives	14
<b>2 LITERATURE REVIEW</b>	
2.1 Microalgal Nutritional Properties	18
2.1.1 Pigments	19
2.1.2 Protein	20
2.1.3 Lipid	20
2.1.4 Carbohydrate	21
2.1.5 Vitamins and Minerals	22
2.1.6 Bioactive Compounds	25
2.2 Cell Productivity and Responses To Growth Factors	26
2.2.1 Lights	26
2.2.2 Nutrients	28
2.2.3 Salinity	29
2.2.4 Temperature	31
2.2.5 pH	31
2.2.6 Other Factors	32
2.3 The Economic Feasibility	34
<b>3 GENERAL MATERIALS AND METHODS</b>	
3.1 Microalgal Cultivation	38
3.1.1 Cultivation Site	40
3.1.2 Culture Containers	40
3.1.3 Water Supply/Treatment	41
3.1.4 Water Salinity	42
3.1.5 Culture Illumination	42
3.1.6 Culture Temperature	44
3.1.7 Growth Media	45



3.1.8	Aeration/Water Turbulence	45
3.1.9	Microalgal Species	46
3.1.10	Production Culture	47
3.1.11	Cell Harvesting	49
3.1.12	Preservation And Storing.	53
3.1.13	Sanitation	53
3.2	Growth Measurements	53
3.3	Biochemical Composition Analyses	55
3.3.1	Chlorophyll <sub>la</sub> Determination	55
3.3.2	Total Protein Determination	56
3.3.3	Total Lipid Determination	57
	(i) Method 1 (Bligh & Dyer, 1959)	58
	(ii) Method 2 (Holland & Hannant, 1973)	59
3.3.4	Lipid Classes Separation	60
3.3.5	Total Carbohydrate Determination	62
3.3.6	Amino Acids Analyses	63
	(i) Acid Hydrolysis	64
	(ii) Performic Acid Oxidation	65
	(iii) Alkaline Hydrolysis	65
	(iv) Derivatisation	66
3.3.7	Fatty Acids Analyses	67
3.4	Data Analyses	69
<b>4</b>	<b>CULTURE OF <i>Chlorella vulgaris</i> AND <i>Isochrysis galbana</i> UNDER LABORATORY CONDITIONS</b>	
4.1	Introduction	70
4.2	Materials and Methods	75
4.2.1	Light Intensity and Photoperiod Study	75
4.2.2	Salinity and Growth Media Study	76
4.2.3	Biochemical Composition Analysis	78
4.3	Results	79
4.3.1	Light Intensity and Photoperiod Study	79
	(i) Cell Density	79
	(ii) Chlorophyll <sub>a</sub> Content	81
	(iii) Total Protein, Total Carbohydrate and Total Lipid Content	92
	(iv) Amino Acids Profile	94
	(v) Fatty Acids Profile	97
4.3.2	Salinity and Growth Media Study	108
	(i) Cell Density	108
	(ii) Chlorophyll <sub>a</sub> Content	113
	(iii) Total Protein, Total Carbohydrate and Total Lipid Content	120
	(iv) Amino Acids Profile	124
	(v) Fatty Acids Profile	129
4.4	Discussions	134
4.4.1	Light Intensity and Photoperiod Study	134
4.4.2	Salinity and Growth Media Study	141

<b>5</b>	<b>MICROALGAL OUTDOOR CULTURE FOR MASS PRODUCTION</b>	
	5.1 Introduction	146
	5.2 Materials and Methods	151
	5.3 Results	155
	5.3.1 Cell Density and Mass Production	155
	5.3.2 Total Protein, Total Carbohydrate and Total Lipid	160
	5.4 Discussions	168
<b>6</b>	<b>THE BIOECONOMIC STUDY OF MICROALGAL PRODUCTION IN OUTDOOR MASS CULTURE</b>	
	6.1 Introduction	173
	6.2 Materials and Methods	179
	6.3 Results	180
	6.3.1 Cell Density and Mass Production	180
	6.3.2 The Bioeconomic Considerations	180
	6.3.3 The Production Cost Estimation	185
	6.3.4 The Production Forecast	191
	6.3.5 Break-even Point	192
	6.5 Discussions	194
<b>7</b>	<b>GENERAL DISCUSSIONS, CONCLUSIONS AND RECOMMENDATIONS.</b>	
	7.1 General Discussions	198
	7.2 Conclusions	207
	7.3 Recommendations	209
	<b>BIBLIOGRAPHY</b>	212
	<b>APPENDIX</b>	232
	<b>BIODATA OF THE AUTHOR</b>	236

