



UNIVERSITI PUTRA MALAYSIA

***EFFECTS OF S-ADENOSYL-L-HOMOCYSTEINE AND TRICHOSTATIN A
ON DEVELOPMENTAL COMPETENCE, EPIGENETIC MODIFICATION,
AND GENE EXPRESSION IN CLONED CATTLE EMBRYOS***

SHAHRAM JAFARI

FP 2012 9

**EFFECTS OF S-ADENOSYL-L-HOMOCYSTEINE AND TRICHOSTATIN A ON
DEVELOPMENTAL COMPETENCE, EPIGENETIC MODIFICATION, AND
GENE EXPRESSION IN CLONED CATTLE EMBRYOS**



**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of
Philosophy**

August 2012

THIS THESIS IS DEDICATED TO
MY WIFE, MY MOTHER, MY DAUGHTER
WITH
LOVE AND GRATITUDE
AND ALSO TO
THE CHILDREN WHO HAVE THE ABILITY
BUT
NOT THE FACILITY AND OPPORTUNITY IN EDUCATION



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

EFFECTS OF S-ADENOSYL-L-HOMOCYSTEINE AND TRICHOSTATIN A ON DEVELOPMENTAL COMPETENCE, EPIGENETIC MODIFICATION, AND GENE EXPRESSION IN CLONED CATTLE EMBRYOS

By

SHAHRAM JAFARI

August 2012

Chairman: Assoc. Professor Halimatun bt Yaakub, PhD

Faculty: Agriculture

With regards to low efficiency of the somatic cell nuclear transfer (SCNT) procedure the hypothesis that epigenetic marks of somatic donor cells are responsible for this inefficiency is proposed. Epigenetic marks such as DNA methylation and histone acetylation are two major participants in nuclear reprogramming which can be used as indicators to assess SCNT efficiency. For this reason treating somatic cells or embryos with epigenetic drugs immediately after fusion or activation may improve the developmental rate of reconstructed embryos. Therefore, in this study we investigated the effect of S-adenosyl-L-homocysteine (SAH), a reversible inhibitor of DNA methyltransferases (DNMTs), at different intervals post SCNT on developmental competence, epigenetic status and gene expression of bovine cloned embryos. Treatment with 1mM SAH for 12 hours resulted in 54.6% blastocyst production which was significantly

higher ($P<0.05$) than *in vitro* fertilized (IVF) embryos (37.2%), cloned embryos treated with SAH for 72 hours (31.0%) and control cloned embryos (34.6%).

Intensity of DNA methylation in cloned embryos treated with SAH for 48 h resembled IVF and was significantly lower ($P<0.05$) than other SCNT groups. The histone H3 Lysine 9 (H3K9) acetylation levels of all SCNT groups were significantly lower ($P<0.05$) than the IVF group. The fluorescent intensity of EGFP-POU5F1 reporter gene at all intervals of SAH treatments, except for the 72 hours, was significantly higher ($P<0.05$) than non-treated SCNT embryos. Real-time analysis of gene expression in cloned blastocysts revealed significantly higher ($P<0.05$) expression of POU5F1 compared with IVF. There was no effect ($P>0.05$) of either embryo production method (SCNT vs. IVF), or among SAH treatment interval on the expression of BCL2 gene. On the contrary, treatment with SAH resulted in significant increase ($P<0.05$) in VEGF expression in comparison with IVF and control SCNT, except for cloned embryos treated with SAH for 24 hours. It was suggested that time interval of DNA methyltransferases (DNMTs) inhibition may have important consequences on the different features of bovine cloned embryos and the improving effects of DNMTs inhibition on developmental competency of cloned embryos is restricted to a specific period of time preceding *de novo* methylation stage. The effect of 0.05 μ M Trichostatin A (TSA), a histone deacetylase inhibitor, post-fusion for 12 h on the diverse aspects of developmental competency in SCNT embryos was investigated. The results of this study showed TSA treatment of SCNT embryos for 12 h after activation, significantly ($P<0.05$) improve blastocysts rate.

A significant reduction and increase ($P<0.05$) were observed in analyse of DNA methylation and histone acetylation levels respectively in the blastocyst stage of the TSA-treated group which is similar to IVF-derived blastocysts. Finally, examination of blastocysts on day 7 revealed the highest expression of OCT4 in the TSA-treated group ($P> 0.05$), while there were no significant changes observed in expressions of BCL2 and VEGF.

The results of this study indicated that post-fusion treatment with SAH has a time dependent effect on DNA-methylation and histone-acetylation, developmental competence and gene expression of the cloned embryos. The results of this study also showed that treatment of SCNT embryos with TSA after activation increased the capability of the embryos to develop to the blastocyst stage and TSA-treated cloned blastocysts are hypomethylated and hyperacetylated and higher relative expression of POU5F1 and VEGF genes. In addition, these results might improve quality of cloned bovine embryos to produce transgenic cattles.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

KESAN S-ADENOSIL-L-HOMOSISTEINA DAN TRIKOSTATIN A TERHADAP KECEKAPAN PERKEMBANGAN, MODIFIKASI EPIGENETIK DAN PENGEKSPRESAN GEN DARIPADA EMBRIO BOVIN TERKLON

Oleh

SHAHRAM JAFARI

Ogos 2012

Pengerusi : Profesor Madya Halimatum Yaakub, PhD

Fakulti : Pertanian

Berdasarkan pada kerendahan kecekapan prosedur pemindahan nukleus sel soma (SCNT), hipotesis yang menyatakan kebanyakan penanda epigenetik sel soma penderma bertanggungjawab terhadap tidak kecekapan ini dicadangkan. Penanda epigenetik seperti pemelilan DNA dan pengasetilan histon adalah dua komponen penting dalam program semula nukleus yang boleh digunakan sebagai penunjuk untuk menilai kecekapan SCNT. Untuk tujuan ini, merawat sel-sel soma atau embrio dengan bahan kimia epigenetik sejurus selepas penggabungan atau pengaktifan mungkin boleh meningkatkan kadar perkembangan embrio yang dibentuk semula.

Oleh itu, dalam kajian ini kami mengkaji kesan SAH, suatu perencat berbalik metiltransferase DNA (DNMTs), pada selang berbeza selepas SCNT terhadap kompetensi perkembangan, status epigenetik dan pengekspresan gen daripada

embrio bovin terklon. Rawatan dengan menggunakan 1 μ l SAH selama 12 jam menghasilkan blastosista sebanyak 54.6% dimana ia lebih tinggi ($P<0.05$) berbanding embrio yang disenyawakan secara *in vitro* (IVF: 37.2%), embrio terklon yang dirawat dengan SAH selama 72 jam (31.0%) dan embrio terklon kawalan (34.6%).

Intensiti pemetilan DNA dalam embrio terklon yang dirawat dengan SAH selama 48 jam menyerupai IVF dan lebih rendah ($P<0.05$) berbanding kumpulan SCNT yang lain. Tahap pengasetilan H3K9 (H3K9-Ac) dari semua kumpulan SCNT nyata sekali lebih rendah ($P<0.05$) daripada kumpulan IVF. Intensiti *fluorescent* (berpendarfluor) gen pelapor EGFP-POU5F1 pada semua selang rawatan SAH, kecuali 72 jam, nyata sekali lebih tinggi ($P<0.05$) berbanding kumpulan kawalan. Analisis masa nyata dan pengekspresan gen dalam blastosista yang diklon menunjukkan ekspresi POU5F1 adalah lebih tinggi ($P<0.05$) berbanding IVF. Tidak terdapat kesan ($P>0.05$) terhadap kaedah pengeluaran embrio (SCNT vs IVF) atau selang masa rawatan SAH ke atas ekspresi gen BCL-2.

Tambahan pula, rawatan menggunakan SAH tidak menunjukkan peningkatan signifikan ($P>0.05$) dalam pengekspresan VEGF berbanding dengan IVF dan SCNT kawalan, kecuali untuk embrio terklon yang dirawat dengan SAH selama 24 jam. Disarankan bahawa selang masa perencatan DNMTs mungkin mempunyai akibat penting terhadap ciri embrio bovin terklon. Kesan peningkatan perencatan DNMTs terhadap kompetensi perkembangan embrio terklon adalah terhad pada jangka masa tertentu sebelum tahap pemetilan *de*

novo. Dengan keputusan ini, penelitian terhadap kesan 0.05 μM TSA, perencat diafetilasi histon, selepas gabungan kuplet selama 12 jam dan pelbagai aspek kompetensi perkembangan pada embrio yang dihasilkan secara SCNT telah dilakukan. Tiada perubahan ($P>0.05$) didapati pada peningkatan belahan, dan kadar blastosista di antara kawalan dan kumpulan yang di rawat.

Walau bagaimanapun, kadar blastosista yang lebih tinggi ($P<0.05$) telah ditunjukkan dalam kumpulan yang dirawat dengan TSA. Kedua-dua penurunan dan peningkatan yang ketara ($P<0.05$) telah di amati dalam analisis pemetilan DNA dan tahap asetilasi histon dalam peringkat blastosista dalam kumpulan yang dirawat adalah mirip dengan blastosista daripada IVF. Akhir sekali, pemeriksaan blastosista pada hari ke 7 mendedahkan pengekspresan OCT4 tertinggi pada kumpulan yang dirawat dengan TSA ($P<0.05$), manakala tiada perubahan ($P>0.05$) didapati dalam pengekspresan BCL2 dan VEGF.

ACKNOWLEDGEMENTS

An effort such as this dissertation is only made possible through the immeasurable support, mentorship, time and patience of many individuals. I am indebted to all who have helped along the way and made the current work a reality.

First and foremost, the author would like to thank Our God for teaching him the way of life. I would like to thank my family for their love and support given to me and most of all their trust that kept me inspired through and my study in oversea.

I wish to convey my heartfelt gratitude to the Malaysian government, Universiti Putra Malaysia, Royan Institute for Reproductive Biomedicine, ACECR, Isfahan, Iran, and my supervisor, Assoc. Professor Dr. Halimatun Yaakub, for their support.

My thanks and appreciations are expressed towards my committee members, especially with patience, encouragement and instruction from Assoc. Professor Dr. Halimatun Yaakub, Assoc. Professor Dr. Nasr-Esfahani Mohammad, Assoc. Professor Dr. ABD. Wahid Haron, Assist. Professor Dr. Noorshariza Nordin and Assist. Professor Dr. Anjomshoa Maryam. I have enhanced my knowledge in both my class work and research.

Additionally, my thanks goes to Professor. Dr. Gourabi Hamid and Assoc. Professor. Dr. Shahverdi Abdolhossein from the Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran, have helped a lot in my research.

The work would not have been made possible without the efforts of Iranian specialist in Iran.

Finally, I am grateful to have found someone to stand beside me and continually remind me not to take life so seriously, especially when an experiment fails. So my final thanks goes to my devoted wife Mary. The only way to thank you would be to reciprocate to always give you the level of love and support that you've given me.

I certify that a Thesis Examination Committee has met on Date of Viva Voce to conduct the final examination of Shahram Jafari on his thesis entitled. "Effects of S-adenosylhomocysteine and Trichostatin A on Developmental Competence, Epigenetic Modification and Gene Expression of Cloned Cattle Embryos" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the degree of Doctor of Philosophy.

Members of the Examination Committee are as follows:

Name of Chairperson, PhD

Associate Professor Dr. Ismail b Idris
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Name of Examiner 1, PhD

Professor Dr. Jothi Malar a/p P.V. Panandam
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Name of Examiner 2, PhD

Associate Professor Dr. Rosnina bt Haji Yusoff
Faculty of Veterinary
Universiti Putra Malaysia
(Member)

Name of External Examiner, PhD

Dr. Bjorn Johannes Obak
Agresearch Limited
Ruakura Research Center
New Zealand
(Independent Examiner)

SEOW HENG FONG, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirements for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Halimatun Bt Yaakub, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Chairman)

Abd. Wahid B Haron, PhD

Associate Professor

Faculty of Veterinary Medicine

Universiti Putra Malaysia

(Member)

Norshariza Binti Nordin, PhD

Assistant Professor

Faculty of Medicine and Health Sciences

Universiti Putra Malaysia

(Member)

Mohammad Hossein Nasr-Esfahani, PhD

Associate Professor

Department of Reproduction and Development,
Cell Sciences Research Center,

Royan Institute for Animal Biotechnology,

ACECR, Isfahan, Iran.

(Member)

Maryam Anjomshoaa, PhD

Assistant Professor

Faculty of Medicine Science

Rafsanjan University of Medical Science,

Rafsanjan, Iran

(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

TABLE OF CONTENTS

	Page
DEDICATION	i
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF ABBREVIATION	xix
 CHAPTER	
1. 1.1 INTRODUCTION	1
1.2 General objectives	1
1.3 Specific objectives	4
1.4 Hypothesis	4
5	5
2. LITERATURE REVIEW	6
2.1 Nuclear transfer	6
2.2 History of nuclear transfer	6
2.2.1 Nuclear transfer in amphibians and reptiles	6
2.2.2 Mammalian nuclear transfer	8
2.3 Importance and application of nuclear transfer	10
2.3.1 Maintain and enhance species on the brink of extinction	10
2.3.2 Production of transgenic animals	11
2.3.3 Generation of stem cells	12
2.3.4 Production and multiplication of animals	13
2.4 Factors affecting SCNT efficiency	13
2.4.1 Source of the donor cells	14
2.4.1.1 Fibroblast cell line	16
2.4.2 Stages of donor cell cycle	17
2.4.3 Method of oocyte enucleation	18
2.4.4 Reconstruction of embryos	18
2.4.5 Embryo culture	19
2.5 Epigenetic modification and nuclear reprogramming	20
2.5.1 Epigenetic modifications: DNA methylation	22
2.5.1.1 DNA methyltransferase	24
2.5.2 Histone modification	24
2.5.3 Nuclear reprogramming in early development of mammalian embryos	26
2.5.4 Nuclear reprogramming in SCNT embryos	28
2.5.5 Nuclear reprogramming approaches	32
2.5.5.1 Cell-fusion based reprogramming	32

2.5.5.2	Cell extracts	33
2.5.5.3	Transfection of pluripotential factors into somatic cells	33
2.5.5.4	Treatment of donor cells or SCNT embryos	35
2.5.5.4.1	Trichostatin A (TSA)	36
2.5.5.4.2	S-adenosyl-L-homocysteine (SAH)	39
3.	EFFECTS OF S-ADENOSYL-L-HOMOCYSTEINE AND TRICHOSTATIN A ON DEVELOPMENTAL COMPETENCY, EPIGENETIC MODIFICATION AND GENE EXPRESSION IN CLONED CATTLE EMBRYOS	
3.1	Introduction	41
3.2	Materials and Methods	44
3.2.1	Preparation of somatic donor cells	44
3.2.1.1	Cell-specific marker staining	45
3.2.2	Donor cells transfection	47
3.2.3	Oocytes collection	48
3.2.4	Oocyte denudation	49
3.2.5	Zona removal	50
3.2.6	Oocyte enucleation	50
3.2.7	Nuclear transfer	51
3.2.8	Electrofusion	52
3.2.9	Reconstructed embryo activation	52
3.2.10	Culture of reconstructed embryos and treatment with SAH	53
3.2.11	<i>In vitro</i> fertilization (IVF)	54
3.2.12	<i>In vitro</i> development of bovine SCNT embryos	55
3.2.13	EGFP- POU5F1 fluorescence	56
3.2.14	Immunofluorescence staining	57
3.2.15	Quantitative analysis of transcripts by real time reverse transcriptase- polymerase chain reaction (RT-PCR)	59
3.2.15.1	RNA extraction	59
3.2.15.2	Reverse transcriptase and cDNA synthesis	60
3.2.15.3	RT-PCR	61
3.2.15.4	Relative quantification	62
3.2.16	Statistical analysis	63
3.3	Results	64
3.3.1	<i>In vitro</i> development of SCNT embryos	64
3.3.2	DNA methylation and histone acetylation levels	66
3.3.3	EGFP- POU5F1 reporter expression	69
3.3.4	Real-time quantification of gene expression	72
3.4	Discussion	73
3.5	Conclusion	80

4.	EFFECTS OF POST ACTIVATION TREATMENT WITH TRICHOSTATIN A ON DEVELOPMENTAL COMPETENCY, EPIGENETIC STATUS AND GENE EXPRESSION OF SOMATIC CELL NUCLEAR DRIVEN BOVINE EMBRYOS	
4.1	Introduction	81
4.2	Materials and Methods	83
4.2.1	Preparation of somatic donor cells, cell-specific marker staining and donor cells transfection	83
4.2.2	Oocytes enucleation and nuclear transfer	83
4.2.3	Electrofusion	83
4.2.4	Reconstructed embryo activation and treatment with TSA	84
4.2.5	Culture of reconstructed embryos	84
4.2.6	<i>In vitro</i> fertilization (IVF)	85
4.2.7	<i>In vitro</i> development of bovine SCNT embryos	85
4.2.8	EGFP- POU5F1 fluorescence	86
4.2.9	Immunofluorescence staining	86
4.2.10	Quantitative analysis of transcripts by real time reverse transcriptase- polymerase chain reaction (RT-PCR)	87
4.2.11	Statistical analysis	87
4.3	Results	88
4.3.1	<i>In vitro</i> development of SCNT embryos	88
4.3.2	EGFP expression of blastocysts on day 7	89
4.3.3	DNA methylation and histone acetylation levels in SCNT embryos	91
4.3.4	Effect of TSA on genes expression of SCNT embryos	95
4.4	Discussion	96
4.5	Conclusion	99
5.	GENERAL DISCUSSION AND RECOMMENDATIONS FOR FUTURE WORK	
5.1	General discussion	100
5.1.1	Effects of SAH	100
5.1.2	Effects of TSA	104
5.2	Recommendation for future work	107
		108
REFERENCES		
APPENDICES		
BIODATA OF STUDENT		
		132
		142