



**UNIVERSITI PUTRA MALAYSIA**

***PRODUCTION OF THERMOSTABLE L2 LIPASE  
USING RECOMBINANT *Pichia pastoris****

**NOOR HIDAYAH SHAHIDAN**

**IB 2012 7**

**PRODUCTION OF THERMOSTABLE L2 LIPASE  
USING RECOMBINANT *Pichia pastoris***

The logo of Universiti Putra Malaysia (UPM) is a shield-shaped emblem. It features a central vertical element resembling a stylized tree or a column with a book at the top. The shield is divided into four quadrants by a vertical and a horizontal line. The top-left quadrant contains the letters 'UPM' in white on a red background. The rest of the shield is filled with a pattern of red and white geometric shapes, including a large 'M' shape in the center. The entire logo is rendered in a light grey, semi-transparent watermark style.

**NOOR HIDAYAH SHAHIDAN**

**MASTER OF SCIENCE  
UNIVERSITI PUTRA MALAYSIA**

**2012**

**PRODUCTION OF THERMOSTABLE L2 LIPASE USING RECOMBINANT  
*Pichia pastoris***

By

**NOOR HIDAYAH SHAHIDAN**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in  
Fulfilment of the Requirements for the Degree of Master of Science**

**May 2012**



*This thesis is especially dedicated to:*

*Fauzi Hanif Hamzah*

*Thank you for being my source of inspirations*

© COPYRIGHT UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**PRODUCTION OF THERMOSTABLE L2 LIPASE USING RECOMBINANT  
*Pichia pastoris***

By

**NOOR HIDAYAH SHAHIDAN**

**May 2012**

**Chairman: Professor Raja Noor Zaliha Raja Abd Rahman, D. Eng**

**Institute: Institute of Bioscience**

Thermostable lipases are widely used for various industrial applications, such as in the production of detergents, oleo chemical industry, food-processing as well in dairy and milk productions. Recombinant *Pichia pastoris* strain GS115 harbouring L2 lipase gene under constitutive expression of glyceraldehyde-3-phosphate (GAP) was subjected to a series of batch cultivation studies at shake-flask scale (100 mL) and in 7.5 L bioreactor. Batch cultivation studies at shake-flask scale were conducted to investigate the effect of different carbon sources on the expression level of L2 lipase and growth of *P. pastoris*. Growth of *P. pastoris* at 2% (w/v) glucose concentration gave the highest lipase activity. Other carbon sources were also tested for growth; oleic acid, trehalose, glycerol, lactose, methanol and sorbitol in which it was observed that 1% (v/v) glycerol produced the highest L2 lipase activity of 34 U/mL. In addition, study on fifth grade molasses as potential carbon source revealed that the highest activity of thermostable L2 lipase of 40 U/mL occurred when the concentration of fifth grade molasses was set at 2% (w/v).

Growth of the recombinant on the remnant of sugar refinery process, mud cake produced little or zero expression level of L2 lipase and poor growth of *P. pastoris*, showing that it was not an effective carbon source. Study conducted on different sugar components of the molasses which were sucrose, maltose, glucose, and fructose showed that the cultivation on fructose led to the highest expression of L2 lipase of 50 U/mL among growth on other carbon sources, followed by growth of *P. pastoris* cells on glucose.

A preliminary study of batch cultivation of recombinant *P. pastoris* was performed using medium containing 2% (w/v) glucose as carbon source in 7.5 L bioreactor with the following parameters: controlled dissolved oxygen tension (DOT) 35%, controlled pH 6.0, 1 vvm, temperature 30°C, 30 h inoculum age, and resulted in 60 U/mL of lipase activity at 24 h of cultivation time. Batch cultivation studies were also conducted using different carbon sources namely glycerol, fifth grade molasses and mixture of glycerol and the fifth grade molasses, which produced 56, 120, and 60 U/mL of lipase activity, respectively. Further optimization studies were conducted for the selected parameters with cultivation medium containing fifth grade molasses as main carbon source: controlled pH at 6.0 and uncontrolled pH (with initial pH at 6.0), inoculum age, controlled DOT, and molasses concentration. Optimized condition with fifth grade molasses as main carbon source produced 120 U/mL when cultivation conditions were set at 35% air saturation, initial pH 6.0, 30 h of inoculum age, and 2% (w/v) of fifth grade molasses concentration. This study showed that the constitutive pGAP expression system of *P. pastoris* is inducible by various carbon sources; therefore, it offers flexibility for diverse expressions of heterologous proteins.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGHASILAN LIPASE L2 TERMOSTABIL MENGGUNAKAN  
REKOMBINAN *Pichia pastoris***

Oleh

**NOOR HIDAYAH SHAHIDAN**

Mei 2012

**Pengerusi: Profesor Raja Noor Zaliha Raja Abd Rahman, D. Eng**

**Institut: Institut Biosains**

Lipase termostabil telah digunakan secara meluas untuk pelbagai kegunaan industri seperti pembuatan detergen, industri oleo kimia, pemprosesan makanan serta untuk penghasilan tenusu dan susu. Rekombinan *Pichia pastoris* strain GS115 yang mengandungi gen L2 lipase di bawah kawalan promoter konstitutif gliseraldehida-3-fosfat dehydrogenase (GAP) adalah tertakluk kepada satu siri kajian pertumbuhan sesekelompok pada skala kelalang goncang (100 mL) dan di dalam 7.5 L bioreaktor. Kajian pembiakan berkumpulan pada skala kelalang goncang telah dijalankan untuk mengkaji kesan pelbagai sumber karbon ke atas penghasilan L2 lipase dan pertumbuhan *P. pastoris*. Pertumbuhan pada kepekatan glukosa 2% (w/v) menghasilkan aktiviti lipase tertinggi. Sumber karbon lain yang dikaji termasuk asid oleik, trehalose, gliserol, laktosa, metanol dan sorbitol, di mana bagi sumber karbon lain, gliserol menghasilkan aktiviti L2 lipase tertinggi iaitu 34 U/mL bagi rekombinan *P. pastoris*. Di samping itu, kajian mengenai sirap pekat (*molasses*) gred kelima sebagai potensi sumber karbon

menunjukkan bahawa aktiviti tertinggi L2 lipase iaitu 40 U/mL berlaku pada kepekatan sirap pekat 2% (b/i) gred kelima. Pertumbuhan rekombinan menggunakan sisa proses kilang penapisan gula, kek lumpur (*mud cake*) menghasilkan aktiviti L2 lipase yang rendah serta kadar pertumbuhan *P. pastoris* yang rendah, menunjukkan ia bukan sumber karbon yang efektif. Kajian yang dijalankan ke atas komponen gula yang berbeza terkandung dalam sirap pekat iaitu sukrosa, maltosa, glukosa, dan fruktosa menunjukkan bahawa pembiakan yis di dalam medium mengandungi fruktosa menghasilkan kadar tertinggi L2 lipase iaitu 50 U/mL berbanding sumber-sumber karbon yang lain, diikuti oleh pertumbuhan sel-sel *P. pastoris* di dalam glukosa.

Kajian awal pembiakan berkumpulan rekombinan *P. pastoris* dijalankan menggunakan medium yang mengandungi 2% (b/i) glukosa sebagai sumber karbon dengan parameter berikut: ketegangan terkawal oksigen terlarut (DOT) 35%, pH terkawal 6.0, 1 vvm, suhu 30°C, 30 jam umur inoculum didapati menghasilkan kadar aktiviti L2 lipase sebanyak 60 U/mL pada 24 jam masa pertumbuhan. Kajian pertumbuhan sesekelompok juga dijalankan ke atas sumber karbon yang berbeza iaitu gliserol, sirap pekat gred kelima dan campuran gliserol dan sirap pekat gred kelima, yang masing-masing menghasilkan 56, 120 dan 60 U/mL lipase aktiviti. Kajian pengoptimuman seterusnya adalah menggunakan sirap pekat gred kelima sebagai sumber karbon utama: pH 6.0 terkawal dan pH yang tidak terkawal (dengan pH awal pada kadar 6.0), umur inoculum, DOT, dan kepekatan sirap pekat. Keadaan optimum bagi sirap pekat gred kelima sebagai sumber karbon utama menghasilkan 120 U/mL apabila parameter telah ditetapkan pada 35% keterlarutan oksigen, pH awal 6.0, 30 jam umur inoculum dan 2% (b/i) kepekatan sirap pekat gred kelima. Kajian ini telah menunjukkan bahawa sistem ekspresi pGAP *P.*



*pastoris* boleh diaruh oleh pelbagai sumber karbon, menjadikannya fleksibel untuk ekspresi protein asing dengan lebih meluas.



## ACKNOWLEDGEMENTS

I wish to express my sincere appreciation to my supervisor, Prof. Dr. Raja Noor Zaliha Raja Abd Rahman, for her valuable support and guidance during my Master study. I would also like to extend my appreciations to my co-supervisors: Prof. Dr. Abu Bakar Salleh, Assoc. Prof. Dr. Rosfarizan Mohamed, and Dr. Adam Leow for their valuable advices regarding my experimental work and this thesis. Appreciations are also extended to my fellow colleagues from Enzyme and Microbial Technology Research (EMTECH) group for the assistance and encouragement I received throughout my study.

I would also like to thank Mr. Zainal Rahmat for the initial idea of using fifth grade molasses as carbon source for yeast cultivation, and to the Central Sugar Refinery Sdn. Bhd. (Malaysia) for the supply of molasses. I feel indebted to Siti Nurbaya Oslan for technical advices. Special thanks to my best buddy, Izuddin Abdul Rahman, for making my life in UPM even more memorable. To Fairolniza Shariff, thank you for a sincere friendship. To my parents and siblings, I am grateful for the continuous support. Last but not least, I would like to thank everyone whom has involved either directly or indirectly throughout the journey of completing my degree.

This study has been financially aided by National Science Fellowship (NSF) from the Ministry of Science, Technology, and Innovation of Malaysia.

Ultimately, it is Allah who has given me this life and driven me to success. All praises are due to Him. *Alhamdulillah.*

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Raja Noor Zaliha Raja Abd Rahman, D. Eng**

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

**Abu Bakar Salleh, PhD**

Professor Dato'

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

**Rosfarizan Mohamed, PhD**

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

**Adam Leow Thean Chor, PhD**

Senior Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

---

**BUJANG BIN KIM HUAT, PhD**

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

## DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

---

**NOOR HIDAYAH SHAHIDAN**

Date: 18 May 2012

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	iii
<b>ABSTRAK</b>	v
<b>ACKNOWLEDGEMENTS</b>	viii
<b>APPROVAL</b>	ix
<b>DECLARATION</b>	x
<b>LIST OF TABLES</b>	xiv
<b>LIST OF FIGURES</b>	xv
<b>LIST OF ABBREVIATIONS</b>	xvii
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	<b>4</b>
2.1 Lipases	4
2.2 Thermostable lipases	7
2.3 Thermostable L2 lipase	8
2.4 Metabolisms of <i>Pichia pastoris</i>	10
2.5 <i>P. pastoris</i> expression system	14
2.6 Production of heterologous protein using <i>P. pastoris</i> as expression host	16
2.6.1 Expression of heterologous proteins using AOX promoters	18
2.6.2 Expression of heterologous proteins through AOX inducible system using mixture of carbon sources	19
2.6.3 Expression of heterologous proteins using GAP promoter	21
2.6.4 Expression of lipase genes in <i>P. pastoris</i>	23
2.7 Mode of bioreactor operations for <i>P. pastoris</i> cultivations	24
2.7.1 Batch culture	24
2.7.2 Fed-batch culture	25
2.7.3 Continuous culture	27
2.8 Effects of physical factors on heterologous protein production by recombinant <i>P. pastoris</i>	28
2.8.1 Temperature	28
2.8.2 Culture pH	31
2.8.3 Dissolved oxygen tension (DOT)	32
2.9 Effects of nutritional factors on heterologous protein production by recombinant <i>P. pastoris</i>	34
2.9.1 Carbon source	34
2.9.1.1 Alternative carbon sources	36
2.9.1.1.1 Molasses	36
2.9.1.1.2 Mud cake	39
2.9.2 Nitrogen source	40
2.9.3 Other potential supplements	41

2.10	Concluding remarks	42
<b>3</b>	<b>MATERIALS AND METHODS</b>	<b>43</b>
3.1	Materials	43
3.2	Overall view of experimental plan	44
3.3	Microorganism	45
3.4	Expression of L2 lipase in <i>P. pastoris</i>	45
	3.4.1 Transformation of recombinant plasmid into <i>P. pastoris</i>	45
	3.4.2 Direct screening of multicopy transformants	47
	3.4.3 Direct PCR analysis of transformants	47
3.5	Gene expression in <i>P. pastoris</i>	48
	3.5.1 Constitutive expression of pGAPZ $\alpha$ A/L2	48
	3.5.1.1 Screening for lipase activity of positive transformants	48
3.6	Batch cultivation studies of recombinant <i>P. pastoris</i> in shake flasks	49
	3.6.1 Effect of various carbon sources for <i>P. pastoris</i> growth and L2 lipase expression	49
	3.6.1.1 Characterization of glucose concentration	49
	3.6.1.2 Characterization of other carbon sources	50
	3.6.2 Effect of fifth grade molasses concentrations as potential carbon source	50
	3.6.3 Effects of sugar components of molasses on <i>P. pastoris</i> growth and L2 lipase expression	51
	3.6.4 Study on mud cake as potential carbon source for <i>P. pastoris</i> growth and L2 lipase expression	51
3.7	Batch cultivation studies of recombinant <i>P. pastoris</i> in bioreactor	52
	3.7.1 Preparation of seed culture	52
	3.7.2 Bioreactor	52
	3.7.3 Effect of media on recombinant <i>P. pastoris</i>	56
	3.7.3.1 Cultivation on 2% glucose	56
	3.7.3.2 Cultivation on 2% molasses	57
	3.7.3.3 Cultivation on 2% glycerol	57
	3.7.3.4 Cultivation on 2% glycerol and 2% molasses	58
	3.7.4 Optimization studies of recombinant <i>P. pastoris</i> with fifth grade molasses	58
	3.7.5.1 Effect of pH	58
	3.7.5.2 Effect of dissolved oxygen tension (DOT)	58
	3.7.5.3 Effect of inoculum age	59
	3.7.5.4 Effect of molasses concentration	59
3.8	Analytical procedures	60
	3.8.1 Cell density measurement	60
	3.8.2 Lipase assay	60
	3.8.3 Determination of protein concentration	61
	3.8.4 Determination of glucose concentration	61
3.9	Statistical analysis	62

<b>4</b>	<b>RESULTS AND DISCUSSION</b>	<b>63</b>
4.1	Transformation and selection of recombinant <i>P. pastoris</i>	63
4.2	Expression of thermostable L2 lipase in <i>P. pastoris</i> under constitutive glyceraldehyde-3-phosphate dehydrogenase (GAP) promoter	66
4.2.1	Screening for L2 lipase activity	67
4.3	Batch cultivation studies of constitutive lipase expression in shake flasks	69
4.3.1	Effect of carbon source	69
4.3.1.1	Effect of glucose	69
4.3.1.2	Effect of other carbon sources	72
4.4	Use of fifth grade molasses as potential carbon source	77
4.4.1	Analysis of molasses: physical characteristics and sugar content	77
4.4.2	Effect of fifth grade molasses on <i>P. pastoris</i> growth and L2 lipase expression	78
4.4.3	Effect of sugar components of molasses on <i>P. pastoris</i> growth and L2 lipase expression	82
4.5	Use of mud cake as potential carbon source	85
4.6	Cultivation studies of <i>P. pastoris</i> expressing L2 lipase in bioreactor	87
4.6.1	Effect of preliminary bioreactor culturing condition	88
4.6.2	Batch cultivation studies of recombinant <i>P. pastoris</i> in bioreactor using molasses as carbon source	91
4.6.2.1	Effect of pH	92
4.6.2.2	Effect of inoculum age	96
4.6.2.3	Effect of dissolved oxygen tension (DOT)	100
4.6.2.4	Effect of molasses concentration	104
4.6.3	Time course study using optimized parameters	108
4.6.4	Effect of different media	111
<b>5</b>	<b>CONCLUSIONS AND RECOMMENDATIONS</b>	<b>117</b>
	<b>REFERENCES</b>	<b>120</b>
	<b>APPENDICES</b>	<b>134</b>
	<b>BIODATA OF STUDENT</b>	<b>153</b>
	<b>LIST OF PUBLICATIONS</b>	<b>154</b>