



UNIVERSITI PUTRA MALAYSIA

**EFFECTS OF SELECTED PROBIOTICS ON THE GROWTH, SURVIVAL AND
GASTRO-INTESTINAL BACTERIAL POPULATION OF JUVENILE GIANT
FRESHWATER PRAWN (*Macrobrachium rosenbergii*, de Man)**

SARA RAVAN RAMZANI

FP 2011 55

**EFFECTS OF SELECTED PROBIOTICS ON THE GROWTH, SURVIVAL AND
GASTRO-INTESTINAL BACTERIAL POPULATION OF JUVENILE GIANT
FRESHWATER PRAWN (*Macrobrachium rosenbergii*, de Man)**



**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfillment of the Requirements for the Master of Science**

December 2011

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment
of the requirements for degree of Masters of Science

**EFFECTS OF SELECTED PROBIOTICS ON THE GROWTH, SURVIVAL AND
GASTRO-INTESTINAL BACTERIAL POPULATION OF JUVENILE GIANT
FRESHWATER PRAWN (*Macrobrachium rosenbergii*, de Man)**

By

SARA RAVAN RAMZANI

December 2011

Chairman: Associate Professor Che Roos Bin Saad, PhD

Faculty: Agriculture

Two experiments were carried out in this project. The first study was done to determine the effectiveness of putative bacteria as a probiotics for growth enhancement and survival of juvenile *Macrobrachium rosenbergii* and protection against *Aeromonas hydrophila* challenge. A total of 360 juveniles *M. rosenbergii* were individually weighed and kept in twenty one 60 L aquaria. Four treatments were conducted as follows; T₁ (Control group), T₂ (*B. subtilis*), T₃ (*Lactobacillus acidophilus*) and T₄ (Commercial probiotic bacteria). The final concentration of the probiotic bacteria in the treated feeds was set at 8×10^9 CFU/g feed. All prawn treated with probiotic-supplemented diets showed better growth performances and feed utilization as compared to those in the control fed with basal diet. The higher growth parameter and survival rate of the prawns fed with *B. subtilis*-added diets over the control feed indicated the probiotic-ability of the *B. subtilis*. There was a significant different (P<0.05) in weight gain, specific growth

rate, feed intake and FCR between the treated and control tanks. The prawns fed with the control diet with no addition of bacteria, recorded the lowest feed intake. The best FCR value (2.56 ± 0.09) was derived from the juvenile fed diet containing *B. subtilis*. After 60 days of comparison among putative *B. subtilis*, *Lactobacillus acidophilus*, commercial probiotic bacteria and probiotic free diets in juvenile *M. rosenbergii*, there was no significant difference ($P > 0.05$) in biochemical composition between groups treated putative *B. subtilis* and *Lactobacillus acidophilus* and commercial probiotic treated groups. There was significant difference between groups receiving putative *B. subtilis*, *Lactobacillus acidophilus* and commercial probiotic bacteria diets in growth enhancement of juvenile *M. rosenbergii*. This study indicated the increase in the growth and survival of the treated prawns, with reduced mortality as compared to the control group.

The second study was conducted to determine the effects of periodical application of probiotics and pathogenic effects of *A. hydrophila* on the survival and growth enhancement of juvenile *M. rosenbergii* with a total of 840 *M. rosenbergii* post-larvae procured from Center of Marine Science (COMAS), Port Dickson, UPM. These prawns were individually weighed and kept in twenty one 60 L aquaria. Two trials were carried out. The first trial had seven feeding regimes designed as: T₁ (Control group); T₂ (*B. subtilis* bacteria once every day), T₃ (*B. subtilis* once every 5 day), T₄ (*B. subtilis* once every 10 days), T₅ (Commercial Probiotic (CP) once every day), T₆ (CP once every 5 days) and T₇ (CP once every 10 days). The final concentration of the probiotic bacteria in the feeds was determined at 8×10^9 CFU/g feed. After 60 days of feeding, the prawns in each treatment were measured for growth performance. Then the prawns in each

treatment were challenged with *A. hydrophila* at 10^7 CFU/mL for 21 days, and measurements for growth and survival rates were recorded at end of the challenge experiment. Parameters such as histopathology, bacteriological study, growth measurement and water quality were taken. Significant differences ($P < 0.05$) in weight gain, specific growth rate, feed intake and FCR between the treated *B. subtilis* and control groups were observed. Results showed a significant difference between control and treatments in the log of *Bacillus* count (Control: Mean = 1.40 ± 0.03 , CPF; Mean= 4.10 ± 0.064 , PF; Mean = 3.46 ± 0.084) and the log of gram negative count (Control; Mean= 3.29 ± 0.08 , CPF; Mean = 1.91 ± 0.045 , PF; Mean = 2.22 ± 0.059) ($P < 0.001$). Appropriate probiotic applications have been shown to improve intestinal microbial balance, thus leading to improved food digestion, and reduced pathogenic bacteria multiplication problems in the gastrointestinal tract.

Abstrak tesis dipersembahkan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains.

KESAN PROBIOTIK TERPILIH KE ATAS PERTUMBUHAN, KEMANDIRIAN DAN POPULASI BAKTERIA GASTROINTESTIN BAGI JUVANA UDANG GALAH (*Macrobrachium rosenbergii*, de Man)

Oleh

SARA RAVAN RAMZANI

Disember 2011

Pengerusi : Prof. Madya Che Roos Bin Saad, PhD

Fakulti : Pertanian

Dua kajian telah dilakukan dalam projek ini. Kajian pertama telah dijalankan untuk menentukan keberkesanan bakteria yang dianggap sebagai probiotik bagi meningkatkan pertumbuhan dan kemandirian juvana *M. rosenbergii* dan perlindungan daripada cabaran bakterium *Aeromonas hydrophila*. Sejumlah 360 juvana *M. rosenbergii* telah ditimbang secara berasingan dan dimasukkan ke dalam 21 tangki akuarium berkapasiti 60L. Empat rawatan telah dijalankan seperti berikut; T1 (Tiada penambahan *B. subtilis*), T2 (Penambahan *B. subtilis*), T3 (*Lactobacillus acidophilus*), dan T4 (Komersial bakteria probiotik). Kepekatan akhir bakteria probiotik dalam makanan yang dirawat telah ditetapkan pada tahap 8×10^9 CFU/g makanan. Kesemua udang yang mendapat probiotik tambahan menunjukkan prestasi pertumbuhan dan penggunaan makanan yang lebih baik berbanding udang yang memakan diet kawalan. Terdapat perbezaan yang bererti ($P<0.05$) untuk kadar pertumbuhan dan kemandirian hidup yang lebih tinggi bagi

udang yang diberi makan dengan diet yang ditambah *B. subtilis* berbanding dengan udang kawalan. Ini menunjukkan bahawa keupayaan probiotik dimiliki oleh *B. subtilis*. Nilai FCR terbaik (2.56 ± 0.09) diperolehi dari pada juvana dengan diet yang mengandungi *B. subtilis*. Selepas perbandingan selama 60 hari di antara *B. subtilis*, *Lactobacillus acidophilus*, probiotik komersial dan diet tanpa probiotik dengan juvana *M. rosenbergii*, tiada perbezaan ketara antara diet makanan tersebut. Namun, terdapat perbezaan ketara antara *B. subtilis*, *Lactobacillus acidophilus* dan probitik komersial yang mempengaruhi pertumbuhan juvana *M. rosenbergii*. Kajian ini menunjukkan peningkatan pertumbuhan dan kemandiriaan udang yang dirawat, dengan pengurangan dalam kadar kematian berbanding kumpulan kawalan.

Kajian kedua telah dijalankan bagi menentukan kesan patogenik oleh *A. hydrophila* diatas pemberian probiotik secara berkala dengan menggunakan 840 ekor pasca larva *M. rosenbergii* yang diperolehi dari Pusat Sains Marin, UPM Port Dickson (COMAS). Kesemua udang telah ditimbang secara berasingan dan diagihkan kedalam 21 tangki akuarium berkapasiti 60 L. Dua percubaan telah dijalankan. Percubaan pertama telah direka bentuk dengan tujuh rejim pemberian makanan: T1 (Kawalan), T2 (*B. subtilis* setiap hari sekali), T3 (*B. subtilis* setiap 5 hari sekali), T4 (*B. subtilis* setiap 10 hari sekali), T5 (Commercial Probiotik (CP) setiap hari sekali), T6 (CP setiap 5 hari sekali), dan T7 (CP setiap 10 hari sekali). Kepekatan akhir bakteria probiotik di dalam makanan telah ditentukan pada tahap 8×10^9 CFU/g makanan. Selepas 60 hari pemberian makanan mengikut setiap rawatan, kesemua udang tersebut telah diukur bagi prestasi pertumbuhan. Kemudian semua udang telah diuji ketahanannya dengan *A. hydrophila* pada tahap 10^7 CFU/mL selama 21 hari dan ukuran berangka untuk kadar pertumbuhan

dan kemandirian setiap rawatan ditentukan pada akhir kajian ini. Parameter seperti histopatologi, kajian bakteriologi, pengukuran pertumbuhan, dan kualiti air telah diambil. Perbezaan bererti ($P < 0.05$) dalam pertambahan berat badan, kadar pertumbuhan spesifik, pengambilan makanan, dan nisbah pertukaran makanan (FCR) di antara kumpulan yang dirawat *B. subtilis* dengan kumpulan kawalan telah diamati. Hasil kajian menunjukkan terdapat perbezaan bererti antara kumpulan kawalan dan rawatan bagi kiraan log *Bacillus* (Kawalan: Min = 1.40 ± 0.03 , CPF; Min= 4.10 ± 0.064 , PF; Min = 3.46 ± 0.084) dan kiraan log gram negatif (Kawalan: Min= 3.29 ± 0.08 , CPF; Min = 1.91 ± 0.045 , PF; Min = 2.22 ± 0.059) ($P < 0.001$). Aplikasi probiotik yang sesuai menunjukkan peningkatan keseimbangan mikrob usus, sekali gus membawa kepada pencernaan makanan yang lebih baik dan mengurangkan masalah pembiakan bakteria patogen di dalam saluran gastrousus.

ACKNOWLEDGEMENTS

In the name of Allah, The Beneficent, The Merciful

First and most important, I would like to offer my earnest appreciation to the Assoc. Professor Dr. Che Roos Saad, chairman of my advisory committee, for providing me with a great opportunity to finish my Master studies under his excellent supervision. Apart from the provision of the required funding, this work would not have been possible without his patience, never-ending inspiration, advice and knowledge. On account of recurrent meeting and his open door policy, Assoc. Prof Dr. Che Roos Saad made a great contribution to this dissertation and my educational development, together with my professional and individual life.

My deepest gratitude is also extended to Assoc. Prof Dr. Kamaruzaman Sijam who is a member of my supervisory committee and provided me some equipment for laboratory and assisted throughout the course of the study. Furthermore, I would like to extend my most heartfelt gratefulness and thanks to Assoc. Prof Dr. Hassan Mohd Daud, for helpful advice and guidance during the study time. I am also thankful for their desire to serve on my committee, provide me support whenever essential, participation in my qualifying examination, and for assessing this dissertation.

I certify that an Examination Committee met on 16 December 2011 of viva to conduct the final examination of Sara Ravan Ramzani on his Master of Science thesis entitled “Effects of Selected Probiotics on the Growth, Survival and Gastrno-Intestinal Bacterial population of Juvenile Giant Freshwater Prawn (*Macrobrachium rosenbergii*)” in accordance with University Colleges Act 1971 and the constitution of the Universiti Putra Malaysia [P.U.(A) 106 15 March 1998. The Committee recommends that the candidate be awarded the Masters of Science.

Member of the Thesis Examination Committee are as follows:

Dr Aziz Arshad, PhD

Professor

Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Mohd Salleh Kamarudin, PhD

Associate Professor

Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Siti Shapor Siraj, PhD

Professor

Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Ng Wing Keong, PhD

Professor

School of Biological Sciences
Universiti Sains Malaysia
(Independent Examiner)

Seow Heng Fong, PhD

Professor

and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia
Date

This thesis was submitted to the Senate of University Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Che Roos Saad, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Chairman)

Hassan Mohd Daud, PhD

Associate Professor

Faculty of Veterinary Medicine

Universiti Putra Malaysia

(Member)

Kamaruzaman Sijam, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean,

School of Graduate Studies

Universiti Putra Malaysia

Date:

DECLARATION

I declare that this thesis is my original work except for quotations and citation except as cited in references, which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or other institutions.

SARA RAVAN RAMZANI

Date: 16 December 2011



TABLE OF CONTENTS

	Page
DEDICATION	ii
ACKNOWLEDGEMENTS	iii
APPROVAL	iv
ABSTRACT	v
ABSTRAK	vii
DECLARATION	ix
LIST OF TABLES	x
LIST OF FIGURES	xi
 CHAPTER	
 I INTRODUCTION	
1.1 Background of Study	1
1.2 Problem Statement	1
1.3 Significance of the Study	4
1.4 Objectives	5
6	
 2 LITERATURE REVIEW	
2.1 General Morphology	7
2.1.1 Taxonomy and Life history of <i>M. rosenbergii</i>	7
2.1.2 Prawn species	8
2.1.3 Feeding behavior and Environmental factors of <i>M. rosenbergii</i>	9
11	
2.1.4 Growth	15
2.1.5 Breeding	15
2.1.6 Seed Supply	17
2.2 Larval Culture	17
2.3 Nursery phase	18
2.3.1 Effect of stocking different Size- Grade of Juvenile Prawns	20
20	
2.4 Stocking of Juveniles	20
2.5 General Review on Probiotics	21
21	
2.5.1 Definition of Probiotics	21
2.5.2 Studies on Probiotic Quality control in Aquaculture	23
23	
2.5.3 Modes of Action and Competitive Exclusion of Probiotics	24
24	
2.5.4 Digestion Enhancement of Probiotics	26
2.5.4.1 Immune Response Enhancement	27
27	

2.5.4.2 Water Quality Improvement	27
2.5.4.3 Colonization	29
3 GENERAL METHODOLOGY	31
3.1 Location of the Study	31
3.2 Experimental Set-up	31
3.3 Bacterial Enumeration .	32
3.4 Growth and Feeding Efficiency	33
3.5 Proximate analysis	33
3.5.1 Crude Protein	34
3.5.2 Lipids	34
3.5.3 Crude Fiber	35
3.5.4 Moisture Determination	36
3.5.5 Ash Determination	37
3.6 Water Quality Management	37
3.7 Staining Method	38
3.8 Histopathology	38
3.8.1 Hematoxylin & Eosin staining	39
3.9 Statistical Analysis	39
4 EFFICIENCY OF <i>Bacillus subtilis</i>, <i>Lactobacillus acidophilus</i> AND COMMERCIAL PROBIOTIC BACTERIA ON THE GROWTH AND SURVIVAL OF JUVENILE <i>Macrobrachium rosenbergii</i>	40
4.1 Introduction	40
4.2 Material and Methods	42
4.2.1 Bacterial strain	43
4.2.2 The preparation of the diets	44
4.2.3 Feeding	45
4.2.4 Growth and Feeding Efficiency	45
4.2.5 Proximate Analyses	46
4.2.6 Water Quality Parameters	47
4.2.7 Bacteriological Study	47
4.2.8 Statistical Analysis	47
4.3 Results	48
4.4 Discussion	54

5 RESPONSE OF JUVENILE <i>Macrobrachiall rosenbergii</i> TO DIFFERENT DIETARY LEVELS OF PROBIOTICS IN FEED AND CHALLENGE TEST WITH <i>Aerobacillus hydropltila</i>	58
5.1 Introduction	58
5.2 Material and Methods	60
5.2.1 Bacterial strain	61
5.2.2 The preparation of the Diets	62
5.2.3 Feeding	63
5.2.4 Pathogen Challenge Trial	64
5.2.5 Bacteriological Study	64
5.2.6 Histopathological examination	65
5.2.7 Growth Measurement	65
5.2.8 Water Quality Parameters	65
5.2.9 Statistical Analysis	66
5.3 Results	66
5.4 Discussion	80
6 GENERAL DISCUSSION CONCLUSION AND RECOMMENDATIONS	83
REFERENCES	86
APPENDICES	106
BIODATA OF STUDENT	116